



**CNX Three Mile Hill
Coolgardie Gold Project**

Biological Surveys

**Prepared for
Focus Minerals Limited**

June 2022

● people ● planet ● professional

Document Reference	Revision	Prepared by	Reviewed by	Admin Review	Submitted to Client	
					Copies	Date
4794AA_Rev0	Internal Draft	LC, JW	SC, EW, ML	LI	-	28/04/2022
4794AA_Rev1	Client Draft	360 Environmental	Focus Minerals Limited	-	1 electronic	02/05/2022
4794AA_Rev2	Final	360 Environmental	Focus Minerals Limited	LI	1 electronic	20/06/2022

Disclaimer

This report is issued in accordance with, and is subject to, the terms of the contract between the Client and 360 Environmental Pty Ltd, including, without limitation, the agreed scope of the report. To the extent permitted by law, 360 Environmental Pty Ltd shall not be liable in contract, tort (including, without limitation, negligence) or otherwise for any use of, or reliance on, parts of this report without taking into account the report in its entirety and all previous and subsequent reports. 360 Environmental Pty Ltd considers the contents of this report to be current as at the date it was produced. This report, including each opinion, conclusion, and recommendation it contains, should be considered in the context of the report as a whole. The opinions, conclusions and recommendations in this report are limited by its agreed scope. More extensive, or different, investigation, sampling and testing may have produced different results and therefore different opinions, conclusions, and recommendations. Subject to the terms of the contract between the Client and 360 Environmental Pty Ltd, copying, reproducing, disclosing, or disseminating parts of this report is prohibited (except to the extent required by law) unless the report is produced in its entirety including this cover page, without the prior written consent of 360 Environmental Pty Ltd.

© Copyright 2022 360 Environmental Pty Ltd ACN 109 499 041

Executive Summary

Focused Mineral Limited commissioned 360 Environmental (part of SLR Consulting) to undertake a detailed flora and vegetation survey and basic vertebrate fauna and habitat survey at selected sites within the Three Mile Mine Project, Coolgardie. The Survey Area consists of four defined survey areas adjacent to the townsite of Coolgardie, Western Australia, and covers approximately 1360.72 ha.

The purpose of the assessment was to identify key biological values to inform decisions regarding the specific lease areas for proposed construction sites within the Survey Area.

Flora and Vegetation

The flora desktop assessment identified 90 conservation significant taxa occurring within 50 km of the Survey Area, including three Threatened taxa (*Gastrolobium graniticum*, *Tetratheca spenceri* and *Thelymitra stellata*). A pre-survey likelihood of occurrence assessment was undertaken and determined 20 taxa as having a high likelihood of occurrence, 18 taxa as having a medium likelihood of occurrence, and 51 taxa as having a low likelihood of occurrence. Following the survey, two taxa retained a high likelihood of occurrence.

The detailed flora and vegetation survey was undertaken in between the 11 - 15 October 2021 (Trip 1) and 15 - 19 November 2021 (Trip 2), with the survey recording the floristic composition and vegetation types from 74 quadrats, 22 relevés and 91 mapping notes. A total of 149 taxa were recorded from 78 genera across 35 families.

No Threatened flora taxa pursuant to the Environment Protection and Biodiversity Conservation Act 1999 and/or gazetted as Threatened pursuant to the Biodiversity and Conservation Act 2016 were recorded during the survey.

One Priority listed flora taxon was identified; a single population of *Austrostipa blackii* (P3) consisting of approximately 10 individuals was recorded within a single quadrat. The vegetation it was recorded in was heavily surveyed and no further populations were recorded.

Thirteen introduced taxa were recorded in the Survey Area, of which one, the Common Prickly Pear (**Opuntia stricta*) is listed as a Declared Pest, and a Weed of National Significance.

Ten vegetation types were described and mapped within the Survey Area. Vegetation in the Survey Area was representative of existing broad scale vegetation, as a well as soil and land system mapping for the area.

Vegetation condition ranged from Excellent to Completely Degraded with the majority considered to be in Excellent condition. Evidence of disturbance across the Survey Area included mine works, cattle grazing, vehicle tracks, weeds, and litter.

Vertebrate Fauna

The fauna desktop assessment identified 20 conservation significant species occurring within 50 km of the Coolgardie Survey Area comprising 17 bird species and 3 mammal species.

Fauna habitat mapping was undertaken based on a combination of aerial imagery, field observations and fauna habitat assessment data. Five fauna habitats were mapped within the Survey Area, of which the *Eucalyptus* Woodland and the *Acacia* Shrubland habitat types represent the most value to conservation significant fauna and overall fauna assemblage.

The basic terrestrial vertebrate fauna survey was undertaken in between the 11 – 15 October 2021 (Trip 1) and 15 – 19 November 2021 (Trip 2). This is considered suitable timing to survey for birds, mammals, reptiles, and summer breeding amphibians, but outside the optimal survey period for autumn and winter breeding amphibians. A total of 61 fauna taxa from 33 families comprising of: 42 birds from 21 families, ten mammals from seven families and nine reptiles from five families. No amphibian taxa were recorded.

One conservation significant taxon, the Western Quoll, Chuditch (*Dasyurus geoffroii fortis*) which is listed as Vulnerable under the Biodiversity Conservation Act and Environment Protection and Biodiversity Conservation Act, was recorded within the Survey Area. Chuditch scat was observed during the field survey.

Seven introduced mammal taxa were recorded during the survey, Goat (*Capra hircus*), European Cattle (*Bos primigenius taurus*), Dog/Dingo (*Canis familiaris*), Red Fox (*Vulpes vulpes*), Horse (*Equus caballus*), Cat (*Felis catus*) and Rabbit (*Oryctolagus cuniculus*).

Although outside the scope of this report, potential habitat for Arid Bronze Azure Butterfly (*Ogyris subterrestris petrina*), which is listed as Critically Endangered under the Biodiversity Conservation Act and Environment Protection and Biodiversity Conservation Act, and the Inland Hairstreak butterfly (*Jalmenus aridus*), which is listed as Priority 1 by the Department of Biodiversity, Conservation and Attractions, was observed within the Survey Area.

Abbreviations

Abbreviations used through the report are described below in Table 1.

Table 1: Abbreviations

Abbreviation	Description
BAM Act	Biosecurity and Agriculture Management Act 2007
BC Act	Biodiversity Conservation Act 2016
BoM	Bureau of Meteorology
CD	Conservation Dependent Fauna
CR	Critically Endangered
DAWE	Department of Agriculture, Water, and the Environment
DBCA	Department of Biodiversity, Conservation and Attractions
DoE	Department of Environment
DP	Declared Pest
DWER	Department of Water and Environmental Regulation
EIA	Environmental Impact Assessment
EN	Endangered
EP Act	Environmental Protection Act 1986
EPA	Environmental Protection Authority
EPBC Act	Environment Protection Biodiversity and Conservation Act 1999
ESA	Environmentally Sensitive Area
FML	Focus Minerals Limited
GDE	Groundwater Dependent Ecosystem
IBRA	Interim Biogeographic Regionalisation for Australia
IBSA	Index of Biodiversity Surveys for Assessments
MI	Migratory
MNES	Matters of National Environmental Significance
NVIS	National Vegetation Information System
OS	Other Specially Protected Fauna
P	Priority
PEC	Priority Ecological Community
PMST	Protected Matters Search Tool
T	Threatened
TEC	Threatened Ecological Community
TPFL	Threatened and Priority Flora Database
VU	Vulnerable
WAH	Western Australian Herbarium

Abbreviation	Description
WAM	Western Australian Museum
WoNS	Weeds of National Significance

Table of Contents

1	Introduction.....	1
1.1	The Project.....	1
1.2	Objectives and Scope	1
2	Background.....	2
2.1	Protection of Flora, Vegetation and Fauna	2
2.2	Existing Environment	3
3	Methods.....	7
3.1	Desktop Assessment	7
3.2	Field Survey Effort	9
3.3	Flora and Vegetation.....	10
3.4	Vertebrate Fauna	13
4	Results.....	14
4.1	Limitations	14
4.2	Flora and Vegetation.....	16
4.3	Vertebrate Fauna	28
5	Discussion	45
5.1	Flora and Vegetation.....	45
5.2	Vertebrate Fauna	47
5.3	Invertebrates	51
6	Conclusion	53
7	Report Disclaimer	54
8	References	55

List of Tables

Table 1: Abbreviations.....	iii
Table 2: Broad Vegetation Types within the Survey Area and their Representation at the State, Regional and Local Levels (Government of Western Australia, 2019).....	5
Table 3 Database Searches of the Survey Area	8
Table 4 Likelihood of Occurrence Criteria.....	9
Table 5: Limitations and Constraints Associated with the Survey	14
Table 6: Introduced Flora Species within the Survey Area.....	18
Table 7 Area (ha) covered by each vegetation condition category within the Survey Area	19
Table 8: Vegetation Types Occurring within the Survey Area.....	21
Table 9: Species Richness Indicators	28
Table 10: Fauna Habitat Types within the Survey Area	29
Table 11: Overview of Vertebrate Fauna Species Recorded	34
Table 12: Conservation Significant Fauna Likelihood of Occurrence	37

List of Plates

Plate 1: <i>Austrostipa blackii</i> (P3).....	18
Plate 2: Scat of Western Quoll, Chuditch (<i>Dasyurus geoffroii fortis</i>).....	35

List of Graphs

Graph 1: Long term and 2021 monthly weather and climate data for Coolgardie and Koorarawalyee	3
Graph 2: Flora Species Accumulation Curve (Sample rarefaction)	27

List of Figures (out of text)

Figure 1: Survey Area
Figure 2: Soil Landscapes and Land Systems, and Hydrography
Figure 3: Broad Vegetation Types
Figure 4: Conservation Areas
Figure 5: Survey Effort
Figure 6: Threatened and Priority Flora Locations Identified by DBCA Database Searches
Figure 7: Vegetation Types and Priority Flora locations within the Survey Area
Figure 8: Vegetation Condition within the Survey Area
Figure 9: DBCA Threatened and Priority Fauna Records
Figure 10: Fauna Habitat and Conservation Significant Fauna Records

List of Appendices

Appendix A Flora and Fauna Literature Review
Appendix B Flora and Fauna Database Searches
Appendix C Coolgardie Flora Likelihood Table
Appendix D Inventory of Vascular Flora
Appendix E Flora Site Data
Appendix F Fauna Habitat Assessments
Appendix G Fauna Inventory

1 Introduction

1.1 The Project

Focus Mineral Limited (FML) commissioned 360 Environmental (part of SLR Consulting) to undertake a detailed flora, vegetation, and basic terrestrial vertebrate fauna assessment to support the environmental and planning approvals for proposed expansions to the FML mining operation in Coolgardie Western Australia (the Project).

The 1360.72 ha FML Coolgardie (the Survey Area) is located approximately 40 km southwest of Kalgoorlie in the Coolgardie bioregion of Western Australia. The Survey Area includes multiple polygons representing potential deposits, waste pits, haul roads, and associated infrastructure.

1.2 Objectives and Scope

The purpose of the biological survey was to delineate key flora, vegetation and terrestrial vertebrate fauna values within the Survey Area and identify potential environmental sensitivities that may impact The Project.

The scope of works includes:

- Undertake a desktop assessment including relevant database searches and a literature review to compile and summarise existing records of flora, vegetation, and terrestrial vertebrate fauna (including conservation significant species and communities) within the vicinity of the Survey Area
- Undertake a detailed flora and vegetation survey in accordance with EPA Technical Guidance (Environmental Protection Authority, 2016)
- Undertake targeted searching for flora of conservation significance within the Survey Area
- Undertake a basic terrestrial vertebrate fauna survey in accordance with EPA Technical Guidance (Environmental Protection Authority, 2020)
- Identify and assess conservation significant terrestrial vertebrate fauna or suitable conservation significant terrestrial vertebrate fauna habitat potentially occurring within the Survey Area
- Produce a technical report based on the findings of the above
- Supply a geospatial data package prepared in accordance with IBSA requirements.

This report presents the outcomes of the FML Coolgardie flora, vegetation, and terrestrial vertebrate fauna assessment undertaken to support the above objectives.

2 Background

2.1 Protection of Flora, Vegetation and Fauna

Western Australian flora and fauna is protected formally and informally by legislative and non-legislative measures:

Legislative measures:

- *Commonwealth Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act)
- *WA Biodiversity Conservation Act 2016* (BC Act)
- *WA Environmental Protection Act 1986* (EP Act)
- *WA Biosecurity and Agriculture Management Act 2007* (BAM Act).

Non-legislative measures:

- WA Department of Biodiversity Conservation and Attractions (DBCA) Priority lists for fauna, flora, and ecological communities
- Weeds of National Significance (WoNS)
- Recognition of locally significant populations by DBCA.

These protection mechanisms are supported by guidance documents published by the Environmental Protection Authority (EPA) and Department of Agriculture, Water, and the Environment (DAWE):

- *Technical Guidance – Flora and Vegetation Surveys for Environmental Impact Assessment* (Environmental Protection Authority, 2016)
- *Technical Guidance – Terrestrial Vertebrate Fauna Surveys for Environmental Impact Assessment* (Environmental Protection Authority, 2020)
- *Matters of National Environmental Significance Significant impact guidelines 1.1 Environment Protection and Biodiversity Conservation Act 1999* (Department of the Environment, 2013)
- *Survey Guidelines for Australia's Threatened Mammals* (Department of Sustainability Environment Population and Communities, 1999)
- *Survey Guidelines for Australia's Threatened Reptiles* (Department of Sustainability Environment Water Population and Communities, 2011)
- *Survey Guidelines for Australia's Threatened Birds Under the Environment Protection And Biodiversity Conservation Act 1999* (Department of the Environment Water Heritage and the Arts, 2010).


2.2 Existing Environment

2.2.1 Climate

The closest long-term Bureau of Meteorology weather station with a complete dataset is Coolgardie weather station (Station 012018) for climate data (i.e. temperature) and Koorarawalyee weather station (Station 012152) for rainfall data. The Coolgardie weather station is located approximately 75 km south of the Survey Area. Koorarawalyee weather station is located approximately 124 km southwest of the Survey Area.

The long-term mean minimum temperature for Coolgardie Weather Station ranges from 5.2°C (July) to 16.8°C (February) (1893 to 2020) and the long-term mean maximum temperature ranges from 16.1°C (July) to 33.3°C (January) (Graph 1) (Bureau of Meteorology, 2021).

The Koorarawalyee weather station recorded 336 mm of rainfall in the 12 months prior to the survey (October 2021 to November 2021), which is 66.4 mm above the long-term average of 269.6 for the same period (Bureau of Meteorology, 2021). In the three months prior to the survey (July 2021 to September 2021), 68.6 mm of rainfall was recorded, which is 7.6 mm above the long-term average of 61 mm for the same period (Bureau of Meteorology, 2021).



Graph 1: Long term and 2021 monthly weather and climate data for Coolgardie and Koorarawalyee

2.2.2 Interim Biogeographic Regionalisation of Australia

The Interim Biogeographic Regionalisation of Australia (IBRA) divides Australia into 89 bioregions based on major biological, geographical, and geological attributes. These bioregions are subdivided into 419 subregions as part of a refinement of the IBRA framework (Department of the Environment and Energy, 2016). The Survey Area occurs within the Coolgardie bioregion and Eastern Goldfield (COO3) subregion.

The Eastern Goldfield (COO3) subregion lies on the Yilgarn Cratons 'Eastern Goldfields Terrains'. The relief is subdued and comprises of gently undulating plains interrupted in the west with low hills and ridges of Archaean greenstones and in the east by a horst of Proterozoic basic granulite.

2.2.3 Soils and Land Systems

Soil landscapes and land system mapping of Western Australia describes broad soil and landscape characteristics from regional to local scales, ranging from 1:20,000 to 1:250,000 (Department of Primary Industries and Regional Development, 2018). The Survey Area occurs entirely within the BB5 land system, characterised by rocky ranges and hills of greenstones with basic igneous rocks as well as sandplains with brown calcareous loam (Figure 2) (Department of Primary Industries and Regional Development, 2018).

2.2.4 Hydrography

The Survey Area contains one small-permanent pool which is just south of Coolgardie-Esperance Highway, named Coolgardie Gorge. The Survey Area also contains four decommissioned deep abandoned mining pits which contain water and several minor watercourses that don't influence vegetation types present. The nearest major hydro geographical feature is a seasonally filled salt lake approximately 6 km east of the Survey area. The salt lake joins the Red, White and Douglas lakes as part of a larger interconnected salt lake system, which flows in a south westerly direction (Figure 2) (Department of Water and Environmental Regulation, 2018).

2.2.5 Broad Vegetation Types

Mapping of pre-European vegetation in Western Australia was completed on a broad scale (1:1,000,000) by Beard, (1976). These vegetation types were later refined by Shepherd, Beeston and Hopkins, (2002) resulting in 819 vegetation types.

Two broad vegetation associations are mapped over the Survey Area (Figure 3). Representation of the vegetation associations at a local, regional, and state level is shown in Table 2.

- **Coolgardie 9** – Medium woodland; Eucalyptus woodland / Eremophila sparse shrubland. Associated species are coral gum (*Eucalyptus torquata*) and goldfields blackbutt (*E. lesouefii*)
- **Coolgardie 1294** – Eucalyptus woodland. Wheatbelt; York gum, salmon gum etc. (*Eucalyptus loxophleba*, *E. salmonophloia*). Goldfields; gimlet, redwood etc. (*E. salubris*, *E. oleosa*). Riverine; rivergum (*E. camaldulensis*). Tropical; messmate, woolybush.

Table 2: Broad Vegetation Types within the Survey Area and their Representation at the State, Regional and Local Levels (Government of Western Australia, 2019)

System and Vegetation Association	Extent			
	Pre-European (ha)	Current (ha)	Remaining (%)	Managed in DBCA Lands (%) [*]
Representation across Western Australia				
9	240,509.33	235,161.94	97.78	8.07
1294	6,295.55	6,047.45	96.06	1.90
Representation across the Coolgardie Bioregion				
9	240,441.99	235,100.97	97.78	8.07
1294	6,295.55	6,047.45	96.06	1.90
Representation across the Eastern Goldfields Subregion				
9	235,047.15	229,757.07	97.75	8.26
1294	6,295.55	6,047.45	96.06	1.90
Representation across the Shire of Coolgardie				
9	166,572.37	163,720.39	98.29	9.81
1294	3,385.95	3,378.94	99.79	3.40

^{*}as a portion of the current extent

2.2.6 Environmentally Sensitive Areas

Environmentally Sensitive Areas (ESAs) are declared by the Department of Water and Environmental Regulation (DWER) to prevent the degradation of important environmental values such as Threatened flora, Threatened Ecological Communities (TECs) or significant wetlands.

The Survey Area does not occur within a mapped ESA. The nearest ESA is the Rowles Lagoon Conservation Park located approximately 52 km north-west of the Survey Area (Department of Water and Environmental Regulation, 2020).

2.2.7 Conservation Areas

The Survey Area is not located within any listed conservation area (Department of Biodiversity Conservation and Attractions, 2021a). The nearest conservation area is the Kangaroo Hills Timber Reserve which is situated adjacent to the southernmost section of the Survey Area separated by Nepean Road (Figure 4).

2.2.8 Land Use

The dominant land use of the Eastern Goldfields subregion include: Unoccupied Crown Land (UCL) and Crown reserves, Grazing-Native pastures-leasehold, freehold, conservation, and Mining leases (Cowan, 2001).

The Survey Area covers sections of land classified as: mining lease, prospecting licence, and miscellaneous licence (Data WA, 2021). The Survey Area has a long history of mining activities, with extensive mine diggings including deep open pits and deep localised mine shafts.

3 Methods

3.1 Desktop Assessment

3.1.1 Literature Review

Background information on the Survey Area and surrounds was compiled prior to the field survey (see Section 3.1.1). Historical vegetation mapping (Beard, 1976; Shepherd, Beeston and Hopkins, 2002), land systems mapping (Department of Agriculture and Food WA, 2012), and the IBRA classification system (Cowan, 2001) were consulted to provide broad contextual knowledge of the vegetation units and habitat likely to be encountered within the Survey Area.

The literature review also considered a selection of biological reports detailing assessments undertaken in the region that were publicly available:

- Coolgardie Landfill - Flora, vegetation and fauna habitat assessment (Strategen Environmental, 2019)
- Fauna survey for Mungari Gold Operations Cutters Ridge Project (Phoenix Environmental Sciences, 2019)
- Flora and Vegetation Impact Assessment - Medcalf Project (Botanica Consulting, 2020)
- Level 1 Vertebrate Fauna Risk Assessment for Lot 500 Kalgoorlie West (Terrestrial Ecosystems, 2018)
- Reconnaissance Flora and Vegetation Survey for the Mt Marion Project Area (Native Vegetation Solutions, 2019)
- Reconnaissance Flora and Vegetation Survey of Lot 500 Great Eastern Highway Kalgoorlie (Native Vegetation Solutions, 2018)
- Reconnaissance Flora and Vegetation Survey of the Spargos Project - October 2020 (Native Vegetation Solutions, 2020)
- Vegetation Clearing - Fauna Assessment (Davyhurst, Callion, Waihi, Siberia, Riverina Clearance Areas) (Biostat Pty Ltd, 2020).

3.1.2 Database Searches

Database searches were undertaken to compile a list of potential flora and fauna and identify potential conservation significant flora, fauna, and ecological communities within or surrounding the Survey Areas (Table 3). In addition, an EPBC Protected Matters Search (PMST) was undertaken to identify the potential for Matters of National Environmental Significance (MNES) to occur within or surrounding the Survey Areas (Department of Agriculture Water and the Environment, 2020).

The search area for each parameter was varied to reflect distances recommended by DBCA. The search areas are herein referred to collectively as the Study Area.

Table 3 Database Searches of the Survey Area

Database Name	Date Received	Search Target	Search Area
Threatened and Priority Ecological Communities database search (Department of Biodiversity Conservation and Attractions, 2022)	7 October 2021	TECs and PECs	50 km buffer around the Survey Area
Threatened and Priority Flora (TPFL) database search (Department of Biodiversity Conservation and Attractions, 2021d)	7 October 2021	Threatened and Priority Flora	60 km buffer around the Survey Area
Western Australian Herbarium flora database search (Department of Biodiversity Conservation and Attractions, 2021e)			60 km buffer around the Survey Area
DBCA Threatened and Priority Fauna database search (Department of Biodiversity Conservation and Attractions, 2021c)	18 October 2021	Threatened and Priority Fauna	50 km buffer around Survey Area
NatureMap (Department of Biodiversity Conservation and Attractions, 2021b)	27 October 2021 (flora)	Threatened and Priority flora and fauna, and inventory of potential flora and fauna	20 km buffer around the Survey Area (flora)
	02 November 2021 (fauna)		40 km buffer around Survey Area (fauna)
Protected Matters Search Tool (PMST) (Department of Agriculture Water and the Environment, 2021a)	25 October 2021	Commonwealth listed Threatened flora and fauna and TECs	50 km buffer around the Survey Area

3.1.3 Likelihood of Occurrence

Conservation significant flora and fauna species identified from the desktop assessment were assessed to determine the likelihood of their occurrence within the Survey Area, both prior to and post field survey. The assessment was completed based on the likelihood of occurrence criteria presented in Table 4.

Only species either recorded within the Survey Area or considered as having a high or medium likelihood of occurrence will be discussed in detail. Species classified as having a low likelihood of occurrence based on the above criteria will not be discussed unless a justification for this classification is required.

For fauna, taxa listed as Marine only under the EPBC Act were not included as conservation significant taxa because the Marine only listed taxa identified by the desktop assessment and field survey were common and widespread, taxa listed as Marine only do not constitute matters

of national environmental significance (MNES) under the EPBC Act, and the Survey Area does not contain any marine habitat.

Table 4 Likelihood of Occurrence Criteria

Rank	Criteria
Previously Recorded	The species has been previously recorded in the Survey Area
High (Likely to occur)	- Preferred habitat capable of supporting individuals or populations is present within the Survey Area - The survey Areas is within the taxon's known distribution - There are existing records of the species near the Survey Area (within 15 km) (flora) - There are existing records of the species near the Survey Area recorded within the last 15 years (fauna) - The species is strongly linked to a specific habitat, which is present in the Survey Area - The species has more general habitat preferences, and suitable habitat is present.
Medium (May occur)	- There is suitable (not necessarily preferred) habitat in the Survey Area, but the species is recorded infrequently in the locality - The Survey Area is within or near the taxon's known distribution - They Survey Area and surrounding area may support individuals or populations present within the Survey Area - There are existing records of the species from the locality (within 30 km for flora and 50 km for fauna), however: - The species is strongly linked to a specific habitat, of which only a small amount is present in the Survey Area - The species has more general habitat preferences, but only some suitable habitat is present.
Low (Unlikely to occur)	- No suitable habitat is present within the Survey Area, or the Survey Area is well outside the taxon's known distribution for the taxon is considered locally or regionally extinct - The Survey Area and surrounding habitat are unlikely to support individuals or populations of the taxon, however individuals may rarely occur as transients or vagrants - The species is linked to a specific habitat, which is absent from the Survey Area; or - Suitable habitat is present, however there are no existing records of the species from the locality despite reasonable previous search effort in suitable habitat; or - There is some suitable habitat in the Survey Area, however the species is very infrequently recorded in the locality.

3.2 Field Survey Effort

3.2.1 Trip 1

The biological survey was undertaken between the 11 - 15 October 2021.

The flora, vegetation and fauna survey was undertaken by Senior Botanist Jason Webb, Principal Zoologist Dr Michael Lohr and Ecologist Lachlan Crossley. The field team has over 20 years of combined experience conducting surveys of similar scope throughout Western Australia.

3.2.2 Trip 2

The biological survey was undertaken between the 15 - 19 November 2021.

The flora, vegetation and fauna survey was undertaken by Senior Botanist Jason Webb, Principal Ecologist Scott Walker and Ecologists Bridget Duncan and Lachlan Crossley. The field team has over 20 years of combined experience conducting surveys of similar scope throughout Western Australia.

The survey effort from both trips is shown in Figure 5.

3.3 Flora and Vegetation

3.3.1 Establishment of Flora Sites

Indicative flora sites were identified prior to the survey using aerial photography to estimate broad vegetation patterns within the Survey Area. The location and number of flora sites completed were adjusted on site to achieve sites most representative of the vegetation present.

Where possible, at least three flora sites were sampled for each vegetation type observed within the Survey Area. Some vegetation types that were not large enough to accommodate three flora sites had only one or two sites sampled.

Flora sites consisted of quadrats (20 m x 20 m) and relevés of approximately 400 m² where possible, or alternate configurations approximately equating to 400 m² (as required in areas such as drainage lines, gullies, and narrow ridge lines). A comprehensive record of the flora present at the time of sampling was recorded for both quadrat and relevé sites.

Flora site locations were recorded using a handheld GPS, the start and finish point of linear relevés, and the central point of circular relevés. At each flora site, the following was recorded:

- Site code
- Date and personnel
- Landform and soil description
- Relevant site descriptors including, slope, aspect, litter cover, bare ground cover and fire history
- Inventory of vascular flora including the approximate average height and percent foliar cover for each taxon recorded
- Vegetation description in accordance with the National Vegetation Information System (NVIS), Level 5 'association', whereby the dominant growth form, height, cover, and species (three species) for the three traditional strata (upper, mid, and ground) are described

- Vegetation condition in accordance with the Southwest and Interzone Botanical Provinces/Eremaean and Northern Botanical Provinces vegetation condition scale (Environmental Protection Authority, 2016a), and evidence of disturbance (for example clearing, rubbish, feral animals, weed incursion and evidence of feral animals and dieback) where present, and
- Photograph of the vegetation occurring within the site.

A total of 74 quadrats, and 22 relevés were established within the Survey Area. An additional 91 mapping notes were completed to aid vegetation mapping delineation.

3.3.2 Opportunistic Flora

Additional flora taxa observed opportunistically around flora sites or while traversing on foot within the Survey Area were also recorded. Where populations of conservation significant flora taxa, Declared Pests (DPs) or WoNS were encountered, a GPS location and a count of the individuals present was recorded.

3.3.3 Targeted Searching

Prior to the survey conservation significant flora with the likelihood or potential to occur within the Survey Area was compiled (see section 3.1.3). Field personnel familiarised themselves with photographs, reference samples and descriptions of these taxa before conducting the survey.

The entire Survey Area was not systematically searched. Rather, targeted searching focussed on habitat suitable for Conservation Significant Flora. Furthermore, potential habitat within the proposed footprint was prioritised for targeted searching over areas outside the proposed footprint.

Personnel also actively searched for conservation significant flora species in and around flora sites, while traversing on foot within the Survey Area and in known locations or preferred habitat encountered in the Survey Area.

Where Threatened or Priority flora were encountered in the field a GPS location was taken and a count of individuals was recorded, followed by a search in the local vicinity to determine if any other individuals were present nearby and delineate population boundaries where relevant. Specimens of any potential conservation significant flora that could not be identified in the field were collected for identification and lodgement at the Western Australian Herbarium (WAH).

3.3.4 Statistical Analyses

Statistical analysis of quadrat data was undertaken in accordance with EPA guidelines for a detailed flora and vegetation survey.

3.3.4.1 Species Accumulation Curve

Species accumulation curves were plotted using Primer-E version 6.1.5. to determine the adequacy of the survey (Clarke and Gorley, 2006). The treatments comprised Sobs (Mao Tao), to reflect the number of species observed (based on a given total of species recorded), and richness estimators Chao 1, Chao 2, Jackknife 1, Bootstrap and Michaelis-Menton to estimate the total sample size of flora taxa present. Species accumulation curves for this survey were calculated using data collected from the flora sites within the Survey Area. All flora taxa, both annual and perennial, within each flora site were used in generating the species accumulation curve. Unconfirmed flora were included if they were the only species present from a Genus.

3.3.5 Taxonomy and Nomenclature

Where field identification of plant taxa was not possible, specimens were collected for identification using resources of the WAH. Identification of flora collections was completed by experienced Taxonomist Frank Obbens.

The finalised species list was cross-referenced against current flora databases and relevant taxonomic literature (Western Australian Herbarium, 2021) to determine name currency, conservation status and known distribution of each taxon. Introduced species were compared against the current BAM Act Declared Plants list the WoNS list to determine their control status (Department of Agriculture Water and the Environment, 2021b; Department of Primary Industries and Regional Development, 2021).

Any conservation significant flora taxa, including potential Threatened and Priority species, range extensions and potential new taxa were submitted to the WAH for verification and lodgement. Where relevant, Threatened and Priority Flora Report Forms (TPFRFs) were submitted to DBCA.

3.3.6 Vegetation Unit and Condition Mapping

Broad vegetation and condition mapping was conducted in the field, with boundaries delineated over aerial photography, at a scale of 1:5,000. Broad vegetation units were refined based on taxonomic identification of flora collections, statistical analysis of data collected from the quadrats and relevés, and mapping notes taken during the field survey. Vegetation condition mapping was refined based on site data and mapping notes. Finalised polygons were digitised and produced as electronic mapping data using GIS software.

3.4 Vertebrate Fauna

3.4.1 Vertebrate Fauna Habitat Assessment

Fauna habitat assessments were undertaken throughout the Survey Area to identify fauna habitat values. Habitat assessment locations are displayed in Figure 5. The following information, which has been adapted from the habitat attributes listed in the Technical Guidance (Environmental Protection Authority, 2020), was collected at each habitat assessment site using Fulcrum, a mobile data collection app:

- Site photo
- Landform
- Soil type and colour
- Rock types, surface stone cover and size classes
- Key habitat and microhabitat features including leaf litter, logs, burrows, rocky outcrops, rock crevices, hollows, water sources
- Habitat quality, fire history and evidence of disturbance
- General description of vegetation structure.

Fauna habitat mapping boundaries were delineated over aerial imagery at a scale of approximately 1:5,000 based on field observations and fauna habitat assessment data. Polygons were digitised and produced as electronic mapping data using GIS software.

3.4.2 Opportunistic Fauna Records and Active Searches

Opportunistic observations of fauna were recorded throughout the Survey Area. Observations of primary evidence (direct sightings, calls) and secondary evidence (tracks, scats, diggings etc.) were recorded. Active searches were undertaken throughout the Survey Area in microhabitats likely to contain fauna. They primarily involved raking leaf litter, peeling bark, and splitting dead wood.

3.4.3 Identification and Taxonomy

Where there was doubt on a species name (through subsequent name changes or taxonomic reviews), an effort was made to determine the current scientific name for each taxon. Taxonomy and nomenclature in this report follows the WA Museum checklist 2021 (Western Australian Museum, 2021) where relevant.

4 Results

4.1 Limitations

Limitations and constraints of the flora, vegetation and fauna survey are detailed below in Table 5

Table 5: Limitations and Constraints Associated with the Survey

Variable	Degree of Limitation	Potential Constraints on Survey Outcomes
Survey Scope	Not a constraint	The detailed flora and vegetation survey was undertaken in accordance with EPA Technical Guidance (Environmental Protection Authority, 2016) and was considered appropriate to support approvals applications. Targeted searching for flora of conservation significance was undertaken, however, systematic searches were not feasible. Rather, targeted searching focused on habitat suitable for conservation listed flora within the proposed development footprint. A basic terrestrial vertebrate fauna survey was undertaken in accordance with EPA Technical Guidance (Environmental Protection Authority, 2020) and was considered appropriate to support approvals applications.
Availability of Data	Not a constraint	All data required to complete the scope of works including regional and local contextual information was available.
Site Access	Not a constraint	The Survey Area was able to be accessed by vehicle and on foot.
Survey Intensity and Resources	Not a constraint	A total of 96 flora sites (comprising 74 quadrats and 22 relevés) were sampled across the Survey Area. An additional 91 mapping notes were undertaken to aid vegetation mapping and delineation. Sufficient time was allocated to the flora and vegetation survey, given the size and complexity of the Survey Area, and the expected level of survey intensity. The survey effort was considered adequate to assess the flora and vegetation values of the Survey Area and provide information required to support approvals applications. Eighty fauna habitat assessments were completed, and 61 fauna taxa were recorded opportunistically throughout the Survey Area. The survey effort was appropriate for a basic terrestrial vertebrate fauna survey.

Variable	Degree of Limitation	Potential Constraints on Survey Outcomes
Experience	Not a constraint	The flora and vegetation surveys were undertaken by Senior Botanist Jason Webb and Botanist Bridget Duncan, who have over 5 years combined experience conducting surveys of similar scope throughout Western Australia. Identification of flora collections was completed by experienced taxonomist Frank Obbens at the WAH. Any specimens with novel characteristics were submitted to the WAH for formal identification. The fauna survey was undertaken by Lachlan Crossley (Ecologist), Dr. Michael Lohr (Principal Zoologist) and Scott Walker (Principal Ecologist), who have 25 years of combined experience conducting fauna surveys. Fauna identification was reviewed by Dr. Michael Lohr (Principal Zoologist) and Evan Webb (Senior Zoologist), using photos of tracks and scats taken during the field survey.
Timing, weather, season	Not a constraint Partial limitation for fauna (Amphibians)	The recommended primary survey period for the region as per the EPA Technical Guidance, is September to November. Survey was conducted in October and November 2021. Rainfall was sufficient prior to the commencement of the field survey and air temperatures were slightly below average. Therefore, weather conditions were not a constraint. According to the EPA Technical Guidance, mammals do not have a preferred time of year for optimal observation. For reptiles, the guidance suggests October to December (for primary survey). As the surveys were conducted in October and November, timing was suitable for reptiles. The optimum timing for observation of amphibians is between May and August (autumn-winter breeders) and November-December (summer breeders). Therefore, the timing of the surveys was not suitable for autumn-winter breeders. The optimum time to observe birds is between September and December for most bush birds and between November and March for migratory birds. As the surveys were conducted in October and November, timing was suitable for birds.
Life Forms Sampled	Not a constraint	The Survey Area was traversed by vehicle and on foot and representative sites of all remnant vegetation were sampled. All flora species encountered within the Survey Area were recorded. A total of 145 vascular flora taxa were recorded from the Survey Area, comprising 93.8% native flora taxa and 6.2% introduced flora taxa. Of the 145 flora taxa recorded, 18 taxa (12.4%), could not be identified to species level because they were sterile at

Variable	Degree of Limitation	Potential Constraints on Survey Outcomes
		the time of the survey. This was not considered a constraint as it represented a very small portion of the flora sampled. None of the unknown flora taxa collected were analogous to Threatened or Priority flora taxa identified by the database searches as likely to occur within the Survey Area, nor were they representative of flora of other significance. The basic vertebrate fauna survey used a range of techniques to detect fauna taxa within the Survey Area. A total of 61 vertebrate fauna taxa were recorded within the Survey Area, comprising 54 confirmed native taxa and seven introduced taxa. All fauna taxa recorded were able to be identified to species level by direct or indirect observation with a high level of confidence.
Mapping Reliability	Not a constraint	Vegetation types were described and mapped based on quadrat/relevé data and additional mapping notes taken during the field survey. The number of sites and coverage of the Survey Area was considered sufficient to inform the survey. Fauna habitat mapping was based largely on vegetation mapping and there were no constraints on mapping reliability.
Disturbances (fire, flood etc.)	Not a constraint	No disturbances occurred during any of the surveys. Areas of disturbance associated with clearing, mining, rubbish, weeds, and the presence of introduced fauna were recorded but were not a constraint on the results of the survey.
Completeness	Not a constraint	The survey was considered complete for a detailed flora and vegetation survey and basic vertebrate fauna survey. All vegetation types and fauna habitats were surveyed and delineated within the Survey Area, however not all vegetation types were surveyed with three flora sites; five vegetation types were surveyed with one or two flora sites.

4.2 Flora and Vegetation

4.2.1 Literature Review

The key findings of the flora and vegetation reports reviewed are summarised in Appendix A1.

4.2.2 Database Searches

Database searches identified 90 conservation significant flora species occurring within 50 km of the Survey Area (Figure 6, Appendix B), comprising:

- Three Threatened taxa
- Twenty-seven Priority 1 taxa

- Fourteen Priority 2 taxa
- Forty Priority 3 taxa, and
- Six Priority 4 taxa.

4.2.3 Likelihood of Occurrence

The pre-survey likelihood of occurrence assessment identified that of the 90 conservation significant flora species identified by the desktop assessment:

- 20 were considered to have a high likelihood of occurrence
- 18 were considered to have a medium likelihood of occurrence
- 51 were considered to have a low likelihood of occurrence.

Following the survey, the likelihood of occurrence was re-assessed and found:

- Two taxa were considered to have a high likelihood of occurrence, due to close records and existing, contiguous habitat within the Survey Area.

The likelihood of occurrence assessment is provided in Appendix C.

4.2.4 Vegetation of Conservation Significance

Database searches did not return any listed Threatened Ecological Communities or Protected Ecological Communities within 50 km of the Survey Area.

4.2.5 Flora Composition

The survey recorded a total of 149 taxa from 78 genera across 35 families (Appendix D). The dominant families were Chenopodiaceae (26 taxa) and Myrtaceae (15 taxa). The most dominant genera were *Eucalyptus* (12 taxa) and *Eremophila* (10 taxa).

4.2.6 Flora of Conservation Significance

4.2.6.1 Threatened or Priority Flora

No Threatened flora species pursuant to the EPBC Act 1999 and/or gazetted as Threatened pursuant to the BC Act 2016 were recorded during the survey.

One Priority taxon was recorded within the Survey Area (Figure 7a, Plate 1). A population consisting of approximately 10 individuals of *Austrostipa blackii* (P3) was recorded in quadrat C1Q01. This taxon is a tufted, perennial, grass, typically growing up to 1 m high that flowers between September and November. The WAH has 41 specimens lodged with records across the Avon Wheatbelt, Coolgardie and Yalgoo bioregions (Western Australian Herbarium, 2022). The habitat descriptions for this taxon are typically mixed Eucalypt species, tall Acacia shrublands and open woodlands of *Allocasuarina dielsiana* and *Allocasuarina tessellata* (Western Australian Herbarium, 2022). *Austrostipa blackii* (P3) grows on gently inclined lower slopes of basalt with red-brown deep sandy clay loam soils (Western Australian Herbarium, 2022).



Plate 1: *Austrostipa blackii* (P3)

Within the Survey Area it was recorded within vegetation unit AcEoaDI (quadrat C1Q01), in association with *Acacia collegialis*, *Eremophila* spp. and *Dodonaea lobulata*. This vegetation type was sampled heavily and was not recorded at any other site.

4.2.7 Introduced Taxa

A total of 13 introduced taxa were recorded within the Survey Area, representing 8.7% of the total taxa recorded (Table 6). One taxon (**Opuntia stricta*) is a Declared Pest and listed as a WoNS (DAWE, 2021b; DPIRD, 2021). Eight individuals of **O. stricta* were recorded (Figure 8).

Three introduced taxa are unlisted under the BAM Act, which prohibits them access to Western Australia, however they are not assigned a control management category (DPIRD, 2021).

Table 6: Introduced Flora Species within the Survey Area

Species	Common Name	Status under BAM Act	WoNS
<i>*Agave americana</i>	Century plant	Permitted - s11	No
<i>*Asphodelus fistulosus</i>	Onion Weed	Permitted - s11	No
<i>*Carrichtera annua</i>	Ward's Weed	Permitted - s11	No
<i>*Centaurea melitensis</i>	Maltese Cockspur	Permitted - s11	No
<i>*Crassula ovata</i>	Jade Tree	Permitted - s11	No

Species	Common Name	Status under BAM Act	WoNS
* <i>Heliotropium europaeum</i>	Common Heliotrope	Permitted - s11	No
* <i>Nicotiana glauca</i>	Tree Tobacco	Permitted - s11	No
* <i>Oligocarpus calendulaceus</i>	-	Unlisted – s14	No
* <i>Rumex vesicaria</i>	Ruby Dock	Unlisted – s14	No
* <i>Salvia verbenaca</i>	Wild Sage	Permitted - s11	No
* <i>Schinus molle</i> var. <i>areira</i>	-	Unlisted – s14	No
* <i>Sonchus asper</i>	Rough Sowthistle	Permitted - s11	No
* <i>Opuntia stricta</i>	Common Prickly Pear	Declared Pest - s22(2)	Yes

4.2.8 Unconfirmed Flora

Eighteen specimens (12.1% of the taxa recorded) could not be identified to species level because the taxa were sterile at the time of the survey (Appendix D). Of these, all but one, *Malvaceae* sp., were able to be identified to genus level, and two were assigned a tentative genus, *?Enchylaena tomentosa* and *?Pimelea* sp.

None of the unconfirmed flora taxa were analogous to Priority flora taxa identified by the database searches.

4.2.9 Vegetation Condition

The Survey Area has been subjected to medium to high level disturbances, including historical small and large scale mine excavations, makeshift tracks, cattle grazing, weeds, and litter. As such, vegetation condition with the Survey Area was predominantly Excellent, and ranged to Completely Degraded (where all vegetation had been completely cleared) (Keighery, 1994) (Table 7, Figure 7).

Table 7 Area (ha) covered by each vegetation condition category within the Survey Area

Vegetation Condition	Area (ha)	% of Survey Area
Excellent	790.6	58.1%
Very Good	327.9	21.2%
Good	78.4	5.8%
Completely Degraded	163.6	12%
Total	1360.7	100

4.2.10 Vegetation Types

Ten vegetation types were described and mapped across four broad landforms within the Survey Area (plains, rocky hills, claypan, and lake) (Table 8, Figure 8). The majority of the Survey Area was dominated by open plains with a mosaic of low *Eucalyptus* spp. (34.8%) which grew together across the plains, and Salmon Gum (*Eucalyptus salmonophloia*) open woodlands (31.4%). Small rocky hills dominated by *Acacia collegialis* shrubland also covered 13.1% of the Survey Area. While 11.5% of the Survey Area was considered to be 'Cleared', some of this was showing signs of revegetation with both native species (such as *Atriplex* and *Maireana*) and introduced (weed) species. These areas did not appear to have been intentionally revegetated.

Detailed data sheets for each flora site are provided in Appendix E.

Table 8: Vegetation Types Occurring within the Survey Area

Vegetation Unit and Description*	Local Landform	Total Area, Proportion of the Survey Area	Sites	Photograph
<u>AcEoaDI</u> <i>Acacia collegialis</i> (<i>A. acuminata</i>) tall shrubland over <i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>, <i>E. georgei</i>, <i>A. tetragonophylla</i> (<i>Senna artemisioides</i> subsp. <i>filifolia</i>, <i>Exocarpos aphyllus</i>) mid shrubland over <i>Dodonaea lobulata</i> (<i>Atriplex vesicaria</i>, <i>Ptilotus obovatus</i> var. <i>obovatus</i>) low shrubland	Rocky hills	178.5 ha 13.1%	C1Q01, C1Q04, C1Q06, C1Q09, C1Q21, C1Q25, C1Q26, C1Q28, C1Q29, C1Q30, C1R04, C2Q07, C2Q09, C2Q14, C2Q18, C2Q32, C2Q33, C2Q34, C2Q35, C2Q38, C2Q40	
<u>Ec</u> <i>Eucalyptus celastroides</i> low woodland	Low slopes of rocky hills	7.4 ha 0.5%	C2Q43, C2R17	

Vegetation Unit and Description*	Local Landform	Total Area, Proportion of the Survey Area	Sites	Photograph
<u>EgAhSaf</u> <i>Eucalyptus griffithsii</i> low open woodland over <i>Acacia hemiteles</i> and <i>Dodonaea stenozyga</i> (<i>A. jennerae</i>, <i>Alyxia buxifolia</i>) mid shrubland over <i>Senna artemisioides</i> subsp. <i>filifolia</i> and <i>Atriplex vesicaria</i> (<i>Olearia muelleri</i>) low open shrubland	Plains	75.6 ha 5.6%	C1Q10, C1Q20, C1R13, C2Q27, C2Q29, C2Q31, C2Q41, C2Q50, C3Q01, C3Q02A, C3R01	
<u>EooEiiDs</u> <i>Eucalyptus oleosa</i> subsp. <i>oleosa</i> low open woodland over <i>Eremophila interstans</i> subsp. <i>interstans</i> mid isolated shrubs over <i>Dodonaea stenozyga</i>, <i>Eremophila glabra</i> subsp. <i>glabra</i>, and <i>Olearia muelleri</i> low open shrubland	Plains	11.1 ha 0.8%	C2Q11, C2Q37	

Vegetation Unit and Description*	Local Landform	Total Area, Proportion of the Survey Area	Sites	Photograph
<u>ErMhOm</u> <i>Eucalyptus ?ravidia</i> low open woodland over <i>Melaleuca ?hamata</i> (<i>M. pauperiflora</i> subsp. <i>fastigiata</i>, <i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>) tall open shrubland over <i>Olearia muelleri</i> low isolated shrubs	Plains	4.7 ha 0.3%	C2Q02, C2Q04, C2Q06	
<u>EsEiiAv</u> <i>Eucalyptus salmonophloia</i> mid open woodland over <i>Eremophila intertans</i> subsp. <i>interstans</i> (<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>, <i>Senna artemisioides</i> subsp. <i>filifolia</i>) tall to mid isolated shrubs over <i>Atriplex vesicaria</i> low open shrubland	Plains	428.0 ha 31.4%	C1Q03, C1Q22, C1Q23, C1Q24, C2Q12, C2Q19, C2Q20, C2Q21, C2Q24, C2Q26, C2Q28, C2Q46, C2Q52, C2Q54	

Vegetation Unit and Description*	Local Landform	Total Area, Proportion of the Survey Area	Sites	Photograph
<u>EspEiiSaa</u> <i>Eucalyptus salmonophloia</i> mid isolated trees over a mosaic of <i>E. celastroides</i>, <i>E. clelandiorum</i>, and <i>E. torquata</i> low open woodland over <i>Eremophila interstans</i> subsp. <i>interstans</i> (<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>) mid isolated shrubs over <i>Senna artemisioides</i> subsp. <i>artemisioides</i>, <i>S. artemisioides</i> subsp. <i>filifolia</i>, and <i>Atriplex vesicaria</i> low open shrubland	Plains, low hills	472.3 ha 34.8%	C1Q02, C1Q05, C1R05, C1R09, C1R10, C2Q01, C2Q03, C2Q05, C2Q08, C2Q10, C2Q13, C2Q15, C2Q16, C2Q36, C2Q39, C2Q42, C2Q45, C2R09, C3Q02, C3Q02B, C3Q06	
<u>EtEaEpa</u> <i>Eucalyptus torquata</i> low open woodland over <i>Exocarpos aphyllus</i> mid isolated shrubs over <i>Eremophila parvifolia</i> subsp. <i>auricampi</i>, <i>Westringia rigida</i>, and <i>Scaevola spinescens</i> (<i>Olearia muelleri</i>) low open shrubland	Plains, lower slopes of rocky hills	20.7 ha 1.5%	C1Q07, C1Q08, C1Q27, C1R01, C1R02, C1R03, C1R11	

Vegetation Unit and Description*	Local Landform	Total Area, Proportion of the Survey Area	Sites	Photograph
SI <i>Streptoglossa liatroides</i> low open herbland	Clay	0.3 ha <0.1%	C2R44	
Lake Coolgardie Gorge, natural semi-permanent pool system, with a fringe of native and introduced flora taxa including <i>Eucalyptus</i> spp., <i>*Schinus molle</i> var. <i>areira</i>, <i>*Heliotropium europaeum</i>, and <i>*Sonchus asper</i>.	Lake system	5.6 ha 0.4%	C2R07	

Vegetation Unit and Description*	Local Landform	Total Area, Proportion of the Survey Area	Sites	Photograph
Cleared Cleared or historically cleared areas including mine pits and borrow pits (often filled with water), bitumen roads, and dirt tracks. Some of these areas were showing signs of revegetation. With occasional <i>Eucalyptus griffithsii</i>, <i>Atriplex vesicaria</i>, <i>Maireana</i> spp., and assorted weed species.	Plains	156.9 ha 11.5%	C2Q17, C2Q25	

*Brackets indicate species that may or may not be present, but were observed as dominant/sub-dominant at some of the sites that make up the vegetation type


4.2.11 Groundwater Dependent Ecosystems

The Groundwater Dependent Ecosystem (GDE) Atlas identified the Survey Area did not contain any GDEs (Bureau of Meteorology, 2022). The results from the flora and vegetation survey support the GDE Atlas as vegetation in the Survey Area comprised xerophytic species that are not groundwater dependent.

4.2.12 Survey Adequacy

Ninety-six flora sites (74 quadrats and 22 relevés) were sampled across the Survey Area. This was adequate to ensure sufficient coverage given the overall size of the Survey Area. Two vegetation types were sampled with only two sites, while a further two were sampled with a single site. This was due to the vegetation being highly restricted in the Survey Area.

The sample rarefaction (Mao Tau) for the Survey Area produced a smooth S_{obs} curve, steadily increasing towards asymptote indicating that the survey captured the majority of flora taxa present with the Survey Area (Graph 2).



Graph 2: Flora Species Accumulation Curve (Sample rarefaction)

Estimated species richness for the Survey Area ranged from 122.1 to 167.6, with an observed value of 130 taxa (Table 9). Richness estimators indicated that the survey was approximately 77.5% to 106.4% adequate in recording the full complement of vascular flora taxa within the Survey Area (Table 9).

The data used to produce the species accumulation curve was conservative because opportunistic species (which are not associated with a site) were not included. Unconfirmed flora taxa were included if they were the only species present from a Genus.

Table 9: Species Richness Indicators

Treatment	Expected Species Richness	Percentage Adequate
Chao 1	162.8	79.8
Chao 2	162.8	79.8
Jackknife 1	167.6	77.5
Bootstrap	147.6	88.0
Michaelis-Menton	122.14	106.4

4.3 Vertebrate Fauna

4.3.1 Desktop Assessment

The desktop assessment identified 20 conservation significant terrestrial vertebrate fauna species potentially occurring within the Survey Area, comprising 17 birds and three mammals.

Key findings of the literature review are summarized in Appendix A2. Database search results are presented in Figure 9 and Appendix B.

4.3.2 Fauna Habitat

Five broad fauna habitats were identified and mapped within the Survey Area (Figure 10). Habitat condition varied throughout the Survey Area with the most prolific forms of disturbance being previous mining activity, litter, and vehicle tracks. Other disturbances included clearing, erosion, weeds, and introduced fauna.

A description, extent within the Survey Area, and a representative photo is provided for each fauna habitat in Table 10. Small discrepancies in fauna habitat extents (i.e., not adding up to the exact area extent of the Survey Area) are due to rounding. Fauna habitat mapping is presented in Figure 10 and site sheets for each habitat assessment are shown in Appendix F.

Table 10: Fauna Habitat Types within the Survey Area

Fauna Habitat	Total Area, Proportion of the Survey Area	Habitat Description	Representative Photo
Acacia Shrublands	75.8 ha	<i>Eucalyptus griffithsii</i> low open woodland over <i>Acacia hemiteles</i> and <i>Dodonaea stenozyga</i> (<i>A. jennerae</i>, <i>Alyxia buxifolia</i>) mid shrubland over <i>Senna artemisioides</i> subsp. <i>filifolia</i> and <i>Atriplex vesicaria</i> (<i>Olearia muelleri</i>) low open shrubland Hollow logs, leaf litter, and peeling bark are present throughout this habitat type. These microhabitat features provide shelter and foraging opportunities for small reptiles, birds, and mammals. Conservation significant fauna such as the Malleefowl (<i>Leipoa ocellata</i>) and the Western Quoll, Chuditch (<i>Dasyurus geoffroii fortis</i>) may use this habitat. Habitat condition varied from disturbed to very good. Evidence of disturbance includes previous mining activity, vehicle tracks, litter, and the presence of introduced fauna.	

Fauna Habitat	Total Area, Proportion of the Survey Area	Habitat Description	Representative Photo
Eucalyptus Woodlands	943.9 ha	Mixed <i>Eucalyptus</i> sp. woodlands over <i>Acacia</i> sp., <i>Dodonaea</i> sp., <i>Eremophila</i> sp. or <i>Melaleuca</i> sp., mixed shrublands. Peeling bark, woody debris, leaf litter and hollow logs were observed throughout this habitat type. These microhabitat features provide shelter for small reptiles and mammals. The canopy of trees provides shelter and foraging habitat for birds. Evidence of the conservation significant Western Quoll, Chuditch (<i>Dasyurus geoffroii fortis</i>) was recorded in this habitat type. The Malleefowl (<i>Leipoa ocellata</i>) may also use this habitat. Habitat condition varied from highly degraded to high quality. Evidence of disturbance includes previous mining activity, vehicle tracks, litter, and the presence of introduced fauna.	

Fauna Habitat	Total Area, Proportion of the Survey Area	Habitat Description	Representative Photo
Open Water	2.7 ha	Coolgardie Gorge was located within the Survey Area, with a fringe of introduced flora taxa including <i>*Schinus molle</i> var. <i>areira</i>, <i>*Heliotropium europaeum</i>, and <i>*Sonchus asper</i>. Additional standing water was identified within remnant mine pits, however this may only be temporary. Microhabitats and habitat features unique to this habitat type include exfoliating rock, steep rocky crevices and water sources. Migratory birds such as the Common Sandpiper (<i>Actitis hypoleucos</i>), Sharp-tailed Sandpiper (<i>Calidris acuminata</i>), Curlew Sandpiper (<i>Calidris ferruginea</i>), Wood Sandpiper (<i>Tringa glareola</i>), Common Greenshank (<i>Tringa nebularia</i>) and Glossy Ibis (<i>Plegadis falcinellus</i>) may use these areas. Habitat condition varies from highly degraded to disturbed. Evidence of disturbance includes previous mining activity, vehicle tracks, litter, and the presence of introduced fauna.	

Fauna Habitat	Total Area, Proportion of the Survey Area	Habitat Description	Representative Photo
Rocky Slopes	178.5	<i>Acacia collegialis</i> (<i>A. acuminata</i>) tall shrubland over <i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>, <i>E. georgei</i>, <i>Acacia tetragonophylla</i> (<i>Senna artemisioides</i> subsp. <i>filifolia</i>, <i>Exocarpos aphyllus</i>) mid shrubland over <i>Dodonaea lobulata</i> (<i>Atriplex vesicaria</i>, <i>Ptilotus obovatus</i> var. <i>obovatus</i>) low shrubland Leaf litter, peeling bark, rock crevices, and woody debris provides shelter for small reptiles and mammals. Shrublands provide shelter and foraging habitat for birds, reptiles, and mammals. Malleefowl (<i>Leipoa ocellata</i>) may forage within this habitat, however, it is unlikely to nest in this habitat due to the rocky substrate. The Western Quoll, Chuditch (<i>Dasyurus geoffroii fortis</i>) may use this habitat. Habitat condition varies from disturbed to very good. Evidence of disturbance includes previous mining activities, vehicle tracks, litter, and the presence of introduced fauna.	

Fauna Habitat	Total Area, Proportion of the Survey Area	Habitat Description	Representative Photo
Cleared Areas	160.9 ha	Cleared or historically cleared areas including mine pits and borrow pits (often filled with water), bitumen roads, and dirt tracks. Areas of completely cleared land include vehicle access tracks and previous mining activity including mining pits. These areas have limited value as habitat for fauna. The mining pits may provide some habitat for small reptiles and mammals in the form of steep rocky crevices and isolated shrubs and grasses. Evidence of disturbance includes previous mining activity, vehicle tracks, litter, and the presence of introduced fauna.	

4.3.3 Fauna Records

The terrestrial vertebrate fauna survey recorded a total of 61 fauna taxa from 33 families. An inventory of fauna recorded during the field survey is provided in Appendix G.

Table 11: Overview of Vertebrate Fauna Species Recorded

Fauna group	Number of species	Number of families
Birds	42	21
Mammals	10	7
Reptiles	9	5
Amphibians	0	0
Total	61	33

4.3.3.1 Birds

A total of 42 avian taxa from 21 families were recorded throughout the Survey Area. The most recorded taxon was the Budgerigar (*Melopsittacus undulatus*), followed by the Australian Ringneck (*Platycercus zonarius*) and the Singing Honeyeater (*Gavicalis virescens*). The most diverse avifauna families were Meliphagidae (seven taxa) and Acanthizidae (six taxa).

4.3.3.2 Mammals

A total of three native mammals were recorded within the Survey Area. The most recorded native mammal taxon was the Western Grey Kangaroo (*Macropus fuliginosus melanops*). Seven introduced mammal taxa were recorded in the Survey Area, Goat (*Capra hircus*), European Cattle (*Bos primigenius taurus*), Dog/Dingo (*Canis familiaris*), Red fox (*Vulpes vulpes*), Horse (*Equus caballus*), Cat (*Felis catus*) and Rabbit (*Oryctolagus cuniculus*).

4.3.3.3 Reptiles

A total of nine reptilian species from five families were recorded throughout the Survey Area. The most recorded species was the Bobtail (*Tiliqua rugosa*) followed by the Bynoe's Gecko (*Heteronotia binoei*). The most diverse reptilian family was Agamidae (four taxa).

4.3.3.4 Amphibians

No amphibians were recorded during the field survey.

4.3.4 Conservation Significant Fauna

One conservation significant taxon was recorded via observation of a scat during the fauna survey, the Western Quoll, Chuditch (*Dasyurus geoffroii fortis*), which is listed Vulnerable under the BC and EPBC Act (Plate 2).



Plate 2: Scat of Western Quoll, Chuditch (*Dasyurus geoffroii fortis*)

GPS Coordinates: Latitude: -30.920324, Longitude: 121.184610. Image Source 360 Environmental.

Three conservation significant fauna taxa were assessed as having a high likelihood of occurrence within the Survey Area:

- Malleefowl (*Leipoa ocellata*), Vulnerable under the BC and EPBC Act
- Common Sandpiper (*Actitis hypoleucos*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act
- Common Greenshank (*Tringa nebularia*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

Four conservation significant taxa were assessed as having a medium likelihood of occurrence within the Survey Area:

- Sharp-tailed Sandpiper (*Calidris acuminata*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act
- Curlew Sandpiper (*Calidris ferruginea*), Critically Endangered, International Agreement under the BC Act and Critically Endangered, Migratory and Marine under the EPBC Act

- Wood Sandpiper (*Tringa glareola*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act
- Glossy Ibis (*Plegadis falcinellus*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

Twelve conservation significant taxa were assessed as having a low likelihood of occurrence within the Survey Area. Further details regarding recorded and potential conservation significant fauna are provided below in Table 12.

Table 12: Conservation Significant Fauna Likelihood of Occurrence

Conservation Status: State - Listed under Biodiversity Conservation Act 2016 or Department of Biodiversity, Conservation and Attractions Conservation List, Federal - Listed under Environmental Protection and Biodiversity Conservation Act 1999. CR - Critically Endangered, EN - Endangered, VU - Vulnerable, IA/MI - Migratory, CD - Conservation Dependent fauna, OS - Other Specially Protected fauna, MA - Marine, P - Listed as Priority by DBCA.

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
AVIAN											
<i>Apodidae</i>	<i>Apus pacificus</i>	Pacific Swift (Fork-tailed Swift)	IA	MI, MA		x				Low	Three records within 100 km on the Survey Area (Atlas of Living Australia, 2022a). Species may fly over the Survey Area as it covers a wide range of airspace over varied habitat (Morcombe, 2003).
<i>Cacatuidae</i>	<i>Calyptrorhynchus latirostris</i>	Carnaby's Cockatoo	EN	EN	x		x	4		Low	This species was recorded 30 km NE of the Survey Area in Kalgoorlie (Department of Biodiversity Conservation and Attractions, 2021c). However, these records occur well outside the known distribution of the species and likely represent a vagrant occurrence of the taxon (Department of Sustainability

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
											Environment Water Population and Communities, 2012).
Charadriidae	<i>Thinornis cucullatus</i>	Hooded Plover (Hooded Dotterel)	P4	MA			x			Low	Closest record 42 km NE of the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). No suitable habitat within the Survey Area (Menkhorst <i>et al.</i> , 2017).
Falconidae	<i>Falco hypoleucos</i>	Grey Falcon	VU	VU		x				Low	No recent nearby records. Some records within 100 km on the Survey Area (Atlas of Living Australia, 2022b). Preferred nesting habitat absent. May use Survey Area for hunting (Menkhorst <i>et al.</i> , 2017).
Megapodiidae	<i>Leipoa ocellata</i>	Malleefowl	VU	VU	x	x	x	79		High	Recent records within 1 km of the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Recorded 40 km north of the Survey Area in 2019 (Phoenix

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
											Environmental Sciences, 2019). Suitable habitat present, unburned mallee and woodland with abundant litter and low scrub (Morcombe, 2003)
Motacillidae	<i>Motacilla cinerea</i>	Grey Wagtail	IA	MI, MA		x				Low	Survey Area is well outside of the distribution of this species Area (Atlas of Living Australia, 2022b). Some suitable habitat present in parts i.e. water bodies (Morcombe, 2003).
Psittaculidae	<i>Pezoporus occidentalis</i>	Night Parrot	CR	EN		x				Low	No records within 100 km on the Survey Area (Atlas of Living Australia, 2022b). No suitable habitat within the Survey Area (Morcombe, 2003).
Scolopacidae	<i>Actitis hypoleucos</i>	Common Sandpiper	IA	MI, MA	x	x	x	3		Previously Recorded	Three recent records within the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Some suitable habitat present i.e. interior wetlands – narrow muddy edges of billabongs (Morcombe, 2003).

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
<i>Scolopacidae</i>	<i>Calidris acuminata</i>	Sharp-tailed Sandpiper	IA	MI, MA	x	x	x	2		Medium	Nearest record 15 km NW of the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Suitable habitat present (water bodies) (Morcombe, 2003).
<i>Scolopacidae</i>	<i>Calidris alba</i>	Sanderling	IA	MI, MA	x		x	1		Low	Nearest record in Kalgoorlie (Department of Biodiversity Conservation and Attractions, 2021c). No suitable habitat within the Survey Area (Menkhorst et al., 2017).
<i>Scolopacidae</i>	<i>Calidris ferruginea</i>	Curlew Sandpiper	CR, IA	CR, MI, MA	x	x	x	1		Medium	Nearest record 15 km NW of Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Suitable habitat within the Survey Area i.e. around lakes, dams (Morcombe, 2003).
<i>Scolopacidae</i>	<i>Calidris melanotos</i>	Pectoral Sandpiper	IA	MI, MA		x				Low	Survey Area is well outside of the distribution of this species Area (Atlas of Living Australia, 2022b). Some suitable habitat present in parts i.e. inland

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
											water bodies (Morcombe, 2003).
<i>Scolopacidae</i>	<i>Calidris ruficollis</i>	Red-necked Stint	IA	MI, MA	x		x	1		Low	Nearest record 15 km NW of Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). No suitable habitat within the Survey Area i.e., mudflats (Morcombe, 2003).
<i>Scolopacidae</i>	<i>Tringa brevipes</i>	Grey-tailed Tattler	IA, P4	MI, MA	x		x	1		Low	Nearest record 20 km NE of Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). No suitable habitat within the Survey Area, coastal in Australia (Menkhorst et al., 2017).
<i>Scolopacidae</i>	<i>Tringa glareola</i>	Wood Sandpiper	IA	MI, MA	x		x			Medium	Nearest record 30 km NE of the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Suitable habitat within the Survey Area i.e., freshwater

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
											wetlands (Menkhorst et al., 2017).
Scolopacidae	<i>Tringa nebularia</i>	Common Greenshank	IA	MI, MA	x	x	x	2		Previously Recorded	Recent record within the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Suitable habitat within the survey areas i.e. temporary inland wetlands (Morcombe, 2003).
Threskiornithidae	<i>Plegadis falcinellus</i>	Glossy Ibis	IA	MI, MA			x			Medium	Only record 27 km NE of the Survey Area in Kalgoorlie (Department of Biodiversity Conservation and Attractions, 2021c). Some suitable habitat present i.e. temporary wetlands (Morcombe, 2003).
MAMMALIAN											
Dasyuridae	<i>Dasyurus geoffroii fortis</i>	Western Quoll, Chuditch	VU	VU		x	x		x	Recorded	A scat was found within the Survey Area during the current survey, identified to be this species. Some suitable habitat present mallee

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
											shrubland (Van Dyck and Strahan, 2008).
Myrmecobiidae	<i>Myrmecobius fasciatus</i>	Numbat, Walpurti	EN	EN	x					Low	No nearby records (Atlas of Living Australia, 2022b). Some suitable habitat is present i.e. eucalypts and wandoo woodland (Van Dyck and Strahan, 2008). The only natural populations exist well outside the Survey Area in the far north of WA (Department of Biodiversity Conservation and Attractions, 2017b).
Thylacomyidae	<i>Macrotis lagotis</i>	Bilby, Dalgyte	VU	VU	x					Low	No nearby records (Atlas of Living Australia, 2022b). No suitable habitat is present (Van Dyck and Strahan, 2008).
INSECTS											
Lycaenidae	<i>Jalmenus aridus</i>	Inland Hairstreak	P1 (not WAM)		x		x			Medium	Nearest record 19 km NE of the Survey Area (Department of Biodiversity Conservation and Attractions, 2021c). Suitable habitat present within the Survey Area

Family	Scientific Name	Common Name	Conservation Status		Source					Likelihood of Occurrence	Justification
			State	Federal	NM	PMST	DBCA	DBCA 15 yrs	Field Survey		
											(Williams, Williams and Lundstrom, 1998)
Lycaenidae	<i>Ogyris subterrestris petrina</i>	Arid Bronze Azure Butterfly	CR (not WAM)	CR (not WAM)	x	x	x			Medium	No recent records (Department of Biodiversity Conservation and Attractions, 2021c). Suitable habitat present (smooth bark <i>Eucalyptus</i> sp.) (Department of Biodiversity Conservation and Attractions, 2020b).

5 Discussion

5.1 Flora and Vegetation

5.1.1 Flora Composition

The suite of flora taxa recorded during the survey is considered typical for the area (Beard 1976) and aligns with the database search results obtained and with previous surveys conducted around the Survey Area.

5.1.2 Survey Adequacy

The flora and vegetation survey effort was in accordance with the scope of works, and appropriate for a detailed flora and vegetation survey in the Eastern Goldfields region.

The Survey Area was sampled with 74 quadrats, 22 relevés and an additional 91 mapping notes. Of the 10 vegetation types defined, four were not sampled with three flora sites. Vegetation types Ec and EooEiiDs were sampled with two sites only due to limited distribution of the vegetation (<1% each). Vegetation types Sl, and Lake were sampled only once due to the landforms (clay and lake) being very limited in distribution.

The inventory of vascular flora was compiled using site data and opportunistic observations made while traversing between sites and during targeted searching within the Survey Area. The entire Survey Area was not systematically searched, and therefore additional flora taxa, and records of conservation significant flora and weed species may be recorded with additional survey effort.

5.1.3 Flora of Conservation Significance

No Threatened flora species pursuant to the EPBC Act 1999 and/or gazetted as Threatened Flora pursuant to the BC Act 2016 were recorded within the Survey Area.

Ten individuals of *Austrostipa blackii* (P3) were recorded from quadrat C1Q01, within vegetation type AcEoaDI. This vegetation type was sampled heavily, and no other records of the taxon were recorded. With more intensive surveys additional individuals may be recorded, as this vegetation type and landform (rocky hills) cover 13.1% of the Survey Area.

5.1.4 Likelihood of Occurrence

Of the 90 Priority flora identified by the database searches, only one was recorded from the Survey Area (*Austrostipa blackii* (P3)). Of the remaining 89 taxa, two were considered to retain a high likelihood of occurrence following the survey:

- *Acacia websteri* (P1): A tall shrub or tree is known only from the Bencubbin and Coolgardie-Kambalda areas, and grows mostly in red loam, sand, and clay in drainage depressions among shrubland and scrub. The habitat for this taxon is present within the Survey Area, and the closest location is <1 km from the survey boundary.

- *Eremophila veronica* (P3): A record of this erect shrub to 40cm high (80cm wide) was located 500 m from the Survey Area, in vegetation similar in appearance to that within the survey boundary. The record was noted as growing with *E. salubris*, *A. burkittii*, and *Cylindropuntia tunicata*.

A further 18 taxa were considered to have a medium likelihood of occurrence due to presence of habitat and records within 15 km of the Survey Area boundary. The remaining taxa were considered to have a low likelihood of occurrence.

5.1.5 Introduced Flora

Thirteen weed species were recorded in the Survey Area, of which one, **Opuntia stricta* (Common Prickly Pear), is a Declared Pest and listed as a WoNS.

Weed diversity and population abundance are consistent with the region and land use (mining operations, broadacre cropping and sheep farming).

Weed species richness and abundance was greatest in previously cleared areas such as old drill pads/ excavation works and associated tracks. Weed species, particularly **Opuntia stricta* were present in relatively high abundance near Great Eastern Highway. The publicly accessible semi-pool 'Coolgardie Gorge' contained dense weed numbers along the edge and in the immediate surrounds.

5.1.6 Unconfirmed Flora

Eighteen specimens collected during the field survey were sterile and could not be confidently identified beyond genus level. None of these were analogous to Priority flora taxa identified by the database searches.

5.1.7 Vegetation Types

No vegetation representative of any TECs or PECs was recorded in the Survey Area.

Mapping reliability across the survey area was high, and the majority of the Survey Area was traversed on foot. The plains through the majority of the Survey Area showed a mosaic of different *Eucalyptus* spp. that could not be easily separated via quadrating or aerial imagery. However, *E. salmonophloia* was scattered across the majority of the plains, with *E. celastroides*, *E. clelandiorum*, and *E. torquate* at varying densities across the plains. This is common across the region, as underlying geology affects the vegetation on a fine scale. The Survey Area was dominated by these plains, along with rocky hills dominated by *Acacia collegialis* tall shrubs. While this vegetation type AcEoaDI rocky hills did support *Austropstipa blackii* (P3), only one population was recorded through 21 quadrats in the vegetation. As such it was not considered to be a distinctive habitat of the taxon, and not considered to be vegetation of local significance.

Coolgardie Gorge is a natural semi-pool landform present within the Survey Area, which is also a public recreation area, as such it contains many introduced (weed) taxa. The landform is limited in size, with the vegetation unique and restricted to a small radius around the semi-pool.

Numerous patches within the Survey Area were considered either recently or historically cleared. Some of these historically cleared areas contained some regrowing native and weed taxa, however, were still considered to be cleared as they were predominantly bare, and any native taxa did not represent the surrounding vegetation. One small area of claypan vegetation was recorded and sampled with a single quadrat. After recent rains, this claypan landform may contain an increased load of annual and cryptic taxa.

5.2 Vertebrate Fauna

5.2.1 Fauna Habitat

The five broad fauna habitats identified within the Survey Area are typical of the Coolgardie bioregion and consistent with habitats identified by previous studies in the region (Terrestrial Ecosystems, 2018; Phoenix Environmental Sciences, 2019; Strategen Environmental, 2019; Biostat Pty Ltd, 2020). Of the habitats within the Survey Area, the *Eucalyptus* Woodlands, *Acacia* Shrublands and Open Water habitats, particularly the semi-permanent pool at Coolgardie Gorge, represent the most value to the overall fauna assemblage within the Survey Area. The *Eucalyptus* Woodlands and the *Acacia* Shrubland habitats are valuable due to the variety of micro habitats present. Peeling bark, woody debris, leaf litter and hollow logs were observed throughout these habitat types. The dense understorey observed in these habitat types may provide refuge and shelter for a variety of mammals, reptiles, and birds. Specifically, these habitat types may provide suitable breeding and nesting habitat for the Malleefowl (*Leipoa ocellata*) and foraging habitat for the Western Quoll, Chuditch (*Dasyurus geoffroyi fortis*).

The Open Water habitat provides an important water source for fauna within the Survey Area and is particularly important for water birds, including several significant species. The Open Water habitat at Coolgardie Gorge is surrounded by vegetation that provides potential foraging, roosting, and nesting habitat for a wide variety of fauna taxa. Old mine pits throughout the Survey Area may have water present, however it will be of less value to most fauna species due to a lack of surrounding vegetation.

The Rocky Slopes habitat contains crevices that may provide refuge and shelter for small fauna taxa. This habitat type generally lacks the dense understory found in the *Eucalyptus* Woodlands and the *Acacia* Shrubland habitats, making it less suitable for the Malleefowl and its slope reduces its value as Malleefowl nesting habitat.

Cleared Areas provide limited habitat value for fauna; however, some small mammals and reptiles may use the rocky crevices within the steep walls of mining pits and the low isolated shrubs occasionally found in other Cleared Areas for shelter and foraging.

The habitat condition varied widely throughout the Survey Area from highly degraded to high quality. The most common forms of disturbance included previous mining activity, litter, and vehicle tracks. Species with a limited dispersal capability may be restricted by these impacts, whilst movements of birds, larger reptiles, and mammals between surrounding habitats are unlikely to be severely impeded by these disturbances.

5.2.2 Conservation Significant Fauna

5.2.2.1 Recorded

Western Quoll, Chuditch (*Dasyurus geoffroii fortis*), Vulnerable under the BC Act and EPBC Act

The Chuditch inhabits areas dominated by sclerophyll forest or drier woodland, heath and mallee shrubland (Van Dyck and Strahan, 2008). The species is generally highly mobile and uses bush remnants as corridors (Woinarski, Burbidge and Harrison, 2014). The Chuditch is a largely nocturnal animal, feeding on a carnivorous diet of mammals, birds, lizards, and frogs. Although they have been recorded foraging during the day at particular times such as during the breeding season or when cold and wet weather restricts their nocturnal movements (Van Dyck and Strahan, 2008). Most diurnal nesting sites in sclerophyll forest consist of hollow logs or earth burrows, although bandicoot nests and hollow tree bases may be used (Van Dyck and Strahan, 2008). The Chuditch was abundant prior to European settlement, and it is now largely restricted to the south-west of Western Australia, with small numbers in the Midwest, Wheatbelt and South Coast regions (Department of Biodiversity Conservation and Attractions, 2017a).

Chuditch scat was recorded within the Survey Area. Of the habitat types present with the Survey Area the *Eucalyptus* Woodland and *Acacia* Shrubland present the most value to this species. These habitats contain mallee woodland and shrubland, which the Chuditch often inhabits as well hollow logs and tree bases which are used for denning (Van Dyck and Strahan, 2008). The suitability of these habits likely varies with the degree of disturbance which may alter prevalence of both denning sites and availability of prey.

5.2.2.2 High Likelihood

Malleefowl (*Leipoa ocellata*), Vulnerable under the BC Act and EPBC Act

The Malleefowl is found in semi-arid to arid shrublands and low woodlands, especially those dominated by mallee and/or *Acacia*. The species nests in large mounds of dirt and leaf litter up to five metres wide and one metre tall (Menkhorst et al., 2017). Sandy substrates and abundance of leaf litter are required for breeding (Department of the Environment and Energy, 2018). Densities of the birds are generally greatest in areas of higher rainfall and on more fertile soils where there is an abundance of food plants.

The Malleefowl has recently been recorded within 1 km of the Survey Area and is therefore highly likely to use suitable habitat within the Survey Area (Department of Biodiversity Conservation and Attractions, 2022). The *Eucalyptus* Woodland and *Acacia* Shrubland habitats constitute suitable habitat for the taxon. No mounds, tracks, or other signs of the taxon observed during the field survey. When present, these signs are obvious and detectable (National Malleefowl Recovery Team, 2016). While breeding may occur in the Survey Area, it is unlikely that high densities of breeding individuals are present.

Common Sandpiper (*Actitis hypoleucos*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

The Common Sandpiper typically feeds on mollusks and crustaceans as well as a variety of insects. It is a migratory species that uses varied coastal and interior wetlands including narrow muddy edges of billabongs, river pools, mangroves, among rocks and snags, reefs or rocky beaches (Morcombe, 2003). It migrates from mid-northern latitudes of Asia (Menkhorst *et al.*, 2017) and likely breeds in the Russian far east (Pizzey and Knight, 2001).

The Common Sandpiper was not recorded during the current field survey, however it has been recorded within the Survey Area previously (Department of Biodiversity Conservation and Attractions, 2022). The taxon may use the Open Water habitat within the Survey Area, particularly the Coolgardie Gorge for foraging and roosting.

Common Greenshank (*Tringa nebularia*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

The Common Greenshank is a migratory species to Australia that typically breeds in Norway (BirdLife International, 2016). The species is found in a wide variety of wetlands from coastal to freshwater, where it wades in shallow water foraging for prey, often lunging or probing for fish and invertebrates (Pizzey and Knight, 2001; BirdLife International, 2016; Menkhorst *et al.*, 2017).

The Common Greenshank was not recorded during the current field survey, however, it has been recorded within the Survey Area previously (Department of Biodiversity Conservation and Attractions, 2022). The taxon may use the Open Water habitat within the Survey Area, particularly the Coolgardie Gorge for foraging and roosting.

5.2.2.3 Medium Likelihood

Sharp-tailed Sandpiper (*Calidris acuminata*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

The Sharp-tailed Sandpiper is a small-medium migratory wader, nesting in the northern Siberian Tundra and is one of the most common shorebirds to inhabit in fresh and saline wetlands occurring inland in Australia (Menkhorst *et al.*, 2017). In Western Australia there are scattered records of the species occurring along the Nullarbor Plain and the southern areas of the Great Victoria Desert. They are widespread from Cape Arid to Carnarvon, around coastal and subcoastal plains of the Pilbara Region to south-west and east Kimberley (Pizzey and Knight,

2001). It also forages in grasslands and tidal areas. The species typically prefers muddy edges of shallow fresh or brackish wetlands, with inundated or emergent sedges, grass, saltmarsh or other low vegetation (Pizzey and Knight, 2001).

The Sharp-tailed Sandpiper was not recorded during the field survey. The nearest record was 15 km northwest of the Survey Area (Department of Biodiversity Conservation and Attractions, 2022). The taxon may use the Open Water habitat within the Survey Area, particularly the Coolgardie Gorge for foraging and roosting.

Curlew Sandpiper (*Calidris ferruginea*), Critically Endangered, International Agreement under the BC Act and Critically Endangered, Migratory and Marine under the EPBC Act.

The Curlew Sandpiper is a small, slim sandpiper that is most commonly found on large tidal flats, occasionally using brackish and freshwater wetland systems (Menkhorst *et al.*, 2017). They typically forage on invertebrates throughout mudflats and nearby shallow water. The species is migratory and nests in Arctic Siberia, and in Western Australia are widespread around coastal and subcoastal plains from Cape Arid to the south-west Kimberley (Pizzey and Knight, 2001).

The Curlew Sandpiper was not recorded during the field survey. The nearest record was 15 km northwest of the Survey Area (Department of Biodiversity Conservation and Attractions, 2022). The taxon may use the Open Water habitat within the Survey Area, particularly the Coolgardie Gorge for foraging and roosting.

Wood Sandpiper (*Tringa glareola*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

The Wood Sandpiper is a small, thin wader that uses well-vegetated, shallow, freshwater wetlands such as swamps, billabongs, lakes, pools and waterholes (Menkhorst *et al.*, 2017). The species is generally associated with emergent aquatic plants or grass, and they forage on moist or dry mud at the edges of wetlands. The Wood Sandpiper is a migratory species, breeding in the subarctic Siberia with some individuals migrating to Australia (Menkhorst *et al.*, 2017). In Western Australia the species is widespread but scattered in most regions (Pizzey and Knight, 2001).

The Wood Sandpiper was not recorded during the field survey. The nearest record was 30 km northeast of the Survey Area (Department of Biodiversity Conservation and Attractions, 2022). The taxon may use the Open Water habitat within the Survey Area, particularly the Coolgardie Gorge for foraging and roosting.

Glossy Ibis (*Plegadis falcinellus*), International Agreement under the BC Act and Migratory and Marine under the EPBC Act.

The preferred foraging and breeding habitat of the Glossy Ibis includes fresh water marshes at the edges of lakes and rivers, lagoons, flood-plains, wet meadows, swamps, reservoirs, sewage ponds, rice-fields and cultivated areas under irrigation (Marchant and Higgins, 1990; del Hoyo, Elliot and Sargatal, 1992). The Glossy Ibis builds a platform nest of sticks in trees or shrubs above

water and typically nests in colonies (Pizzey and Knight, 2013). The distribution of the Glossy Ibis is generally east of the Kimberley in Western Australia and Eyre Peninsula in South Australia. The species is also known to be patchily distributed in the rest of Western Australia (Department of the Environment and Energy, 2020).

The Glossy Ibis was not recorded during the field survey, however, it has been recorded 27 km northeast of the Survey Area in Kalgoorlie (Department of Biodiversity Conservation and Attractions, 2022). The taxon may use the Open Water habitat within the Survey Area, particularly the Coolgardie Gorge for foraging and roosting.

5.3 Invertebrates

Although outside the scope of this report, two conservation significant invertebrates were identified as potentially occurring within the Survey Area.

Arid Bronze Azure Butterfly (*Ogyris subterrestris petrina*), Critically Endangered under the BC Act and EPBC Act.

Following the extirpation of the only known population in the early 1990s, the Arid Bronze Azure Butterfly (ABAB) was rediscovered at Barbalin Nature Reserve in 2006. The ABAB can only reproduce where nests of the sugar ant, *Camponotus* sp. nr. *terebrans* are present. The ants protect the larvae from predators and in return, the ants feed on secretions produced by the ABAB larvae. Unlike other species in this genus, the ABAB larva do not eat vegetation and are entirely dependent upon the host ant. The ABAB larvae require large ant colonies that are typically found at the base of many species of smooth-barked eucalypts including *Eucalyptus salubris* and *E. salmonophloia* (Department of Biodiversity Conservation and Attractions, 2020b).

The presence of multiple smooth-barked eucalypt species within the known distribution of the host ant *Camponotus* sp. nr. *terebrans* suggests that suitable habitat for the ABAB may occur within the survey area. The presence of potentially suitable ABAB habitat within the range of the host ant requires surveys to determine the presence and density of *Camponotus* sp. nr. *terebrans* colonies (Department of Biodiversity Conservation and Attractions, 2020a, 2020b). Consequently, a targeted survey for host ant colonies was conducted in Dec 2021. The results of this survey are presented in a separate report in preparation (360 Environmental 2022, in prep).

Inland Hairstreak (*Jalmenus aridus*), Priority 1 under the BC Act.

Jalmenus aridus is one of ten currently recognised species in the genus *Jalmenus*, which is found only in Australia. *Jalmenus* species feed openly on the foliage of their respective host plants, and all are attended by specific ant species. *Jalmenus aridus* is attended by the ant species *Froggattella kirbii* (Sands and New, 2002). *Jalmenus aridus* is protected by the ants from predators and parasitoids and the butterfly caterpillars reward the ants with sweet liquid produced in special organs.

The butterfly is only present in its adult form for short periods of time (perhaps only two to three weeks at a given site). Timing of adult emergence is linked to rainfall patterns, plant growth stages and other ecological conditions with most likely flight times from mid-October to mid-November.

The ant, *Froggattella kirbii* is found from near Perth to the East coast of Australia and is associated with two known host plants. One host plant, *Senna nemophila* (*artemisioides*), is found over most of central and eastern WA while the other known host plant, *Acacia tetragonophylla*, is found almost everywhere in WA except the very far north and far south.

During the surveys, it was observed that suitable habitat was present within the Survey Area. Additional targeted surveys will be required to confirm the presence of *Jalmenus aridus* within the Survey Area.

6 Conclusion

Flora and Vegetation

- No Threatened Listed flora species pursuant to the EPBC Act 1999 and/or gazetted as Threatened/Declared Rare Flora pursuant to the BC Act 2016 were recorded
- Ten individuals of *Austrostipa blackii* (P3), were recorded in one quadrat (C1Q01)
- Thirteen introduced species were recorded during the survey, of which one, **Opuntia stricta*, is a Declared Pest and listed as a WoNS
- Ten vegetation types were mapped within the Survey Area, none of which were analogous to conservation significant ecological communities.

Vertebrate Fauna

- Five fauna habitats were mapped, of which the *Eucalyptus* Woodland, *Acacia* Shrubland, and Open Water habitat types represent the most value to conservation significant fauna and overall fauna assemblages. A naturally occurring pool at Coolgardie Gorge was of particularly value to water birds and to the overall fauna assemblage.
- One conservation significant species, the Western Quoll, Chuditch (*Dasyurus geoffroii fortis*), which is listed as Vulnerable under the BC Act and EPBC Act, was recorded within the Survey Area. Chuditch scat was observed during the field survey.
- Seven introduced species were recorded during the survey, Goat (*Capra hircus*), European Cattle (*Bos primigenius taurus*), Dog/Dingo (*Canis familiaris*), Red fox (*Vulpes vulpes*), Horse (*Equus caballus*), Cat (*Felis catus*) and Rabbit (*Oryctolagus cuniculus*).
- One significant fauna species, the Western Quoll, Chuditch (*Dasyurus geoffroii fortis*) which is listed as Vulnerable under the BC Act and EPBC Act, was recorded within the Survey Area based on a scat observed during the field survey.
- Although outside the scope of this report, potential habitat for Arid Bronze Azure Butterfly (*Ogyris subterrestris petrina*), which is listed as Critically Endangered under the BC Act and EPBC Act, and the Inland Hairstreak butterfly (*Jalmenus aridus*), which is listed as Priority 1 by the DBCA, was observed within the Survey Area.

7 Report Disclaimer

This report is produced strictly in accordance with the scope of services set out in the contract or otherwise agreed in accordance with the contract. 360 Environmental makes no representations or warranties in relation to the nature and quality of soil and water other than the visual observation and analytical data in this report.

In the preparation of this report, 360 Environmental has relied upon documents, information, data, and analyses ('client's information') provided by the client and other individuals and entities. In most cases where client's information has been relied upon, such reliance has been indicated in this report. Unless expressly set out in this report, 360 Environmental has not verified that the client's information is accurate, exhaustive, or current and the validity and accuracy of any aspect of the report including, or based upon, any part of the client's information is contingent upon the accuracy, exhaustiveness, and currency of the client's information. 360 Environmental shall not be liable to the client or any other person in connection with any invalid or inaccurate aspect of this report where that invalidity or inaccuracy arose because the client's information was not accurate, exhaustive, and current or arose because of any information or condition that was concealed, withheld, misrepresented, or otherwise not fully disclosed or available to 360 Environmental.

Aspects of this report, including the opinions, conclusions, and recommendations it contains, are based on the results of the investigation, sampling and testing set out in the contract and otherwise in accordance with normal practices and standards. The investigation, sampling and testing are designed to produce results that represent a reasonable interpretation of the general conditions of the site that is the subject of this report. However, due to the characteristics of the site, including natural variations in site conditions, the results of the investigation, sampling and testing may not accurately represent the actual state of the whole site at all points.

It is important to recognise that site conditions, including the extent and concentration of contaminants, can change with time. This is particularly relevant if this report, including the data, opinions, conclusions, and recommendations it contains, are to be used a considerable time after it was prepared. In these circumstances, further investigation of the site may be necessary.

Subject to the terms of the contract between the Client and 360 Environmental Pty Ltd, copying, reproducing, disclosing, or disseminating parts of this report is prohibited (except to the extent required by law) unless the report is produced in its entirety including this page, without the prior written consent of 360 Environmental Pty Ltd.

8 References

- Atlas of Living Australia (2022a) *No Title*. Available at: <https://www.ala.org.au/>.
- Atlas of Living Australia (2022b) *No Title*.
- Beard, J. S. (1976) *Vegetation survey of Western Australia. Western Australia 1: 1 000 000 vegetation series. Design and cartography by Dept. of Geography, University of W.A.*
- Biostat Pty Ltd (2020) *Vegetation Clearing - Fauna Assessment (Davyhurst, Callion, Waihi, Siberia, Riverina Clearance Areas)*.
- BirdLife International (2016) *Tringa nebularia. The IUCN Red List of Threatened Species 2016*. Available at: <https://www.iucnredlist.org/species/22693220/86684205>.
- Botanica Consulting (2020) *Flora and Vegetation Impact Assessment - Medcalf Project*.
- Bureau of Meteorology (2021) *Monthly climate data statistics*. Available at: www.bom.gov.au/climate/data.
- Bureau of Meteorology (2022) *Groundwater Dependent Ecosystems Atlas*.
- Clarke, K. R. and Gorley, R. N. (2006) 'Primer-E v6'. Plymouth, United Kingdom, United Kingdom.
- Cowan, M. (2001) *Coolgardie 3 (COO3 – Eastern Goldfields subregion)*. Available at: https://www.dpaw.wa.gov.au/images/documents/about/science/projects/waaudit/coolgardie03_p156-169.pdf.
- Data WA (2021) *Mining Tenements (DMIRS-003), Mining Tenements (DMIRS-003)*. Available at: <https://catalogue.data.wa.gov.au/dataset/mining-tenements-dmirs-003>.
- Department of Agriculture and Food WA (2012) *Soil-landscape systems of Western Australia - GIS dataset*. Perth, Australia.
- Department of Agriculture Water and the Environment (2020) *Protected Matters Search Tool*. Canberra, Australia. Available at: <http://www.environment.gov.au/webgis-framework/apps/pmst/pmst.jsf>.
- Department of Agriculture Water and the Environment (2021a) *Protected Matters Search Tool*. Canberra, Australia. Available at: <http://www.environment.gov.au/webgis-framework/apps/pmst/pmst.jsf>.
- Department of Agriculture Water and the Environment (2021b) *Weeds of National Significance*. Available at: <https://weeds.org.au/>.
- Department of Biodiversity Conservation and Attractions (2017a) *Fauna Profile - Chuditch Dasyurus geoffroii*. Available at: https://www.dpaw.wa.gov.au/images/documents/plants-animals/animals/animal_profiles/chuditch_fauna_profile.pdf.
- Department of Biodiversity Conservation and Attractions (2017b) *Fauna Profile - Numbat Myrmecobius fasciatus, 2017*. Available at: https://www.dpaw.wa.gov.au/images/documents/plants-animals/animals/animal_profiles/numbat_fauna_profile.pdf.
- Department of Biodiversity Conservation and Attractions (2020a) *Arid bronze azure butterfly (ABAB) survey in Western Australia additional information*.

Department of Biodiversity Conservation and Attractions (2020b) *Guideline for the survey of arid bronze azure butterfly (ABAB) in Western Australia*.

Department of Biodiversity Conservation and Attractions (2021a) *DBCA - Legislated Lands and Waters (DBCA-011) GIS Dataset*. Perth, Australia. Available at:
<https://catalogue.data.wa.gov.au/dataset/dbca-legislated-lands-and-waters>.

Department of Biodiversity Conservation and Attractions (2021b) *NatureMap*. Perth, Western Australia. Available at: <https://naturemap.dpaw.wa.gov.au/>.

Department of Biodiversity Conservation and Attractions (2021c) *Threatened and Priority Fauna database request (custom search)*. Perth, Australia.

Department of Biodiversity Conservation and Attractions (2021d) *Threatened and Priority Flora List (TPFL) database request (custom search)*. Perth, Australia.

Department of Biodiversity Conservation and Attractions (2021e) *Western Australia Herbarium Flora Database (custom search)*. Perth, Australia.

Department of Biodiversity Conservation and Attractions (2022) *Threatened and Priority Ecological Communities database request (custom search)*.

Department of Primary Industries and Regional Development (2018) *Soil Landscape Mapping - Systems (DPIRD-064) - GIS Dataset*. Perth, Australia. Available at:
<https://catalogue.data.wa.gov.au/dataset/soil-landscape-mapping-systems>.

Department of Primary Industries and Regional Development (2021) *Declared plants*. Available at: <https://www.agric.wa.gov.au/organisms>.

Department of Sustainability Environment Population and Communities (1999) *Survey Guidelines for Australia's Threatened Mammals*. Canberra, Australia. Available at:
<http://www.environment.gov.au/system/files/resources/b1c6b237-12d9-4071-a26e-ee816caa2b39/files/survey-guidelines-mammals.pdf>.

Department of Sustainability Environment Water Population and Communities (2011) *Survey guidelines for Australia's threatened reptiles: Guidelines for detecting reptiles listed as threatened under the EPBC Act*. Canberra, Australia. Available at:
<http://www.environment.gov.au/resource/survey-guidelines-australias-threatened-reptiles-guidelines-detecting-reptiles-listed>.

Department of Sustainability Environment Water Population and Communities (2012) *EPBC Act Referral guidelines for three threatened black cockatoo species: Carnaby's cockatoo, Baudin's cockatoo and Forest red-tailed black cockatoo*. Canberra, Australia.

Department of the Environment (2013) *Matters of National Environmental Significance: Significant impact guidelines 1.1*. Canberra, Australia. Available at:
http://www.environment.gov.au/system/files/resources/42f84df4-720b-4dcf-b262-48679a3aba58/files/nes-guidelines_1.pdf (Accessed: 24 July 2018).

Department of the Environment and Energy (2018) *Leipoa ocellata - Malleefowl*. Available at:
http://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon_id=934.

Department of the Environment and Energy (2020) *Plegadis falcinellus in Species Profile and Threats (SPRAT) database*. Available at: https://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon_id=991.

Department of the Environment Water Heritage and the Arts (2010) *Survey guidelines for*

Australia's threatened birds: Guidelines for detecting birds listed as threatened under the EPBC Act. Canberra, Australia. Available at:

<http://www.environment.gov.au/system/files/resources/107052eb-2041-45b9-9296-b5f514493ae0/files/survey-guidelines-birds-april-2017.pdf> (Accessed: 24 July 2018).

Department of Water and Environmental Regulation (2018) *Hydrography, Linear (Hierarchy) (DWER-031) - GIS Dataset*. Perth, Australia: Landgate. Available at:

<https://catalogue.data.wa.gov.au/dataset/hydrography-linear-hierarchy>.

Department of Water and Environmental Regulation (2020) *Clearing Regulations - Environmentally Sensitive Areas (DWER-046)*. Available at:

<https://catalogue.data.wa.gov.au/dataset/clearing-regulations-environmentally-sensitive-areas-dwer-046>.

Van Dyck, S. and Strahan, R. (2008) *The mammals of Australia*. 3rd edn. Sydney, Australia: New Holland Publishers.

Environmental Protection Authority (2016) *Technical Guidance - Flora and Vegetation surveys for Environmental Impact Assessment*. Perth, Australia. Available at:

[http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EPA Technical Guidance - Flora and Vegetation survey_Dec13.pdf](http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/EPA%20Technical%20Guidance%20-%20Flora%20and%20Vegetation%20survey_Dec13.pdf).

Environmental Protection Authority (2020) *Technical Guidance - Terrestrial vertebrate fauna surveys for environmental impact assessment*. Perth, Australia. Available at:

<https://www.epa.wa.gov.au/policies-guidance/technical-guidance-terrestrial-vertebrate-fauna-surveys-environmental-impact>.

Government of Western Australia (2019) *2018 Statewide Vegetation Statistics - Full Report*.

Available at: <https://catalogue.data.wa.gov.au/dataset/dbca-statewide-vegetation-statistics/resource/b7bd60c2-bff6-4637-b213-ae4706412c7>.

del Hoyo, J., Elliot, A. and Sargatal, J. (). (1992) *Handbook of the Birds of the World. 1. Ostrich to Ducks*. Spain: Lynx Edicions.

Keighery, B. (1994) *Bushland Plant Survey: A Guide to Plant Community Survey for the Community*. Floreat, Western Australia.

Marchant, S. and Higgins, P. J. (1990) *Handbook of Australian, New Zealand and Antarctic Birds. Volume 1 - Ratites to Ducks*. Melbourne: Oxford University Press.

Menkhorst, P. et al. (2017) *The Australian bird guide*. Australia: CSIRO Publishing.

Morcombe, M. (2003) *Field Guide to Australian Birds*. Archerfield: Steve Parish Publishing Pty Ltd.

National Malleefowl Recovery Team (2016) *NATIONAL MALLEEFOWL Monitoring Manual*.

Native Vegetation Solutions (2018) *Reconnaissance Flora and Vegetation Survey of Lot 500 Great Eastern Highway Kalgoorlie*.

Native Vegetation Solutions (2019) *Reconnaissance Flora and Vegetation Survey for the Mt Marion Project Area*.

Native Vegetation Solutions (2020) *Reconnaissance Flora and Vegetation Survey of the Spargos Project - October 2020*.

Phoenix Environmental Sciences (2019) *Fauna survey for Mungari Gold Operations Cutters*

Ridge Project.

Pizzey, G. and Knight, F. (2001) *Field Guide to Birds of Australia*. Pymble, Australia: Harpercollins Australia.

Pizzey and Knight (2013) *Birds of Australia, Digital Edition*. Melbourne, Australia: Gibbon Multimedia.

Sands, D. P. A. and New, T. R. (2002) *The Action Plan for Butterflies*. Canberra: Environment Australia.

Shepherd, D. P., Beeston, G. R. and Hopkins, A. J. M. (2002) *Native Vegetation in Western Australia Technical Report 249*. Perth, Australia.

Strategen Environmental (2019) *Coolgardie Landfill - Flora, vegetation and fauna habitat assessment*.

Terrestrial Ecosystems (2018) *Level 1 Vertebrate Fauna Risk Assessment for Lot 500 Kalgoorlie West*.

Western Australian Herbarium (2021) *FloraBase - The Western Australian Flora*. Perth, Australia. Available at: <https://florabase.dpaw.wa.gov.au>.


Western Australian Herbarium (2022) *FloraBase - The Western Australian Flora*. Available at: <https://florabase.dpaw.wa.gov.au/>.

Western Australian Museum (2019) *Checklist of the Terrestrial Vertebrate Fauna of Western Australia / Western Australian Museum*. Perth, Australia. Available at: <http://museum.wa.gov.au/research/departments/terrestrial-zoology/checklist-terrestrial-vertebrate-fauna-western-australia> (Accessed: 3 August 2018).

Williams, M., Williams, A. and Lundstrom, T. (1998) 'Jewels of the West', *Landscape*, pp. 49–53.

Woinarski, J., Burbidge, A. and Harrison, P. (2014) *The action plan for Australian mammals 2012*. CSIRO Publishing.

Figures



Legend

- Survey Area
- State Road



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
 - OTHER DATA SOURCED LANDGATE 2022
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
 (© Western Australian Land Information Authority 2022)

360
 environmental
 a 10 Demondravy St West Leederville, 6007 WA
 t (08) 9384 8380
 f (08) 9381 2260
 www.360environmental.com.au

PROJECT ID: 4794
 DATE: 21/04/2022


HORIZONTAL DATUM AND PROJECTION
 GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 1
 Survey Area



Legend

Soil Land System


- 266BB5
- 266Mx43

Hydrography

- Watercourse - minor
- Drain - major
- Lake
- Water Reservoir
- Survey Area

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360 environmental
1000 938 8280
1000 938 12260
www.360environmental.com.au

PROJECT ID: 4794 DATE: 21/04/2022


HORIZONTAL DATUM AND PROJECTION: GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 2
Soil Landscapes and Land Systems and Hydrography



Legend


 Survey Area

Broad Vegetation Types

- BOORABBIN_8
- COOLGARDIE_123
- COOLGARDIE_1294
- COOLGARDIE_522
- COOLGARDIE_540
- COOLGARDIE_9
- COOLGARDIE_936

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
 - OTHER DATA SOURCED LANDGATE 2022
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
 (© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360
 environmental
 www.360environmental.com.au

PROJECT ID: 4794
 DATE: 29/04/2022


HORIZONTAL DATUM AND PROJECTION
 GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 3
 Broad Vegetation Types




Legend

- Survey Area
- DBCA Managed Lands and Waters
- DBCA Managed Land

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
 - OTHER DATA SOURCED LANDGATE 2022
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
 (© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360
 environmental
 www.360environmental.com.au

PROJECT ID: 4794
 DATE: 21/04/2022


HORIZONTAL DATUM AND PROJECTION
 GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 4
 Conservation Areas



Legend


- Survey Area
- GPS Tracks

Sample Sites

- Quadrat
- ▲ Relieve

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360 environmental
1000 9384 8380
1000 9381 2260
www.360environmental.com.au

PROJECT ID: 4794 DATE: 29/04/2022


HORIZONTAL DATUM AND PROJECTION: GDA 1994 MGA Zone 51

CREATED: CL CHECKED: NW APPROVED: NW REVISION: 0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 5a
Survey Effort



Legend


- Survey Area
- GPS Tracks

Sample Sites

- Quadrat
- ▲ Relieve

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360 environmental
1000 1000 1000
1000 1000 1000
www.360environmental.com.au

PROJECT ID: 4794 DATE: 29/04/2022


HORIZONTAL DATUM AND PROJECTION: GDA 1994 MGA Zone 51


CREATED: CL CHECKED: NW APPROVED: NW REVISION: 0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 5b
Survey Effort






Legend

- Survey Area ● *Austrosipa blackii* (P3)
- Vegetation Types**
- AcEoaDI
 - Ec
 - EgAhsat
 - EooEiiDs
 - EsEiiAv
 - EspEiiSaa
 - EtEaEpa
 - SI
 - Cleared

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360 environmental
1000 938 8280
1000 938 12260
www.360environmental.com.au

PROJECT ID: 4794 DATE: 13/05/2022


HORIZONTAL DATUM AND PROJECTION: GDA 1994 MGA Zone 51

CREATED: CL CHECKED: NW APPROVED: NW REVISION: 0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 7a
Vegetation Types




Legend

- Survey Area
- Vegetation Types**
- AcEoaDI
- EgAhSaf
- EooEiIDs
- ErMhOm
- EsEiiAv
- EspEiiSaa
- EtEaEpa
- Lake
- Cleared

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
 - OTHER DATA SOURCED LANDGATE 2022
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
 (© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360
 environmental
 10 Dennington St West Leederville, 6007 WA
 t (08) 9384 8380
 f (08) 9381 2260
 www.360environmental.com.au

PROJECT ID: 4794 DATE: 13/05/2022


HORIZONTAL DATUM AND PROJECTION: GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 7b
Vegetation Types



Legend

Weed Species

- *Agave americana*
- *Carrichtera annua*
- *Centaurea melitensis*
- *Crassula ovata*
- *Nicotiana glauca*
- *Oligocarpus calendulaceus*
- *Rumex vesicarius*
- *Salvia verbenaca*
- *Schinus molle* var. *areira*


Survey Area

Vegetation Condition

- Excellent
- Very Good
- Good
- Completely Degraded

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360
environmental
www.360environmental.com.au

PROJECT ID
4794

DATE
29/04/2022


HORIZONTAL DATUM AND PROJECTION
GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 8a
Vegetation Condition



Legend

Weed Species


- *Asphodelus fistulosus*
- *Garrichtera annua*
- *Centaurea melitensis*
- *Euphorbia drummondii*
- *Heliotropium europaeum*
- *Oligocarpus calendulaceus*
- *Opuntia stricta*
- *Salvia verbenaca*
- *Schinus molle* var. *areira*
- *Sonchus asper*

Survey Area

- Survey Area
- Excellent
- Very Good
- Good
- Completely Degraded

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360
environmental
www.360environmental.com.au

10 Demondrery St West Leederville, 6007 WA
t (08) 9384 8380
f (08) 9381 2260

PROJECT ID
4794

DATE
29/04/2022


HORIZONTAL DATUM AND PROJECTION
GDA 1994 MGA Zone 51

CREATED CL CHECKED NW APPROVED NW REVISION 0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 8b
Vegetation Condition




Legend

 Survey Area


Threatened and Priority Fauna Records

- | | |
|---|---|
| ● A Fairy Shrimp (Carnarvon to Kalbarrie) (P3) | ● Glossy Ibis (MI) |
| ● And Bronze Azure Butterfly (CR) | ● Grey-tailed Tattler (P4) |
| ● Carnaby's Cockatoo (EN) | ● Hooded plover, hooded dotterel (P4) |
| ● Chuditch, Western Quoll (VU) | ● Inland Hairstreak, Desert Blue Butterfly (P1) |
| ● Common Greenshank, Greenshank (MI) | ● Malleefowl (VU) |
| ● Common Sandpiper (MI) | ● Red-necked Stint (MI) |
| ● Curlew Sandpiper (CR) | ● Sanderling (MI) |
| | ● Sharp-tailed Sandpiper (MI) |
| | ● Wood Sandpiper (MI) |

SLIP ENABLER



360 environmental ● ● ●		10 Demondweg St West Leederville, 6007 WA T (08) 9384 8380 F (08) 9381 2260 www.360environmental.com.au	
PROJECT ID		DATE	
4794		21/04/2022	
HORIZONTAL DATUM AND PROJECTION GDA 1994 MGA Zone 51			
CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0
Focus Mineral Limited			
CNX Three Mile Hill Coolgardie Gold Project			
Figure 9 DBCA Threatened and Priority Fauna Records			




Legend

- Survey Area
- Habitat Assessment Location
- Fauna Habitat**
 - Acacia Shrubland
 - Eucalyptus Woodland
 - Rocky Slopes
 - Cleared

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
 - OTHER DATA SOURCED LANDGATE 2022
 - AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
 (© Western Australian Land Information Authority 2022)



- NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360 environmental
 10 Denison St West Leederville, 6007 WA
 t (08) 9384 8380
 f (08) 9381 2260
 www.360environmental.com.au

PROJECT ID: 4794
 DATE: 29/04/2022


HORIZONTAL DATUM AND PROJECTION
 GDA 1994 MGA Zone 51

CREATED	CHECKED	APPROVED	REVISION
CL	NW	NW	0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 10a
Fauna Habitat




Legend

- Survey Area
- Habitat Assessment Location
- Fauna Habitat**
- Acacia Shrubland
- Eucalyptus Woodland
- Open Water
- Rocky Slopes
- Cleared

SLIP ENABLER

- LOCALITY MAP SOURCED FROM LANDGATE 2022
- OTHER DATA SOURCED LANDGATE 2022
- AERIAL PHOTOGRAPHY SOURCED LANDGATE 2022
(© Western Australian Land Information Authority 2022)



NOTE THAT POSITION ERRORS CAN BE >5M IN SOME AREAS

LOCALITY MAP



360 environmental
1000 1000 1000
1000 1000 1000
www.360environmental.com.au

PROJECT ID: 4794
DATE: 29/04/2022

HORIZONTAL DATUM AND PROJECTION
GDA 1994 MGA Zone 51

CREATED: CL
CHECKED: NW
APPROVED: NW
REVISION: 0

Focus Mineral Limited

CNX Three Mile Hill Coolgardie Gold Project

Figure 10b
Fauna Habitat

Appendices

Appendix A

Flora and Fauna Literature Review

Appendix A: Fauna Literature Review

Report	Distance to current Survey Area	Survey timing	Survey effort	Recorded conservation significant fauna	Fauna habitats
Coolgardie landfill Flora, Vegetation and Fauna Habitat Assessment (Stategen Environmental 2019)	6.4 kms ESE of Survey Area	18 December 2018	Describe fauna habitat within the Survey Area	None	Open mallee woodland to mallee woodland.
Fauna survey for Mungari Gold Operations Cutters Ridge Project (Phoenix Environmental Sciences, 2019)	40 km North of Survey Area	September 2018	Level 1 Fauna Targeted Malleefowl	Malleefowl (<i>Leipoa ocellata</i>)	Open Eucalypt woodland, shrubland, and salt lake, in addition to a small portion of existing cleared areas.
Level 1 Vertebrate Fauna Risk Assessment for Lot 500 Kalgoorlie West (Terrestrial Ecosystems, 2018)	24 km NE of Survey Area	2018	Level 1 Vertebrate Fauna	None	Open eucalypt woodland with a mixed understory of scattered shrubs and chenopods
Vegetation Clearing - Fauna Assessment (Biostat Pty Ltd, 2020)	40 km N of the Survey Area	December 2019	Level 1 Reconnaissance	None	Open woodland: Acacia, Casuarina, and Eucalypt. Chenopod shrubland, Acacia shrubland. Eucalyptus sp. mallee over spinifex. Stoney/rocky hills Drainage lines Mulga woodland

Appendix B

Flora and Fauna Database Searches

Appendix B: DBCA Threatened and Priority Fauna

CLASS	SCI_NAME	COM_NAME	WA_status	EPBCstatus	YEAR	SOURCE	CERTAINTY	OBS_METHOD	OBS_TYPE	COUNT	LOCALITY	SITE
Actitis hypoleucos	Common Sandpiper	BIRD	MI	MI	2014	BIRDATA				0	coolgardie gorge	coolgardie gorge
Actitis hypoleucos	Common Sandpiper	BIRD	MI	MI	2011	BIRDATA				0	The Gorge (Coolgardie0	The Gorge (Coolgardie0
Actitis hypoleucos	Common Sandpiper	BIRD	MI	MI	2013	BIRDATA				0	Coolgardie Gorge wetland	Coolgardie Gorge wetland
Tringa nebularia	Common greenshank, greenshank	BIRD	MI	MI	2013	BIRDATA				0	Coolgardie Gorge wetland	Coolgardie Gorge wetland
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	2012	BIRDATA				0	Silver Lake	Silver Lake
Calidris alba	Sanderling	BIRD	MI	MI	2016	BIRDATA				0	Hannan Lake	Hannan Lake
Calyptrorhynchus lat	Carnaby's cockatoo	BIRD	EN	EN	2016	BIRDATA				0	367 Collins	367 Collins
Calyptrorhynchus lat	Carnaby's cockatoo	BIRD	EN	EN	2016	BIRDATA				0	Cape Lilac on alley	Cape Lilac on alley
Calyptrorhynchus lat	Carnaby's cockatoo	BIRD	EN	EN	2017	BIRDATA				0	Piccadilly St West	Piccadilly St West
Tringa brevipes	Grey-tailed tattler	BIRD	P4	MI	2017	BIRDATA				0	Lake Douglas	Lake Douglas
Tringa nebularia	Common greenshank, greenshank	BIRD	MI	MI	1980	BIRDATLAS1				0	KANOWNA	KANOWNA
Tringa glareola	Wood sandpiper	BIRD	MI	MI	1980	BIRDATLAS1				0	KARLKURLA	KARLKURLA
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	1980	BIRDATLAS1				0	KARLKURLA	KARLKURLA
Thinornis rubricollis	Hooded plover, hooded dotterel	BIRD	P4		1980	BIRDATLAS1				0		
Tringa nebularia	Common greenshank, greenshank	BIRD	MI	MI	1980	BIRDATLAS1				0	KANOWNA	KANOWNA
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	1980	BIRDATLAS1				0	FEYSVILLE	FEYSVILLE
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	1981	BIRDATLAS1				0	KARLKURLA	KARLKURLA
Plegadis falcinellus	Glossy ibis	BIRD	MI	MI	1981	BIRDATLAS1				0	KARLKURLA	KARLKURLA
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	1981	BIRDATLAS1				0	KANOWNA	KANOWNA
Tringa nebularia	Common greenshank, greenshank	BIRD	MI	MI	2001	BIRDATLAS2				0	Kopai Lake	Kopai Lake
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	2001	BIRDATLAS2				0	Kopai Lake	Kopai Lake
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	2001	BIRDATLAS2				0	Kopai Lake	Kopai Lake
Tringa glareola	Wood sandpiper	BIRD	MI	MI	2005	BIRDATLAS2				0	Kaloorlie Sewerage overflow ponds	Kaloorlie Sewerage overflow ponds
Tringa glareola	Wood sandpiper	BIRD	MI	MI	2005	BIRDATLAS2				0	Kaloorlie Sewerage outlet	Kaloorlie Sewerage outlet
Tringa nebularia	Common greenshank, greenshank	BIRD	MI	MI	2006	BIRDATLAS2				0	Young River Station Lake	Young River Station Lake
Calidris ferruginea	curlew sandpiper	BIRD	CR	CR	2006	BIRDATLAS2				0	Young River Station Lake	Young River Station Lake
Calidris ruficollis	Red-necked stint	BIRD	MI	MI	2006	BIRDATLAS2				0	Young River Station Lake	Young River Station Lake
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	2006	BIRDATLAS2				0	Young River Station Lake	Young River Station Lake
Tringa nebularia	Common greenshank, greenshank	BIRD	MI	MI	1999	BIRDATLAS2				0	Kurnalpi Road	Kurnalpi Road
Calidris ferruginea	curlew sandpiper	BIRD	CR	CR	1999	BIRDATLAS2				0	Kurnalpi Road	Kurnalpi Road
Calidris acuminata	Sharp-tailed sandpiper	BIRD	MI	MI	1999	BIRDATLAS2				0	Kurnalpi Road	Kurnalpi Road
Leipoa ocellata	Malleefowl	BIRD	VU	VU	2012	FAUNASURVEY	Certain	Survey	Unknown	1	FEYSVILLE	Kaloorlie, Goldfields, Mt Martin
Leipoa ocellata	Malleefowl	BIRD	VU	VU	2012	FAUNASURVEY	Certain	Survey	Unknown	1	FEYSVILLE	Kaloorlie Region, Goldfields, Jubilee mine
Leipoa ocellata	Malleefowl	BIRD	VU	VU	2012	FAUNASURVEY	Certain	Survey	Unknown	1	FEYSVILLE	Kaloorlie Region, Goldfields, Jubilee mine
Leipoa ocellata	Malleefowl	BIRD	VU	VU	2012	FAUNASURVEY	Certain	Survey	Unknown	1	FEYSVILLE	Kaloorlie Region, Goldfields, Jubilee mine
Leipoa ocellata	Malleefowl	BIRD	VU	VU	2012	FAUNASURVEY	Certain	Survey	Unknown	1	FEYSVILLE	Kaloorlie Region, Goldfields, Jubilee mine
Leipoa ocellata	Malleefowl	BIRD	VU	VU	2012	FAUNASURVEY	Certain	Survey	Unknown	1	BULONG	Hampton Hill, Bulong Mining Lease M25/333
Dasyurus geoffroi	chuditch, western quoll	MAMMAL	VU	VU	1974	TFAUNA	Certain	Opportunistic sighting	Caught or trapped	1	Kambalda East	1 mile N of Kambalda East townsite, at edge of Lake Lefroy in sandy mulga country
Leipoa ocellata	malleefowl	BIRD	VU	VU	1994	TFAUNA	Certain	Opportunistic sighting	Secondary sign	0	Bullabulling	one active nest at Bullabulling No. 8 Pumping station.
Leipoa ocellata	malleefowl	BIRD	VU	VU	1995	TFAUNA	Moderately certain	Opportunistic sighting	Day sighting	1	Londonderry	Yerilla Sandalwood Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2000	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Mount Burges	access road to Kundana Mining Lease - "30km NW (10km W & 22km N) of Kalgoorlie"
Branchinella dentic	a fairy shrimp (Carnavon to Kalgoorlie)	INVERTEBRATE	P3		1937	TFAUNA	Certain	Historical (written)	Caught or trapped	0	Gidgi Lake	Gidgi Lake, N of Kalgoorlie
Thinornis rubricollis	hooded plover, hooded dotterel	BIRD	P4		1992	TFAUNA	Certain	Survey	Sighting	0	Arrow Lake	Arrow Lake
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1989	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas, 12 km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1991	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas, 12 km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1982	TFAUNA	Certain	Survey	Caught or trapped	2	Lake Douglas	Lake Douglas, 12 km SW of Kalgoorlie

CLASS	SCI_NAME	COM_NAME	WA_status	EPBCstatus	YEAR	SOURCE	CERTAINTY	OBS_METHOD	OBS_TYPE	COUNT	LOCALITY	SITE
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1986	TFAUNA	Certain	Survey	Caught or trapped	15	Lake Douglas	Lake Douglas, 12 km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1986	TFAUNA	Certain	Survey	Caught or trapped	4	Lake Douglas	Lake Douglas, 12 km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1991	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas, Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1987	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1987	TFAUNA	Certain	Survey	Caught or trapped	8	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1987	TFAUNA	Certain	Survey	Caught or trapped	2	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1989	TFAUNA	Certain	Survey	Caught or trapped	2	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1989	TFAUNA	Certain	Survey	Caught or trapped	2	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1989	TFAUNA	Certain	Survey	Caught or trapped	3	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1989	TFAUNA	Certain	Survey	Caught or trapped	6	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1991	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1991	TFAUNA	Certain	Survey	Caught or trapped	4	Lake Douglas	Lake Douglas, 12km SW of Kalgoorlie
Leipoa ocellata	malleefowl	BIRD	VU	VU	1985	TFAUNA	Certain	Opportunistic sighting	Sighting	2	Jaurdie Hills	Jaurdie Hills
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1911	TFAUNA	Certain	Historical (written)	Caught or trapped	1	Kalgoorlie	SW Australia, Kalgoorlie District
Jalmenus aridus	inland hairstreak, desert blue butterfly	INVERTEBRATE	P1		1985	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas 12 km SW of Kalgoorlie
Jalmenus aridus	inland hairstreak, desert blue butterfly	INVERTEBRATE	P1		1989	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas 12 km SW of Kalgoorlie
Jalmenus aridus	inland hairstreak, desert blue butterfly	INVERTEBRATE	P1		1986	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas 12 km SW of Kalgoorlie
Jalmenus aridus	inland hairstreak, desert blue butterfly	INVERTEBRATE	P1		1986	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas 12 km SW of Kalgoorlie
Ogyris subterrestris	arid bronze azure butterfly	INVERTEBRATE	CR	CR	1985	TFAUNA	Certain	Survey	Caught or trapped	1	Lake Douglas	Lake Douglas 12 km SW of Kalgoorlie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2006	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Bullabulling	Bullabulling
Leipoa ocellata	malleefowl	BIRD	VU	VU	2007	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Bullabulling	Bullabulling
Leipoa ocellata	malleefowl	BIRD	VU	VU	2008	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Bullabulling	Bullabulling pastoral lease, 10km west of the Bullabulling Pub and 2 kms south of the highway
Jalmenus aridus	inland hairstreak, desert blue butterfly	INVERTEBRATE	P1		1997	TFAUNA	Certain	Survey	Caught or trapped	1	Karramindie	Lake Douglas (Douglas Lake)
Leipoa ocellata	malleefowl	BIRD	VU	VU	2009	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Mount Burges	North of Mount Burges
Leipoa ocellata	malleefowl	BIRD	VU	VU	2009	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Bullabulling	Bullabulling, road from Bullabulling to Stewart
Leipoa ocellata	malleefowl	BIRD	VU	VU	2010	TFAUNA	Certain	Opportunistic sighting	Sighting	1		Vic Rock Road
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Opportunistic sighting	Day sighting	1		Yallari Timber Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Opportunistic sighting	Secondary sign	0		Yallari Timber Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Opportunistic sighting	Day sighting	1		
Leipoa ocellata	malleefowl	BIRD	VU	VU	2009	TFAUNA	Certain	Opportunistic sighting	Day sighting	1		on road from Bullabulling to Stuart sighting, off Great Eastern Highway
Leipoa ocellata	malleefowl	BIRD	VU	VU	2009	TFAUNA	Certain	Opportunistic sighting	Day sighting	1		north of Mt Burgess
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Survey	Caught or trapped	1		gravel area just of side of the road at Yarri and Barick Kanowna Belle access road intersection.
Leipoa ocellata	malleefowl	BIRD	VU	VU	2012	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Coolgardie	Just off cave hill road in Widgeemooltha close to one of Focus Minerals small operations.
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Dead	1	Bullabulling	48km West of Coolgardie, No 8 pump Denardi Station, Great Eastern Highway
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Moderately certain	Opportunistic sighting	Day sighting	1	Bullabulling	~10km east if Bullabulling (40km east of Coolgardie) on Great Eastern Highway
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Moderately certain	Opportunistic sighting	Night sighting	1	Feysville	Out the front of Pevnatty Crib room on active haul road. A highly disturbed area.
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Secondary sign	0	Schahil Timber reserve	Schahil Timber reserve

CLASS	SCI_NAME	COM_NAME	WA_status	EPBCstatus	YEAR	SOURCE	CERTAINTY	OBS_METHOD	OBS_TYPE	COUNT	LOCALITY	SITE
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Burra Rock Conservation Reserve	Carpark at Burra Rock. Mound found within adjacent bush habitat.
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Ora Banda	500m before Cawse Nickel turn off on Broad Arrow-Ora Banda road, heading towards Ora Banda
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Kanowna	~5km before the Mt Pleasant Office turn off on Mt Pleasant Rd, off Menzies Highway
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Dead	1	Kanowna	3.1km north of Paddington Mill (35km north of Kalgoorlie) on Menzies Highway
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Day sighting	1		3.2km east of cawse nickel turn off on the Ora Banda - Broad Arrow Rd
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Moderately certain	Opportunistic sighting	Day sighting	1	Feysville	Woolibar Station, just south of where Woolibar creek crosses the Goldfields Highway.
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Dead	1	Kanowna	~4.45km east of Cawse Nickel turn off on the Broad Arrow- Ora Banda Rd
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Dead	1	Kanowna	~100-200m W of drilling track access road, on the Broad Arrow to Ora Banda road, Near Cawse Nickel turn off.
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Kanowna	~10km W of Menzies Hwy on Ora Banda - Broad Arrow road
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Day sighting	1		3.26km E of Cawse Nickel turn off on Broad Arrow - Ora banda Rd
Leipoa ocellata	malleefowl	BIRD	VU	VU	2014	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Burra Rock	Burra Rock, Directly west of campsite, south of old east-west track
Leipoa ocellata	malleefowl	BIRD	VU	VU	2015	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Parkeston	100m along a small track - off a major dirt road; approximately 9km NE of Kalgoorlie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2015	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Bulong	10.6km SSE of Hampton Hill Station homestead
Leipoa ocellata	malleefowl	BIRD	VU	VU	2015	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Feysville	Brown Hill on Woolubar Station. Kambolda Rd, Boulder WA
Leipoa ocellata	malleefowl	BIRD	VU	VU	2016	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	Coolgardie	Coolgardie North Rd, approx. 5km N of Coolgardie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2016	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	Coolgardie	Coolgardie North Rd, approx. 7km N of Coolgardie, near Bonnievale
Leipoa ocellata	malleefowl	BIRD	VU	VU	2015	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	Coolgardie	Between Burra Rocks Rd and Coolgardie-Esperance Rd, approx. 5km S of Coolgardie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2016	TFAUNA	Certain	Opportunistic sighting	Dusk sighting	1	Yallari Timber Reserve	Yallari Timber Reserve, central N-S track
Leipoa ocellata	malleefowl	BIRD	VU	VU	2016	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	Mount Burges	Traveling route back from Lake Carnage (Coolgardie Nth Rd??), 10km S of Ora Banda
Leipoa ocellata	malleefowl	BIRD	VU	VU	1996	TFAUNA	Moderately certain	Opportunistic sighting	Secondary sign	0	Coolgardie	4WD Holland Track, 200km NE of Mt Holland (cannot find Holland Track)
Leipoa ocellata	malleefowl	BIRD	VU	VU	1965	TFAUNA	Moderately certain	Historical (written)	Secondary sign	0	MOUNT BURGES	12 miles North of Coolgardie
Leipoa ocellata	malleefowl	BIRD	VU	VU	1994	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	2	BULLABULLING	22 kms west of Coolgardie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2004	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	KANOWNA	50 km nth of kalgoorlie on main hwy nth of Mt Vettters homestead
Leipoa ocellata	malleefowl	BIRD	VU	VU	1902	TFAUNA	Moderately certain	Opportunistic sighting	Secondary sign	0	BOORARA	Boorara
Leipoa ocellata	malleefowl	BIRD	VU	VU	2001	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	BULONG	Corsair Mine 10km E of Kalgoorlie; 1km S of Bulong Rd on graded track heading E
Leipoa ocellata	malleefowl	BIRD	VU	VU	1988	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	MOUNT BURGES	Eight Mile Rock dam
Leipoa ocellata	malleefowl	BIRD	VU	VU	2002	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	2	KARRAMINDIE	grt eastern hwy 1 km kal side of mungarrie industrial area
Leipoa ocellata	malleefowl	BIRD	VU	VU	2002	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	2	BULONG	Hampton Hill Station, 16km ESE of Boulder b/w Boorara Mine Site & Golden Ridge
Leipoa ocellata	malleefowl	BIRD	VU	VU	1910	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	LAMINGTON	Kalgoorlie
Leipoa ocellata	malleefowl	BIRD	VU	VU	1991	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	BULLABULLING	Kangaroo Hills Timber Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	1947	TFAUNA	Moderately certain	Opportunistic sighting	Secondary sign	0	LAMINGTON	PO Kalgoorlie
Leipoa ocellata	malleefowl	BIRD	VU	VU	1988	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	MOUNT BURGES	WMC sand pit Jaurdi Hills Rd
Leipoa ocellata	malleefowl	BIRD	VU	VU	1995	TFAUNA	Moderately certain	Opportunistic sighting	Sighting	1	LONDONDERRY	Yallari Timber Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	1995	TFAUNA	Moderately certain	Opportunistic sighting	Day sighting	1	LONDONDERRY	Yerilla Sandalwood Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2017	TFAUNA	Very Certain (photo, spec)	Opportunistic sighting	Day sighting	1	Goldfields Woodlands Conservation P	5km N of Victoria Rock, on Coolgardie Vic Rock Rd, in Goldfields Woodlands CP

CLASS	SCI_NAME	COM_NAME	WA_status	EPBCstatus	YEAR	SOURCE	CERTAINTY	OBS_METHOD	OBS_TYPE	COUNT	LOCALITY	SITE
Leipoa ocellata	malleefowl	BIRD	VU	VU	2017	TFAUNA	Very Certain (photo, spec)	Monitoring	Remote camera	1	Londonderry	Scahill Rimber Reserve, on track south of reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2017	TFAUNA	Very Certain (photo, spec)	Monitoring	Remote camera	1	Londonderry	Scahill Rimber Reserve, on track south of reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Very Certain (photo, spec)	Opportunistic sighting	Dawn sighting	2	Kanowna	Borad Arrow-Ora Banda Rd, 3km E of Cawse Nickel turnoff
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Not sure	Opportunistic sighting	Sighting	1	Ora Banda	Norton Gold field mining lease road
Leipoa ocellata	malleefowl	BIRD	VU	VU	2017	TFAUNA	Moderately certain	Opportunistic sighting	Day sighting	1	Karramindie	Karramindie State Forest No. 8
Leipoa ocellata	malleefowl	BIRD	VU	VU	2017	TFAUNA	Very Certain (photo, spec)	Opportunistic sighting	Dusk sighting	1	Karramindie	50m E of the S/W corner gate of Karramindie State Forest, where the gully crosses the track
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Very Certain (photo, spec)	Opportunistic sighting	Secondary sign	0	Karramindie	1.7km S of Karramindie State Forest on Hampton Location 53
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Londonberry	Old woodland track running SW away from Scahill Timber Reserve
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Very Certain (photo, spec)	Monitoring	Remote camera	2	Londonderry	3.5km S of Scahill Timber Reserve, Londonderry
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Certain	Opportunistic sighting	Day sighting	1	Kanowna	Golden cities Mine Site haul road, Kanowna
Calyptorhynchus lat	Carnaby's cockatoo	BIRD	EN	EN	2018	TFAUNA	Very Certain (photo, spec)	Opportunistic sighting	Day sighting	1	Sommerville	Southern corner of Hay St and Hutton St
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Very Certain (photo, spec)	Opportunistic sighting	Day sighting	0	Kalgoorlie	West north-west of Kalgoorlie about 20km
Leipoa ocellata	malleefowl	BIRD	VU	VU	2018	TFAUNA	Certain	Opportunistic sighting	Day sighting	2	Golden Cities	Goldfields mining tenure
Leipoa ocellata	malleefowl	BIRD	VU	VU	2007	TFAUNA	Certain	Opportunistic sighting		1	Coolgardie	Victoria Rock Rd, about 15km south of Coolgardie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2007	TFAUNA	Certain	Opportunistic sighting		1	Kalgoorlie	Broad Arrow-Ora Banda Rd, few km east of Cawse Mine turnoff
Leipoa ocellata	malleefowl	BIRD	VU	VU	2007	TFAUNA	Certain	Opportunistic sighting		1	Coolgardie	Burra Rock Road, 11.2km north of DEC Burra Rock Reserve boundary sign
Leipoa ocellata	malleefowl	BIRD	VU	VU	2012	TFAUNA	Certain	Opportunistic sighting		4	Coolgardie	Borefields on Focus owned mine lease, near bore 8
Leipoa ocellata	malleefowl	BIRD	VU	VU	2008	TFAUNA	Certain	Opportunistic sighting		1	Kambalda	
Leipoa ocellata	malleefowl	BIRD	VU	VU	2006	TFAUNA	Certain	Opportunistic sighting		1	Coolgardie	23.2km south of T intersection of Coolgardie Norseman Rd
Leipoa ocellata	malleefowl	BIRD	VU	VU	2008	TFAUNA	Certain	Opportunistic sighting	Dusk sighting	1	Coolgardie	Great Eastern Hwy, 40.5km west of Coolgardie
Leipoa ocellata	malleefowl	BIRD	VU	VU	2009	TFAUNA	Certain	Opportunistic sighting		2	Coolgardie	Juardi Hills Rd, 100m north of crest
Leipoa ocellata	malleefowl	BIRD	VU	VU	2009	TFAUNA	Certain	Opportunistic sighting		1	Londonerry	about 3km south of Burra Rock Reserve along Burra Rock Road
Leipoa ocellata	malleefowl	BIRD	VU	VU	2010	TFAUNA	Certain	Opportunistic sighting		1	Londonberry	Burra Rock Nature Reserve, next to camping area
Leipoa ocellata	malleefowl	BIRD	VU	VU	2010	TFAUNA	Certain	Opportunistic sighting		1	Kambalda	5km along pipeline access road off Cave Rocks mine haul rd
Leipoa ocellata	malleefowl	BIRD	VU	VU	2010	TFAUNA	Certain	Opportunistic sighting	Dead	1	Kambalda	9km N of Kambalda T Intersection along Goldfields Hwy
Leipoa ocellata	malleefowl	BIRD	VU	VU	2010	TFAUNA	Certain	Opportunistic sighting		1	Bullabulling	Great Eastern Hwy, 130km E of Southern Cross, near unnamed gravel road
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Opportunistic sighting		1	Ora Banda	Broad Arrow-Ora Banda Rd, approx. 5.5km from Norilsk Nickel access road toward east
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Opportunistic sighting		2	Londonderry	Burra Rock camping ground
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Survey		0	Ora Banda	
Leipoa ocellata	malleefowl	BIRD	VU	VU	2011	TFAUNA	Certain	Survey		0	Ora Banda	
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Sighting	1	Coolgardie	7km south of Coolgardie on the Victoria Rock Road
Leipoa ocellata	malleefowl	BIRD	VU	VU	0	TFAUNA	Certain	Opportunistic sighting	Secondary sign	0	Burra Rock Conservation Reserve	
Leipoa ocellata	malleefowl	BIRD	VU	VU	2012	TFAUNA	Certain	Opportunistic sighting	Sighting	1	Kalgoorlie	Goldfield HWY
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Sighting	1	Ora Banda	Between Broad Arrow and Ora Banda
Leipoa ocellata	malleefowl	BIRD	VU	VU	2013	TFAUNA	Certain	Opportunistic sighting	Sighting	1	Goldfields- Kalgoorlie/Coolgardie	Great Eastern Highway
Leipoa ocellata	malleefowl	BIRD	VU	VU	2019	TFAUNA	Certain	Opportunistic sighting	Dusk sighting	1	Arrow Lake	
Leipoa ocellata	malleefowl	BIRD	VU	VU	2019	TFAUNA	Certain	Opportunistic sighting	Sighting	1	Mungari	Mungari turnoff from Great Eastern HWY heading South
Leipoa ocellata	Malleefowl	BIRD	VU	VU	0	WAM_BIRDS	WAM Vouchered	Collection	Specimen	1	KALGOORLIE	Kalgoorlie
Leipoa ocellata	Malleefowl	BIRD	VU	VU	0	WAM_BIRDS	WAM Vouchered	Collection	Specimen	1		Kalgoorlie

NatureMap Species Report

Created By Guest user on 02/11/2021

Kingdom Animalia
Current Names Only Yes
Core Datasets Only Yes
Species Group All Animals
Method By Circle
Centre 121° 11' 12" E, 30° 55' 14" S
Buffer 40km
Group By Conservation Status

Conservation Status	Species	Records
Non-conservation taxon	297	4751
Priority 1	1	5
Priority 4	1	1
Protected under international agreement	6	13
Rare or likely to become extinct	6	67
TOTAL	311	4837

Name ID	Species Name	Naturalised	Conservation Code	Endemic To Query Area
Rare or likely to become extinct				
1.	24784 <i>Calidris ferruginea</i> (Curlew Sandpiper)		T	
2.	24734 <i>Calyptrornis latirostris</i> (Carnaby's Cockatoo, White-tailed Short-billed Black Cockatoo)		T	
3.	24557 <i>Leipoa ocellata</i> (Malleefowl)		T	
4.	24168 <i>Macrotis lagotis</i> (Bilby, Dalgite, Niniu)		T	
5.	24148 <i>Mymecobius fasciatus</i> (Numbat, Walpuri)		T	
6.	33987 <i>Ogyris subterrestris</i> subsp. <i>petrina</i> (And Bronze Azure Butterfly)		T	
Protected under international agreement				
7.	41323 <i>Actitis hypoleucos</i> (Common Sandpiper)		IA	
8.	24779 <i>Calidris acuminata</i> (Sharp-tailed Sandpiper)		IA	
9.	24780 <i>Calidris alba</i> (Sanderling)		IA	
10.	24788 <i>Calidris ruficollis</i> (Red-necked Stint)		IA	
11.	24806 <i>Tringa glareola</i> (Wood Sandpiper)		IA	
12.	24808 <i>Tringa nebularia</i> (Common Greenshank, greenshank)		IA	
Priority 1				
13.	33979 <i>Jalmenus aridus</i> (inland hairstreak, desert blue butterfly)		P1	Y
Priority 4				
14.	24803 <i>Tringa brevipes</i> (Grey-tailed Tattler)		P4	
Non-conservation taxon				
15.	24559 <i>Acanthogenys rubrogularis</i> (Spry-cheeked Honeyeater)			
16.	24260 <i>Acanthiza apicalis</i> (Broad-tailed Thornbill, Inland Thornbill)			
17.	24261 <i>Acanthiza chrysorrhoa</i> (Yellow-rumped Thornbill)			
18.	24265 <i>Acanthiza uropygialis</i> (Chestnut-rumped Thornbill)			
19.	25243 <i>Acanthophis pyrrhus</i> (Desert Death Adder)			
20.	25535 <i>Accipiter cirrocephalus</i> (Collared Sparrowhawk)			
21.	25536 <i>Accipiter fasciatus</i> (Brown Goshawk)			
22.	25544 <i>Aegotheles cristatus</i> (Australian Owllet-nightjar)			
23.	<i>Afrallacilla striolator</i>			
24.	<i>Afrostemophorus hirsti</i>			Y
25.	<i>Allodessus bistrigatus</i>			
26.	<i>Aname armigera</i>			
27.	<i>Anamio mainae</i>			
28.	24312 <i>Anas gracilis</i> (Grey Teal)			
29.	24313 <i>Anas platyrhynchos</i> (Mallard)			
30.	24315 <i>Anas rhynchotis</i> (Australasian Shoveler)			
31.	24316 <i>Anas superciliosa</i> (Pacific Black Duck)			
32.	47414 <i>Anhinga novaehollandiae</i> (Australasian Darter)			
33.	<i>Anidiops villosus</i>			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
34.	24561	<i>Anthochaera carunculata</i> (Red Wattlebird)			
35.	25670	<i>Anthus australis</i> (Australian Pipit)			
36.	24599	<i>Anthus australis</i> subsp. <i>australis</i> (Australian Pipit)			
37.	25528	<i>Aphelocephala leucopsis</i> (Southern Whiteface)			
38.	24266	<i>Aphelocephala leucopsis</i> subsp. <i>castaneiventris</i> (Southern Whiteface)			
39.	24285	<i>Aquila audax</i> (Wedge-tailed Eagle)			
40.	41324	<i>Ardea modesta</i> (great egret, white egret)			
41.	24341	<i>Ardea pacifica</i> (White-necked Heron)			
42.	24610	<i>Ardeotis australis</i> (Australian Bustard)			
43.		<i>Argiope protensa</i>			
44.		<i>Argiope trifasciata</i>			
45.	25566	<i>Artamus cinereus</i> (Black-faced Woodswallow)			
46.	24353	<i>Artamus cyanopterus</i> (Dusky Woodswallow)			
47.	24356	<i>Artamus personatus</i> (Masked Woodswallow)			
48.		<i>Austracantha minax</i>			
49.	24318	<i>Aythya australis</i> (Hardhead)			
50.		<i>Backobourkia heroine</i>			
51.		<i>Baiami tegenarioides</i>			
52.		<i>Barnardius zonarius</i>			
53.		<i>Berosus nutans</i>			
54.	24319	<i>Biziura lobata</i> (Musk Duck)			
55.	42380	<i>Brachyurophis fasciolatus</i> subsp. <i>fasciolatus</i> (Narrow-banded Shovel-nosed Snake)			
56.	42381	<i>Brachyurophis semifasciatus</i> (Southern Shovel-nosed Snake)			
57.	25715	<i>Cacatua roseicapilla</i> (Galah)			
58.	25716	<i>Cacatua sanguinea</i> (Little Corella)			
59.	42307	<i>Cacomantis pallidus</i> (Pallid Cuckoo)			
60.	24253	<i>Capra hircus</i> (Goat)	Y		
61.		<i>Carassius auratus</i>			
62.		<i>Celaenia excavata</i>			
63.	24086	<i>Cercartetus concinnus</i> (Western Pygmy-possum, Mundarda)			
64.		<i>Cercophonius michaelsoni</i>			
65.	24186	<i>Chalinolobus gouldii</i> (Gould's Wattle Bat)			
66.	24187	<i>Chalinolobus morio</i> (Chocolate Wattle Bat)			
67.	24377	<i>Charadrius ruficapillus</i> (Red-capped Plover)			
68.	43380	<i>Chelodina colliei</i> (South-western Snake-necked Turtle)			
69.	24321	<i>Chenonetta jubata</i> (Australian Wood Duck, Wood Duck)			
70.	47909	<i>Cheramoeca leucosterna</i> (White-backed Swallow)			
71.		<i>Chroicocephalus novaehollandiae</i>			
72.	24431	<i>Chrysococcyx basalis</i> (Horsfield's Bronze Cuckoo)			
73.	24434	<i>Chrysococcyx osculans</i> (Black-eared Cuckoo)			
74.	24289	<i>Circus assimilis</i> (Spotted Harrier)			
75.	24774	<i>Cladorhynchus leucocephalus</i> (Banded Stilt)			
76.	25675	<i>Colluricincla harmonica</i> (Grey Shrike-thrush)			
77.	24399	<i>Columba livia</i> (Domestic Pigeon)	Y		
78.	24361	<i>Coracina maxima</i> (Ground Cuckoo-shrike)			
79.	25568	<i>Coracina novaehollandiae</i> (Black-faced Cuckoo-shrike)			
80.		<i>Corasoides australis</i>			
81.		<i>Corimaethes campestris</i>			
82.		<i>Cormocephalus bungalbinensis</i>			
83.	24416	<i>Corvus bennetti</i> (Little Crow)			
84.	25592	<i>Corvus coronoides</i> (Australian Raven)			
85.	25593	<i>Corvus orru</i> (Torresian Crow)			
86.	24671	<i>Coturnix pectoralis</i> (Stubble Quail)			
87.	24420	<i>Cracticus nigrogularis</i> (Pied Butcherbird)			
88.	25595	<i>Cracticus tibicen</i> (Australian Magpie)			
89.	25596	<i>Cracticus torquatus</i> (Grey Butcherbird)			
90.	30893	<i>Cryptoblepharus buechananii</i>			
91.	25020	<i>Cryptoblepharus plagiocephalus</i>			
92.	25458	<i>Ctenophorus caudicinctus</i> (Ring-tailed Dragon)			
93.	24871	<i>Ctenophorus cristatus</i> (Bicycle Dragon)			
94.	24873	<i>Ctenophorus fordii</i> (Mallee Sand Dragon)			
95.	24874	<i>Ctenophorus isolepis</i> subsp. <i>citrinus</i> (Yellowy Military Dragon)			
96.	24882	<i>Ctenophorus nuchalis</i> (Central Netted Dragon)			
97.	24886	<i>Ctenophorus reticulatus</i> (Western Netted Dragon)			
98.	24888	<i>Ctenophorus salinarum</i> (Salt Pan Dragon)			
99.	24889	<i>Ctenophorus scutulatus</i> (Lozenge-marked Dragon)			
100.	25026	<i>Ctenotus atlas</i>			
101.	25052	<i>Ctenotus leonhardii</i>			
102.	25074	<i>Ctenotus schomburgkii</i>			
103.	25465	<i>Ctenotus uber</i> (Spotted Ctenotus)			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
104.	25080	<i>Ctenotus uber</i> subsp. <i>uber</i> (Spotted Ctenotus)			
105.	25089	<i>Cyclodomorphus melanops</i> subsp. <i>elongatus</i> (Slender Blue-tongue)			
106.	24322	<i>Cygnus atratus</i> (Black Swan)			
107.		<i>Cyrtophora parnasia</i>			
108.		<i>Daphnia carinata</i>			
109.	25673	<i>Daphoenositta chrysoptera</i> (Varied Sittella)			
110.	24606	<i>Daphoenositta chrysoptera</i> subsp. <i>pileata</i> (Varied Sittella, Black-capped Sittella)			
111.	24995	<i>Delma australis</i>			
112.	25247	<i>Demansia psammophis</i> subsp. <i>psammophis</i> (Yellow-faced Whipsnake)			
113.	25607	<i>Dicaeum hirundinaceum</i> (Mistletoebird)			
114.	25469	<i>Diplodactylus granariensis</i>			
115.	24929	<i>Diplodactylus granariensis</i> subsp. <i>granariensis</i>			
116.	24940	<i>Diplodactylus pulcher</i>			
117.	24470	<i>Dromaius novaehollandiae</i> (Emu)			
118.	24650	<i>Drymodes brunneopygia</i> (Southern Scrub-robin)			
119.	25092	<i>Egernia depressa</i> (Southern Pygmy Spiny-tailed Skink)			
120.	25094	<i>Egernia formosa</i>			
121.	25104	<i>Egernia richardi</i>			
122.		<i>Egretta novaehollandiae</i>			
123.		<i>Elanus axillaris</i>			
124.	24290	<i>Elanus caeruleus</i> subsp. <i>axillaris</i> (Australian Black-shouldered Kite)			
125.	47937	<i>Elseyanis melanops</i> (Black-fronted Dotterel)			
126.		<i>Enochrus elongatulus</i>			
127.		<i>Eolophus roseicapillus</i>			
128.	24651	<i>Eopsaltria australis</i> subsp. <i>griseogularis</i> (Western Yellow Robin)			
129.	24567	<i>Epthianura albiglans</i> (White-fronted Chat)			
130.	24570	<i>Epthianura tricolor</i> (Crimson Chat)			
131.	25109	<i>Eremiascincus richardsonii</i> (Broad-banded Sand Swimmer)			
132.		<i>Eriophora biapicata</i>			
133.	24379	<i>Erythronyx cinctus</i> (Red-kneed Dotterel)			
134.	24368	<i>Eurostopodus argus</i> (Spotted Nightjar)			
135.	25621	<i>Falco berigora</i> (Brown Falcon)			
136.	24471	<i>Falco berigora</i> subsp. <i>berigora</i> (Brown Falcon)			
137.	25622	<i>Falco cenchroides</i> (Australian Kestrel, Nankeen Kestrel)			
138.	25623	<i>Falco longipennis</i> (Australian Hobby)			
139.	24041	<i>Felis catus</i> (Cat)	Y		
140.		<i>Fissarena castanea</i>			
141.	25727	<i>Fulica atra</i> (Eurasian Coot)			
142.	25301	<i>Furina ornata</i> (Moon Snake)			
143.	24957	<i>Gehyra purpurascens</i>			
144.	24959	<i>Gehyra variegata</i>			
145.	25530	<i>Gerygone fusca</i> (Western Gerygone)			
146.	24443	<i>Grallina cyanoleuca</i> (Magpie-lark)			
147.	24295	<i>Haliastur sphenurus</i> (Whistling Kite)			
148.		<i>Hemicloea sublimbata</i>			
149.	25232	<i>Hemidactylus frenatus</i> (Asian House Gecko)	Y		
150.	25115	<i>Hemiergis initialis</i> subsp. <i>initialis</i>			
151.	42408	<i>Hesperoedura reticulata</i>			
152.	24961	<i>Heteronotia binoei</i> (Bynoe's Gecko)			
153.	47965	<i>Hieraaetus morphnoides</i> (Little Eagle)			
154.	25734	<i>Himantopus himantopus</i> (Black-winged Stilt)			
155.	24775	<i>Himantopus himantopus</i> subsp. <i>leucocephalus</i> (Black-winged Stilt)			
156.	24491	<i>Hirundo neoxena</i> (Welcome Swallow)			
157.		<i>Hoggicosa castanea</i>			
158.		<i>Hoggicosa forresti</i>			
159.		<i>Holoplatys kalgoorlie</i>			Y
160.		<i>Holoplatys planissima</i>			
161.	34001	<i>Hylacola cauta</i> subsp. <i>whitlocki</i> (Shy Groundwren)			
162.		<i>Idiomata blackwalli</i>			
163.		<i>Isometroides vesus</i>			
164.		<i>Isopeda magna</i>			
165.		<i>Isopedella saundersi</i>			
166.		<i>Jalmenus icilius</i>			Y
167.	24367	<i>Lalage tricolor</i> (White-winged Triller)			
168.		<i>Lampona cylindrata</i>			
169.		<i>Lamponina scutata</i>			
170.		<i>Latrodectus hasseltii</i>			
171.		<i>Lerista kingi</i>			
172.	25155	<i>Lerista muelleri</i>			
173.	25162	<i>Lerista picturata</i>			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
174.	25172	<i>Lerista stictopleura</i>			
175.	42411	<i>Lerista timida</i>			
176.	25005	<i>Lialis burtonis</i>			
177.	25659	<i>Lichenostomus leucotis</i> (White-eared Honeyeater)			
178.	24576	<i>Lichenostomus leucotis</i> subsp. <i>novaenorcae</i> (White-eared Honeyeater)			
179.	25661	<i>Lichmera indistincta</i> (Brown Honeyeater)			
180.	41411	<i>Liopholis inornata</i> (Desert Skink)			
181.	25388	<i>Litoria moorei</i> (Motorbike Frog)			
182.	30935	<i>Lucasium maini</i>			
183.		<i>Lycosa ariadnae</i>			
184.	24132	<i>Macropus fuliginosus</i> (Western Grey Kangaroo)			
185.	24136	<i>Macropus rufus</i> (Red Kangaroo, Marlu)			
186.	24326	<i>Malacorhynchus membranaceus</i> (Pink-eared Duck)			
187.	25652	<i>Malurus leucopterus</i> (White-winged Fairy-wren)			
188.	24551	<i>Malurus pulcherrimus</i> (Blue-breasted Fairy-wren)			
189.	25654	<i>Malurus splendens</i> (Splendid Fairy-wren)			
190.	24583	<i>Manorina flavigula</i> (Yellow-throated Miner)			
191.	25663	<i>Melithreptus brevirostris</i> (Brown-headed Honeyeater)			
192.	24736	<i>Melopsittacus undulatus</i> (Budgerigar)			
193.	25184	<i>Menetia greyii</i>			
194.	24598	<i>Merops ornatus</i> (Rainbow Bee-eater)			
195.		<i>Microcarbo melanoleucos</i>			
196.	25693	<i>Microeca fascians</i> (Jacky Winter)			
197.	24654	<i>Microeca fascians</i> subsp. <i>assimilis</i> (Jacky Winter)			
198.		<i>Missulena occatoria</i>			
199.	24904	<i>Moloch horridus</i> (Thorny Devil)			
200.	25240	<i>Morelia spilota</i> subsp. <i>imbricata</i> (Carpet Python)			
201.	25188	<i>Morethia adelaidensis</i>			
202.	25190	<i>Morethia butleri</i>			
203.	24223	<i>Mus musculus</i> (House Mouse)	Y		
204.	25248	<i>Neelaps bimaculatus</i> (Black-naped Snake)			
205.	25425	<i>Neobatrachus kunapalari</i> (Kunapalari Frog)			
206.	25426	<i>Neobatrachus pelobatoides</i> (Humming Frog)			
207.	25427	<i>Neobatrachus sutor</i> (Shoemaker Frog)			
208.	25428	<i>Neobatrachus wilsmorei</i> (Plonking Frog)			
209.		<i>Nephila edulis</i>			
210.	24971	<i>Nephurus vertebralis</i>			
211.		<i>Nicodamus mainae</i>			
212.	24096	<i>Ningui yvonneae</i> (Southern Ningui)			
213.	24229	<i>Notomys mitchellii</i> (Mitchell's Hopping-mouse)			
214.	24194	<i>Nyctophilus geoffroyi</i> (Lesser Long-eared Bat)			
215.	24742	<i>Nymphicus hollandicus</i> (Cockatiel)			
216.	24407	<i>Ocyphaps lophotes</i> (Crested Pigeon)			
217.	24618	<i>Oreica gutturalis</i> (Crested Bellbird)			
218.	34011	<i>Oreica gutturalis</i> subsp. <i>gutturalis</i> (Crested Bellbird (southern))			
219.	24085	<i>Oryctolagus cuniculus</i> (Rabbit)	Y		
220.		<i>Ostracoda</i> (unident.)			
221.		<i>Oxyopes amoenus</i>			
222.		<i>Oxyopes dingo</i>			
223.		<i>Oxyopes variabilis</i>			
224.		<i>Ozestheria packardii</i>			
225.	24619	<i>Pachycephala inornata</i> (Gilbert's Whistler)			
226.	25680	<i>Pachycephala rufiventris</i> (Rufous Whistler)			
227.		<i>Parartemia</i> sp.			
228.	25253	<i>Parasuta gouldii</i>			
229.	25254	<i>Parasuta monachus</i>			
230.	25681	<i>Pardalotus punctatus</i> (Spotted Pardalote)			
231.	25682	<i>Pardalotus striatus</i> (Striated Pardalote)			
232.	24630	<i>Pardalotus striatus</i> subsp. <i>westraliensis</i> (Striated Pardalote)			
233.	48060	<i>Petrochelidon ariel</i> (Fairy Martin)			
234.	48061	<i>Petrochelidon nigricans</i> (Tree Martin)			
235.	24659	<i>Petroica goodenovii</i> (Red-capped Robin)			
236.	24667	<i>Phalacrocorax sulcirostris</i> (Little Black Cormorant)			
237.	24409	<i>Phaps chalcoptera</i> (Common Bronzewing)			
238.	24841	<i>Platalea flavipes</i> (Yellow-billed Spoonbill)			
239.	25720	<i>Platycercus icterotis</i> (Western Rosella)			
240.	24748	<i>Platycercus varius</i> (Mulga Parrot)			
241.	25721	<i>Platycercus zonarius</i> (Australian Ringneck, Ring-necked Parrot)			
242.	24751	<i>Platycercus zonarius</i> subsp. <i>zonarius</i> (Port Lincoln Parrot)			
243.	25703	<i>Podargus strigoides</i> (Tawny Frogmouth)			

	Name ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
244.	24907	<i>Pogona minor</i> subsp. <i>minor</i> (Dwarf Bearded Dragon)			
245.	24681	<i>Poliocephalus poliocephalus</i> (Hoary-headed Grebe)			
246.	30854	<i>Polytelis anthopeplus</i> subsp. <i>westralis</i> (Regent Parrot)			
247.	24683	<i>Pomatostomus superciliosus</i> (White-browed Babbler)			
248.	34013	<i>Pomatostomus superciliosus</i> subsp. <i>ashbyi</i> (White-browed Babbler (western wheatbelt))			
249.	24769	<i>Porzana fluminea</i> (Australian Spotted Crane)			
250.	25261	<i>Pseudechis australis</i> (Mulga Snake)			
251.	24232	<i>Pseudomys bolami</i> (Bolam's Mouse)			
252.	24237	<i>Pseudomys hermannsburgensis</i> (Sandy Inland Mouse)			
253.	42416	<i>Pseudonaja mengdeni</i> (Western Brown Snake)			
254.	25263	<i>Pseudonaja modesta</i> (Ringed Brown Snake)			
255.	25434	<i>Pseudophryne occidentalis</i> (Western Toadlet)			
256.	42344	<i>Purnella albifrons</i> (White-fronted Honeyeater)			
257.	25008	<i>Pygopus lepidopodus</i> (Common Scaly Foot)			
258.	25009	<i>Pygopus nigriceps</i>			
259.	24278	<i>Pyrrholaemus brunneus</i> (Redthroat)			
260.	24776	<i>Recurvirostra novaehollandiae</i> (Red-necked Avocet)			
261.	48096	<i>Rhipidura albiscapa</i> (Grey Fantail)			
262.	25614	<i>Rhipidura leucophrys</i> (Willie Wagtail)			
263.	24982	<i>Rhynchoedura ornata</i> (Western Beaked Gecko)			
264.		<i>Sandalodes scopifer</i>			
265.		<i>Scolopendra laeta</i>			
266.		<i>Scolopendra morsitans</i>			
267.	24199	<i>Scotorepens balstoni</i> (Inland Broad-nosed Bat)			
268.		<i>Selenotholus foelschei</i>			
269.	25266	<i>Simoselaps bertholdi</i> (Jan's Banded Snake)			
270.	30948	<i>Smicrornis brevirostris</i> (Weebill)			
271.	24108	<i>Sminthopsis crassicaudata</i> (Fat-tailed Dunnart)			
272.	24109	<i>Sminthopsis dolichura</i> (Little long-tailed Dunnart)			
273.	24111	<i>Sminthopsis gilberti</i> (Gilbert's Dunnart)			
274.	24329	<i>Stictonetta naevosa</i> (Freckled Duck)			
275.		<i>Storena sinuosa</i>			
276.	25597	<i>Strepera versicolor</i> (Grey Currawong)			
277.	25590	<i>Streptopelia senegalensis</i> (Laughing Turtle-Dove)	Y		
278.	24923	<i>Strophurus assimilis</i> (Goldfields Spiny-tailed Gecko)			
279.	24927	<i>Strophurus elderi</i>			
280.	25269	<i>Suta fasciata</i> (Rosen's Snake)			
281.	25705	<i>Tachybaptus novaehollandiae</i> (Australasian Grebe, Black-throated Grebe)			
282.	24207	<i>Tachyglossus aculeatus</i> (Short-beaked Echidna)			
283.	24331	<i>Tadorna tadornoides</i> (Australian Shelduck, Mountain Duck)			
284.	30870	<i>Taeniopygia guttata</i> (Zebra Finch)			
285.		<i>Tamopsis circumvidens</i>			
286.	24176	<i>Taphozous hilli</i> (Hill's Sheath-tail-bat)			
287.		<i>Tasmanicosa leuckartii</i>			
288.	24845	<i>Threskiornis spinicollis</i> (Straw-necked Ibis)			
289.	25203	<i>Tiliqua occipitalis</i> (Western Bluetongue)			
290.	25519	<i>Tiliqua rugosa</i>			
291.	25207	<i>Tiliqua rugosa</i> subsp. <i>rugosa</i>			
292.	42351	<i>Todiramphus pyrrhopygius</i> (Red-backed Kingfisher)			
293.	25549	<i>Todiramphus sanctus</i> (Sacred Kingfisher)			
294.	48141	<i>Tribonyx ventralis</i> (Black-tailed Native-hen)			
295.		<i>Trichocyclops balladong</i>			
296.	39407	<i>Triops australiensis</i> (Shield Shrimp)			
297.	24851	<i>Turnix velox</i> (Little Button-quail)			
298.	30814	<i>Tympanocryptis cephalus</i> (Pebble Dragon)			
299.	39408	<i>Tympanocryptis lineata</i> (Lined Earless Dragon)			
300.	24852	<i>Tyto alba</i> subsp. <i>delicatula</i> (Barn Owl)			
301.	24983	<i>Underwoodisaurus milii</i> (Barking Gecko)			
302.		<i>Urodacus armatus</i>			
303.		<i>Urodacus hoplurus</i>			
304.		<i>Urodacus yaschenkoi</i>			
305.	24386	<i>Vanellus tricolor</i> (Banded Lapwing)			
306.	25211	<i>Varanus caudolineatus</i>			
307.	25218	<i>Varanus gouldii</i> (Bungarra or Sand Monitor)			
308.	25526	<i>Varanus tristis</i> (Racehorse Monitor)			
309.	24202	<i>Vespadelus baverstocki</i> (Inland Forest Bat)			
310.	24206	<i>Vespadelus regulus</i> (Southern Forest Bat)			
311.	25765	<i>Zosterops lateralis</i> (Grey-breasted White-eye, Silvereye)			

Name	ID	Species Name	Naturalised	Conservation Code	¹ Endemic To Query Area
Conservation Codes					
T - Rare or likely to become extinct					
X - Presumed extinct					
IA - Protected under international agreement					
S - Other specially protected fauna					
1 - Priority 1					
2 - Priority 2					
3 - Priority 3					
4 - Priority 4					
5 - Priority 5					

¹ For NatureMap's purposes, species flagged as endemic are those whose records are wholly contained within the search area. Note that only those records complying with the search criterion are included in the calculation. For example, if you limit records to those from a specific datasource, only records from that datasource are used to determine if a species is restricted to the query area.



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.


Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about [Environment Assessments](#) and the EPBC Act including significance guidelines, forms and application process details.

Report created: 25/10/21 19:14:20

- [Summary](#)
- [Details](#)

[Matters of NES](#)[Other Matters Protected by the EPBC Act](#)[Extra Information](#)
- [Caveat](#)
- [Acknowledgements](#)



This map may contain data which are
©Commonwealth of Australia
(Geoscience Australia), ©PSMA 2015

[Coordinates](#)

Buffer: 50.0Km



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the [Administrative Guidelines on Significance](#).

World Heritage Properties:	None
National Heritage Places:	1
Wetlands of International Importance:	None
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	None
Listed Threatened Ecological Communities:	None
Listed Threatened Species:	8
Listed Migratory Species:	7

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at <http://www.environment.gov.au/heritage>

A [permit](#) may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	4
Commonwealth Heritage Places:	None
Listed Marine Species:	11
Whales and Other Cetaceans:	None
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Australian Marine Parks:	None

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	9
Regional Forest Agreements:	None
Invasive Species:	16
Nationally Important Wetlands:	None
Key Ecological Features (Marine)	None

Details

Matters of National Environmental Significance

National Heritage Properties		[Resource Information]
Name	State	Status
Historic		
Goldfields Water Supply Scheme, Western Australia	WA	Listed place

Listed Threatened Species		[Resource Information]
Name	Status	Type of Presence
Birds		
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat likely to occur within area
Falco hypoleucos Grey Falcon [929]	Vulnerable	Species or species habitat may occur within area
Leipoa ocellata Malleefowl [934]	Vulnerable	Species or species habitat known to occur within area
Pezoporus occidentalis Night Parrot [59350]	Endangered	Species or species habitat may occur within area
Insects		
Ogyris subterrestris petrina Arid Bronze Azure [77743]	Critically Endangered	Species or species habitat may occur within area
Mammals		
Dasyurus geoffroii Chuditch, Western Quoll [330]	Vulnerable	Species or species habitat may occur within area
Plants		
Gastrolobium graniticum Granite Poison [14872]	Endangered	Species or species habitat known to occur within area
Thelymitra stellata Star Sun-orchid [7060]	Endangered	Species or species habitat may occur within area

Listed Migratory Species		[Resource Information]
* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.		
Name	Threatened	Type of Presence
Migratory Marine Birds		
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Migratory Terrestrial Species		

Name	Threatened	Type of Presence
Motacilla cinerea Grey Wagtail [642]		Species or species habitat may occur within area
Migratory Wetlands Species		
Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat may occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat known to occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat likely to occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat likely to occur within area

Other Matters Protected by the EPBC Act

Commonwealth Land	[Resource Information]
-------------------	--

The Commonwealth area listed below may indicate the presence of Commonwealth land in this vicinity. Due to the unreliability of the data source, all proposals should be checked as to whether it impacts on a Commonwealth area, before making a definitive decision. Contact the State or Territory government land department for further information.

Name
Commonwealth Land - Defence - AIRTC KALGOORLIE Defence - KALGOORLIE RIFLE RANGE Defence - KALGOORLIE TRAINING DEPOT

Listed Marine Species	[Resource Information]
-----------------------	--

* Species is listed under a different scientific name on the EPBC Act - Threatened Species list.

Name	Threatened	Type of Presence
Birds		
Actitis hypoleucos Common Sandpiper [59309]		Species or species habitat may occur within area
Apus pacificus Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea ibis Cattle Egret [59542]		Species or species habitat may occur within area
Calidris acuminata Sharp-tailed Sandpiper [874]		Species or species habitat known to occur within area
Calidris ferruginea Curlew Sandpiper [856]	Critically Endangered	Species or species habitat likely to occur within area
Calidris melanotos Pectoral Sandpiper [858]		Species or species habitat may occur within area
Chrysococcyx osculans Black-eared Cuckoo [705]		Species or species

Name	Threatened	Type of Presence
		habitat known to occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Motacilla cinerea Grey Wagtail [642]		Species or species habitat may occur within area
Thinornis rubricollis Hooded Plover [59510]		Species or species habitat may occur within area
Tringa nebularia Common Greenshank, Greenshank [832]		Species or species habitat likely to occur within area

Extra Information

State and Territory Reserves	[Resource Information]
Name	State
Goldfields Woodlands	WA
Kalgoorlie Arboretum	WA
Kambalda	WA
Kangaroo Hills Timber Reserve	WA
Kurrawang	WA
Lakeside Timber Reserve	WA
Scahill Timber Reserve	WA
Victoria Rock	WA
Yallari Timber Reserve	WA

Invasive Species	[Resource Information]
Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resouces Audit, 2001.	

Name	Status	Type of Presence
Birds		
Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
Streptopelia chinensis Spotted Turtle-Dove [780]		Species or species habitat likely to occur within area
Streptopelia senegalensis Laughing Turtle-dove, Laughing Dove [781]		Species or species habitat likely to occur within area
Mammals		
Canis lupus familiaris Domestic Dog [82654]		Species or species habitat likely to occur within area
Capra hircus Goat [2]		Species or species

Name	Status	Type of Presence
		habitat likely to occur within area
Equus asinus Donkey, Ass [4]		Species or species habitat likely to occur within area
Equus caballus Horse [5]		Species or species habitat likely to occur within area
Felis catus Cat, House Cat, Domestic Cat [19]		Species or species habitat likely to occur within area
Mus musculus House Mouse [120]		Species or species habitat likely to occur within area
Oryctolagus cuniculus Rabbit, European Rabbit [128]		Species or species habitat likely to occur within area
Vulpes vulpes Red Fox, Fox [18]		Species or species habitat likely to occur within area
Plants		
Carrichtera annua Ward's Weed [9511]		Species or species habitat likely to occur within area
Cenchrus ciliaris Buffel-grass, Black Buffel-grass [20213]		Species or species habitat may occur within area
Cylindropuntia spp. Prickly Pears [85131]		Species or species habitat likely to occur within area
Lycium ferocissimum African Boxthorn, Boxthorn [19235]		Species or species habitat likely to occur within area
Reptiles		
Hemidactylus frenatus Asian House Gecko [1708]		Species or species habitat likely to occur within area

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-30.92058 121.18644

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- [-Office of Environment and Heritage, New South Wales](#)
- [-Department of Environment and Primary Industries, Victoria](#)
- [-Department of Primary Industries, Parks, Water and Environment, Tasmania](#)
- [-Department of Environment, Water and Natural Resources, South Australia](#)
- [-Department of Land and Resource Management, Northern Territory](#)
- [-Department of Environmental and Heritage Protection, Queensland](#)
- [-Department of Parks and Wildlife, Western Australia](#)
- [-Environment and Planning Directorate, ACT](#)
- [-Birdlife Australia](#)
- [-Australian Bird and Bat Banding Scheme](#)
- [-Australian National Wildlife Collection](#)
- Natural history museums of Australia
- [-Museum Victoria](#)
- [-Australian Museum](#)
- [-South Australian Museum](#)
- [-Queensland Museum](#)
- [-Online Zoological Collections of Australian Museums](#)
- [-Queensland Herbarium](#)
- [-National Herbarium of NSW](#)
- [-Royal Botanic Gardens and National Herbarium of Victoria](#)
- [-Tasmanian Herbarium](#)
- [-State Herbarium of South Australia](#)
- [-Northern Territory Herbarium](#)
- [-Western Australian Herbarium](#)
- [-Australian National Herbarium, Canberra](#)
- [-University of New England](#)
- [-Ocean Biogeographic Information System](#)
- [-Australian Government, Department of Defence](#)
- [Forestry Corporation, NSW](#)
- [-Geoscience Australia](#)
- [-CSIRO](#)
- [-Australian Tropical Herbarium, Cairns](#)
- [-eBird Australia](#)
- [-Australian Government – Australian Antarctic Data Centre](#)
- [-Museum and Art Gallery of the Northern Territory](#)
- [-Australian Government National Environmental Science Program](#)
- [-Australian Institute of Marine Science](#)
- [-Reef Life Survey Australia](#)
- [-American Museum of Natural History](#)
- [-Queen Victoria Museum and Art Gallery, Inveresk, Tasmania](#)
- [-Tasmanian Museum and Art Gallery, Hobart, Tasmania](#)
- Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the [Contact Us](#) page.

[© Commonwealth of Australia](#)

[Department of Agriculture Water and the Environment](#)

GPO Box 858

Canberra City ACT 2601 Australia

+61 2 6274 1111

Appendix C

Coolgardie Flora Likelihood Table

Appendix: Assessment of the Likelihood of Occurrence of Threatened and Priority Flora as per Desktop Assessment Database Searches surrounding the Survey Area

Distance to Nearest Record from the Survey Area is based on a distance analysis undertaken against 2021 DBCA database. High = Suitable habitat present and records less than 15 km from the Survey Area, Medium = Suitable habitat present and records between 15 km and 30 km from the Survey Area, and Low = No suitable habitat present and/or records greater than 30 km from the Survey Area, Unknown = Insufficient information available to classify. CR= Listed as Critically Endangered under the EPBC Act, EN = Listed as Endangered under the EPBC Act, VU = listed as Vulnerable under the EPBC Act. T = Threatened under the BC Act, P = Priority Listed, Ranked and Listed by the DBCA. Likelihoods are assessed both pre and post survey based on knowledge of the Survey Area, nearest known records, known flowering period of flora taxa and knowledge gained from the survey effort during ground truthing. 1: Department of the Environment (2021). SPRAT EPBC Threatened Flora in Species Profile and Threats Database, Department of the Environment, Canberra. Available from: <http://www.environment.gov.au/sprat>. 2: Department of Biodiversity, Conservation and Attractions (2021). FloraBase - The Western Australian Flora. <https://florabase.dpaw.wa.gov.au/>

Species	Conservation Status		Source			Distance to Nearest Record (km)	Flowering Period	Preferred Habitat	Habitat occurs within the Survey Area	Pre-Survey Likelihood of Occurrence	Post-Survey Likelihood of Occurrence
	DBCA	EPBC	NatureMap	PMST	DBCA						
<i>Gastrolobium graniticum</i>	T	EN	X	X	X	25.8	Aug to Sep.	Sand, sandy loam, granite. Margins of rock outcrops, along drainage lines. ²	No	Medium	Low
<i>Tetratheca spenceri</i>	T				X	39.1	Mar	Gentle slope on duricrust breakaway. ²	No	Low	Low
<i>Thelymitra stellata</i>	T	EN		X		12.8	Oct - Nov	Grows on ridges and slopes, flats, also on riverbanks and breakaways. Soil types are red, brown, yellow, or grey sandy loams clay or gravel over laterite or gravel. Dry, moist or saline conditions are tolerated. ¹	Yes	High	Medium
<i>Acacia coatesii</i>	P1		X			7.4	Sep	Flat to gentle slope, red sandy loam soils. ²	Yes	High	Medium
<i>Acacia epedunculata</i>	P1				X	18.6	Aug - Oct	Yellow sand. Sandplains. ²	Yes	Medium	Low
<i>Acacia sclerophylla</i> var. <i>teretiuscula</i>	P1		X			16.1	Sep - Oct	Clay and loamy soils. ²	Yes	Medium	Low
<i>Acacia websteri</i>	P1		X		X	0.7	Jun	Red sand, clay or loam. Low-lying areas, flats. ²	Yes	High	High
<i>Calandrinia lefroyensis</i>	P1				X	10.8	Feb or Apr or Oct - Nov	Light brown, orange or red sand, sandy loam, sandy clay loam over sediments. Fine quartz. Gentle slopes, near salt lakes or salt flats, sand dunes. ²	Yes	High	Medium
<i>Chamelaucium</i> sp. <i>Parker Range</i> (B.H. Smith 1255)	P1				X	19	Nov or Dec	Sandplains, Mid slope. Dry, yellow sand over laterite. ²	Yes	Medium	Low
<i>Eremophila xantholaema</i>	P1				X	17.7	Sep - Oct	Hilltop and slopes. Brown/orange/red very rocky loam/granite. ²	Yes	Medium	Low
<i>Cyathostemon divaricatus</i>	P1				X	40.5	Jan or Jun or Aug	Hills, hillslope. ²	Yes	Low	Low
<i>Dampiera plumosa</i>	P1		X			4.5	Oct	Red sandy soils. ²	Yes	High	Medium
<i>Eucalyptus websteriana</i> subsp. <i>norsemanica</i>	P1		X			8.7	Sep to Nov	Rocky rises. ²	Yes	High	Medium
<i>Lepidosperma lyonsii</i>	P1				X	70.6	Jun	Orange skeletal sandy loam with banded ironstone gravel and rock, well-drained shallow stony loam with quartz. Gentle hill slopes, upper slopes of large hill. ²	Yes	Low	Low
<i>Lepidosperma</i> sp. <i>Parker Range</i> (N. Gibson & M. Lyons 2094)	P1		X			8.3	NA	Brown sandy loam, or clay. Gravel, laterite. Banded ironstone formations, ridges, mid-slopes. ²	Yes	High	Medium

¹ Department of Agriculture, Water and Environment (2020) ²Western Australian Herbarium (2020)

Species	Conservation Status		Source			Distance to Nearest Record (km)	Flowering Period	Preferred Habitat	Habitat occurs within the Survey Area	Pre-Survey Likelihood of Occurrence	Post-Survey Likelihood of Occurrence
	DBCA	EPBC	NatureMap	PMST	DBCA						
<i>Melichrus</i> sp. Coolgardie (K.R. Newbey 8698)	P1				X	24.8	April or Jun or Aug - Dec	Yellow loamy sand. Sandplains. ²	Yes	Medium	Low
<i>Phebalium appressum</i>	P1		X			7.8	Jul	Yellow sandplain. ²	Yes	High	Low
<i>Philotheca apiculata</i>	P1				X	45.8	Aug - Nov	Stony clay loam. Rocky outcrops, hillsides. ²	Yes	Low	Low
<i>Philotheca pachyphylla</i>	P1				X	31.9	May or Sep	Sand, red loam, clay loam. Sandplains, hill tops. ²	Yes	Low	Low
<i>Prostanthera splendens</i>	P1				X	66.8	Aug - Oct	Stony loam, shallow soils with ironstone pebbles. Breakaways. ²	No	Low	Low
<i>Pterostylis xerampelina</i>	P1				X	66.1	Sept - Oct	Granite. ²	Yes	Low	Low
<i>Ptilotus procumbens</i>	P1				X	30.2	Nov	Red clay. ²	Yes	Low	Low
<i>Ptilotus rigidus</i>	P1				X	30.4	Mar or May or Oct - Jan	Clay, clay loam on quartz hills and edges of salt lakes. ²	Yes	Low	Low
<i>Ptilotus</i> sp. Kalgoorlie (J. Jackson & B. Moyle 260)	P1				X	36.1	NA	Quartz hills. ²	Yes	Low	Low
<i>Rhodanthe uniflora</i>	P1				X	44.8	Aug - Oct	Brown earth. Open <i>Eucalyptus</i> woodland. ²	Yes	Low	Low
<i>Ricinocarpus digynus</i>	P1				X	50.4	Aug	Red-brown sand-loam. Rocky hillslopes and plains. ²	Yes	Low	Low
<i>Tecticornia flabelliformis</i>	P1	VU		X	X	54.8	Dec	Clay. Saline flats. ²	Yes	Low	Low
<i>Tecticornia mellarium</i>	P1				X	58.4	NA	Close proximity to salt lakes and dunes. ²	Yes	Low	Low
<i>Thryptomene planiflora</i>	P1		X			19.1	Jul or Aug - Sep	Brown, yellow or orange sandy loam. Lateritic gravel. Gentle hillslopes, sandplains, slight depressions. ²	Yes	Medium	Medium
<i>Thryptomene</i> sp. Coolgardie (E. Kelso s.n. 1902)	P1		X			2.2	NA	NA	Unknown	Unknown	Medium
<i>Acacia kerryana</i>	P2				X	39.2	Oct - Dec or Jan - Feb	Granitic loamy sand, stony clayey loam or clayey sand. Low stony ridges, undulating plains. ²	Yes	Low	Low
<i>Austrostipa frankliniae</i>	P2		X			8.5	Oct - Nov	Flat plain, Basalt slope. Dry brown / red loam. ²	Yes	High	Medium
<i>Bossiaea laxa</i>	P2				X	43.3	May	Brown loam over deep granite. Sheltered positions around outcrops. ²	No	Low	Low

¹ Department of Agriculture, Water and Environment (2020) ²Western Australian Herbarium (2020)

Species	Conservation Status		Source			Distance to Nearest Record (km)	Flowering Period	Preferred Habitat	Habitat occurs within the Survey Area	Pre-Survey Likelihood of Occurrence	Post-Survey Likelihood of Occurrence
	DBCA	EPBC	NatureMap	PMST	DBCA						
<i>Elachanthus pusillus</i>	P2				X	32.1	Aug - Oct	Dry brown or red-orange loam clay. Quartz, limestone, granite. Low plains, drainage flats, gentle upper slopes. ²	Yes	Low	Low
<i>Eremophila praecox</i>	P2		X		X	17.2	Oct or Dec.	Red/brown sandy loam. Undulating plains. ²	Yes	Medium	Low
<i>Eucalyptus educta</i>	P2				X	29.7	Apr	Shallow soils. Granite rocks. ²	Yes	Medium	Medium
<i>Goodenia salina</i>	P2				X	23.9	May or Aug - Nov	Well-drained, saline, grey or brown loamy clay. Low gypseous dunes near salt pans. ²	Yes	Medium	Low
<i>Hakea rigida</i>	P2		X		X	61.3	Sep to Oct.	Sandy soils, yellow sand. ²	Yes	Low	Low
<i>Lepidium merrallii</i>	P2		X			2.5	NA	Clay loam. ²	Yes	High	Medium
<i>Lepidosperma</i> sp. <i>Kambalda</i> (A.A. Mitchell 5156)	P2				X	45.2	Dec	Lower footslope of basalt hill. ²	No	Low	Low
<i>Phebalium clavatum</i>	P2		X		X	10.1	Aug - Sep	Sandy soils. Sandplains. ²	Yes	Medium	Low
<i>Phebalium</i> sp. <i>Yerilgee Sandplain</i> (J. Jackson 223)	P2				X	52.1	Aug	Yellow, orange/red sad, loam. Sandplains. ²	Yes	Low	Low
<i>Rumex crystallinus</i>	P2				X	55.3	Annual	Arid & semi-arid areas.	Yes	Low	Low
<i>Thysanotus</i> sp. <i>Yellowdine</i> (A.S. George 6040)	P2				X	61.5	Mar or Dec	Yellow sand, sandy clay. Sandplains, undulating ridges. ²	Yes	Low	Low
<i>Acacia crenulata</i>	P3				X	23.9	April - May or Nov	Clay, sandy clay, yellow sand. ²	Yes	Medium	Low
<i>Acacia cylindrica</i>	P3				X	63.2	Aug to Oct.	Yellow/brown sand, gravelly soils. Undulating plains, flats. ²	Yes	Low	Low
<i>Allocasuarina eriochlamys</i> subsp. <i>grossa</i>	P3		X			12.7	Jul	Stony loam, laterite clay. Granite outcrops. ²	No	High	Low
<i>Alyogyne</i> sp. <i>Great Victoria Desert</i> (D.J. Edinger 6212)	P3				X	2.5	Aug or Dec	Orange, yellow or red sand, sandy loams. Flat plains. ²	Yes	High	Low
<i>Alyxia tetanifolia</i>	P3				X	21.6	May - Jun or Nov	Sandy clay, loam, concretionary gravel. Drainage lines, near lakes. ²	Yes	Medium	Low
<i>Angianthus prostratus</i>	P3				X	46.7	Jul - Sep	Red clay or loamy soils. Saline depressions. ²	Yes	Low	Low
<i>Atriplex lindleyi</i> subsp. <i>conduplicata</i>	P3				X	55.1	NA	Crabhole plains, dry, yellow bare sandy clay, by lakes. ²	Yes	Low	Low

¹ Department of Agriculture, Water and Environment (2020) ²Western Australian Herbarium (2020)

Species	Conservation Status		Source			Distance to Nearest Record (km)	Flowering Period	Preferred Habitat	Habitat occurs within the Survey Area	Pre-Survey Likelihood of Occurrence	Post-Survey Likelihood of Occurrence
	DBCA	EPBC	NatureMap	PMST	DBCA						
<i>Austrostipa blackii</i>	P3		X			1.7	Sep - Nov	Orange, red or brown clay loam, silty sand, sandy clay loam. Gravel, basalt. Winter wet depressions, rocky outcrops, hills sides. ²	Yes	High	Recorded
<i>Austrostipa turbinata</i>	P3		X			8	Sep - Oct	Hill. Brown sandy clay. ²	Yes	High	Medium
<i>Bossiaea celata</i>	P3				X	23.3	Sep - Oct	Deep sand. Open mallee. ²	No	Medium	Low
<i>Bossiaea concinna</i>	P3				X	68.3	Jun - Sep	White or red sand, gravel. ²	Yes	Low	Low
<i>Calytrix creswellii</i>	P3				X	69.3	Sep to Dec.	Yellow sand, sometimes with lateritic gravel. Sandplains. ²	Yes	Low	Low
<i>Chrysocephalum apiculatum</i> subsp. <i>norsemanense</i>	P3		X			2.1	Aug - Oct	Well-drained loamy sand. Moderately exposed, gentle undulating plains, hills. ²	Yes	High	Medium
<i>Cratystylis centralis</i>	P3				X	39.2	Oct	Red sandy loam with ironstone gravel. Flat plains, breakaway country. ²	Yes	Low	Low
<i>Cryptandra crispula</i>	P3				X	42.7	Aug - Sep	Brown sandy clay, yellow loamy sand, red soil, pebbles. Dune ridges, hills, near salt lakes. ²	Yes	Low	Low
<i>Cyathostemon verrucosus</i>	P3				X	43.9	Apr or Jul or Aug or Oct	Yellow sand, gentle undulating plain. ²	Yes	Low	Low
<i>Eremophila annosocaulis</i>	P3				X	69.8	Sep	Stony, flat, sandy plain. Red sand. ²	Yes	Low	Low
<i>Eremophila arachnoides</i> subsp. <i>tenera</i>	P3				X	71.1	Oct - Dec	Undulating plains, Saline plains, drainage, clay. ²	Yes	Low	Low
<i>Eremophila veronica</i>	P3		X			0.5	Apr to May.	Stony clay, clay loam. Lateritic breakaways. ²	No	High	High
<i>Eucalyptus exigua</i>	P3				X	48.7	Mar - Apr or Oct or Dec	Sandy loam, white sand. Sandplains. ²	Yes	Low	Low
<i>Eucalyptus frenchiana</i>	P3				X	42.2	Oct	Brown or orange loam, sand or sandy clay. Granite. Flat plains, low undulating plains. ²	Yes	Low	Low
<i>Eutaxia actinophylla</i>	P3				X	64.8	Sep to Oct.	Red-brown clay loam, red clay loam over granite, gravel. Small depressions.	Yes	Low	Low
<i>Gompholobium cinereum</i>	P3				X	19.9	Feb or May or Sep - Nov	Yellow sand, clayey sand, brown loam, sandy gravel, laterite. Well-drained open sites, slopes, plains, roadsides. ²	Yes	Medium	Low
<i>Grevillea georgeana</i>	P3		X		X	1.1	Jan or Mar or Sep to Nov.	Stony loam/clay. Ironstone hilltops & slopes.	Yes	High	Medium

¹ Department of Agriculture, Water and Environment (2020) ²Western Australian Herbarium (2020)

Appendix D

Inventory of Vascular Flora

Appendix: Inventory of Vascular Flora

Family	Taxon
Aizoaceae	<i>Carpobrotus</i> sp.
Amaranthaceae	<i>Ptilotus exaltatus</i>
	<i>Ptilotus holosericeus</i>
	<i>Ptilotus obovatus</i> var. <i>obovatus</i>
	<i>Ptilotus polystachyus</i>
Anacardiaceae	* <i>Schinus molle</i> var. <i>areira</i>
Apocynaceae	<i>Alyxia buxifolia</i>
	<i>Leichhardtia australis</i>
	<i>Marsdenia australis</i>
	<i>Vincetoxicum lineare</i>
Asparagaceae	* <i>Agave americana</i>
	<i>Thysanotus manglesianus</i>
Asphodelaceae	* <i>Asphodelus fistulosus</i>
Asteraceae	* <i>Centaurea melitensis</i>
	<i>Chrysocephalum puteale</i>
	<i>Cratystylis conocephala</i>
	<i>Olearia muelleri</i>
	<i>Olearia pimeleoides</i>
	* <i>Oligocarpus calendulaceus</i>
	<i>Rhodanthe chlorocephala</i> subsp. <i>rosea</i>
	* <i>Sonchus asper</i>
	<i>Streptoglossa liatroides</i>
	<i>Vittadinia dissecta</i> var. <i>hirta</i>
	<i>Waitzia acuminata</i> var. <i>acuminata</i>
Boraginaceae	<i>Halgania andromedifolia</i>
	<i>Heliotropium curassavicum</i>
	* <i>Heliotropium europaeum</i>
Brassicaceae	* <i>Carrichtera annua</i>
Cactaceae	<i>Opuntia stricta</i>
Casuarinaceae	<i>Allocasuarina ?huegeliana</i>
	<i>Casuarina pauper</i>
Chenopodiaceae	<i>Atriplex ?holocarpa</i>
	<i>Atriplex ?vesicaria</i>
	<i>Atriplex nummularia</i> subsp. <i>spathulata</i>
	<i>Atriplex quadrivalvata</i>
	<i>Atriplex</i> sp.
	<i>Atriplex vesicaria</i>
	<i>Chenopodium curvispicatum</i>
	<i>Dysphania melanocarpa</i> forma <i>leucocarpa</i>
	? <i>Enchylaena tomentosa</i>
	<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>
	<i>Maireana ?georgei</i>
	<i>Maireana ?marginata</i>
	<i>Maireana georgei</i>
	<i>Maireana pentatropis</i>
	<i>Maireana sedifolia</i>
	<i>Maireana tomentosa</i>
	<i>Maireana trichoptera</i>
	<i>Maireana triptera</i>

Appendix: Inventory of Vascular Flora

Family	Taxon
Chenopodiaceae	<i>Osteocarpum salsuginosum</i>
	<i>Rhagodia drummondii</i>
	<i>Salsola australis</i>
	<i>Sclerolaena cuneata</i>
	<i>Sclerolaena diacantha</i>
	<i>Sclerolaena fusiformis</i>
	<i>Sclerolaena obliquicuspis</i>
	<i>Tecticornia halocnemoides</i>
Crassulaceae	* <i>Crassula ovata</i>
Euphorbiaceae	* <i>Euphorbia drummondii</i>
Fabaceae	<i>Acacia acuminata</i>
	<i>Acacia collegialis</i>
	<i>Acacia dissona</i> var. <i>dissona</i>
	<i>Acacia erinacea</i>
	<i>Acacia hemiteles</i>
	<i>Acacia jennerae</i>
	<i>Acacia tetragonophylla</i>
	<i>Acacia xerophila</i> var. <i>brevior</i>
	<i>Senna artemisioides</i> subsp. <i>artemisioides</i>
	<i>Senna artemisioides</i> subsp. <i>filifolia</i>
	<i>Senna artemisioides</i> subsp. x <i>artemisioides</i>
	<i>Senna stowardii</i>
Frankeniaceae	<i>Frankenia ?fecunda</i>
Goodeniaceae	<i>Goodenia havilandii</i>
	<i>Goodenia pinnatifida</i>
	<i>Scaevola spinescens</i>
Haloragaceae	<i>Haloragis trigonocarpa</i>
Hemerocallidaceae	<i>Dianella revoluta</i> var. <i>divaricata</i>
Lamiaceae	* <i>Salvia verbenaca</i>
	<i>Westringia rigida</i>
Malvaceae	<i>Abutilon cryptopetalum</i>
	<i>Hibiscus solanifolius</i>
	<i>Hibiscus sturtii</i> var. <i>grandiflorus</i>
	<i>Lawrenzia repens</i>
	<i>Malvaceae</i> sp.
	<i>Sida petrophila</i>
	<i>Sida ?petrophila</i>
	<i>Calothamnus ?gracilis</i>
	<i>Eucalyptus ?ravida</i>
	<i>Eucalyptus campaspe</i>
	<i>Eucalyptus celastroides</i>
	<i>Eucalyptus clelandiorum</i>
	<i>Eucalyptus griffithsii</i>
	<i>Eucalyptus loxophleba</i> subsp. <i>lissophloia</i>
	<i>Eucalyptus oldfieldii</i>
	<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>
	<i>Eucalyptus salmonophloia</i>
	<i>Eucalyptus salubris</i>
	<i>Eucalyptus torquata</i>

Appendix: Inventory of Vascular Flora

Family	Taxon
Myrtaceae	<i>Eucalyptus websteriana</i> subsp. <i>websteriana</i>
	<i>Melaleuca ?hamata</i>
	<i>Melaleuca pauperiflora</i> subsp. <i>fastigiata</i>
Pittosporaceae	<i>Pittosporum angustifolium</i>
Poaceae	<i>Aristida contorta</i>
	<i>Austrostipa blackii</i> (P3)
	<i>Austrostipa platychaeta</i>
	<i>Austrostipa</i> sp.
	<i>Austrostipa trichophylla</i>
	<i>Chloris truncata</i>
	<i>Enneapogon caerulescens</i>
	<i>Eragrostis dielsii</i>
	<i>Monachather paradoxus</i>
	<i>Rytidosperma</i> sp.
	<i>Triodia desertorum</i>
Polygonaceae	* <i>Rumex vesicarius</i>
Portulacaceae	<i>Calandrinia baccata</i>
	<i>Calandrinia eremaea</i>
Pteridaceae	<i>Cheilanthes lasiophylla</i>
	<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>
Rhamnaceae	<i>Trymalium myrtillus</i> subsp. <i>myrtillus</i>
Rutaceae	<i>Phebalium laevigatum</i>
Santalaceae	<i>Exocarpos aphyllus</i>
	<i>Santalum acuminatum</i>
	<i>Santalum spicatum</i>
Sapindaceae	<i>Dodonaea adenophora</i>
	<i>Dodonaea lobulata</i>
	<i>Dodonaea stenozyga</i>
Scrophulariaceae	<i>Eremophila alternifolia</i>
	<i>Eremophila deserti</i>
	<i>Eremophila forrestii</i>
	<i>Eremophila georgei</i>
	<i>Eremophila glabra</i> subsp. <i>glabra</i>
	<i>Eremophila interstans</i> subsp. <i>interstans</i>
	<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>
	<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>
	<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>
	<i>Eremophila scoparia</i>
	<i>Myoporum platycarpum</i> subsp. <i>platycarpum</i>
Solanaceae	<i>Lycium australe</i>
	* <i>Nicotiana glauca</i>
	<i>Nicotiana rotundifolia</i>
	<i>Solanum cleistogamum</i>
	<i>Solanum lasiophyllum</i>
	<i>Solanum nummularium</i>
Thymelaeaceae	? <i>Pimelea</i> sp.
	<i>Pimelea microcephala</i> subsp. <i>microcephala</i>
	<i>Pimelea spiculigera</i> var. <i>thesioides</i>
Zygophyllaceae	<i>Roepera glauca</i>

Appendix: Inventory of Vascular Flora

Family	Taxon
Zygophyllaceae	<i>Roepera ovata</i>
	<i>Roepera</i> sp.

Appendix E

Flora Site Data

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q01
Location MGA 50 326500 mE 6578134 mN

Described by: JW
Date: 12/10/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite, quartz



Vegetation: *Eremophila georgei* tall open shrubland over *Dodonaea lobulata* mid open shrubland over *Ptilotus obovatus* var. *obovatus*, *Sida petrophila* and *Vincetoxicum lineare* low open herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Abutilon cryptopetalum</i>	10	0.1	
<i>Aristida contorta</i>	10	0.1	
<i>Austrostipa blackii</i> (P3)	10	1	
<i>Calandrinia eremaea</i>	10	0.1	
<i>Dodonaea lobulata</i>	150	12	
<i>Eremophila georgei</i>	300	28	
<i>Nicotiana rotundifolia</i>	10	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	16	
<i>Sida petrophila</i>	25	11	
<i>Solanum cleistogamum</i>	10	0.1	
<i>Solanum lasiophyllum</i>	25	1	
<i>Vincetoxicum lineare</i>	10	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q02
Location MGA 50 326422 mE 6578029 mN

Described by: JW
Date: 12/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus celastroides* and *Eucalyptus clelandiorum* low woodland over *Eremophila scoparia* mid sparse shrubland over *Atriplex* sp., *Atriplex vesicaria* and *Atriplex nummularia* subsp. *spathulata* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	35	6	
<i>Atriplex</i> sp.	35	16	
<i>Atriplex vesicaria</i>	50	7	
<i>Dodonaea stenozyga</i>	25	1	
<i>Eremophila scoparia</i>	120	6	
<i>Eucalyptus celastroides</i>	500	16	
<i>Eucalyptus clelandiorum</i>	1000	5	
<i>Exocarpos aphyllus</i>	45	1	
<i>Haloragis trigonocarpa</i>	25	0.5	
<i>Lycium australe</i>	40	2	
<i>Ptilotus holosericeus</i>	5	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	1	
<i>Roepera</i> sp.	10	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q03
Location MGA 50 327421 mE 6577933 mN

Described by: JW
Date: 12/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus campaspe* low woodland over *Eremophila scoparia* mid sparse shrubland over *Senna artemisioides* subsp. *x artemisioides*, *Atriplex* sp. and *Olearia muelleri* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex</i> sp.	40	6	
* <i>Carrichtera annua</i>	25	2	
<i>Eremophila scoparia</i>	140	3	
<i>Eucalyptus campaspe</i>	300	26	
<i>Olearia muelleri</i>	50	6	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	2	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	100	12	
<i>Streptoglossa liatroides</i>	5	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q04
Location MGA 50 327671 mE 6577991 mN

Described by: JW
Date: 12/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: N/A



Vegetation: *Exocarpos aphyllus*, *Eremophila oldfieldii* subsp. *angustifolia* and *Acacia tetragonophylla* mid shrubland over *Dodonaea lobulata* mid open shrubland over *Scaevola spinescens*, *Olearia muelleri* and *Eremophila alternifolia* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia tetragonophylla</i>	180	6	
<i>Dodonaea lobulata</i>	110	12	
<i>Eremophila alternifolia</i>	50	1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	200	12	
<i>Exocarpos aphyllus</i>	200	26	
<i>Lycium australe</i>	5	0.1	
<i>Maireana georgei</i>	18	1	
<i>Marsdenia australis</i>	10	1	
<i>Olearia muelleri</i>	45	2	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	1	
<i>Scaevola spinescens</i>	100	6	
<i>Thysanotus manglesianus</i>	100	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q05
Location MGA 50 327597 mE 6577741 mN

Described by: JW
Date: 12/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus clelandiorum* and *Eucalyptus campaspe* low open woodland over *Eremophila scoparia* and *Senna artemisioides* subsp. *x artemisioides* mid open shrubland over *Cheilanthes lasiophylla* and *Ptilotus obovatus* var. *obovatus* low sparse herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Cheilanthes lasiophylla</i>	15	4	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	35	0.1	
<i>Eremophila scoparia</i>	200	20	
<i>Eucalyptus campaspe</i>	300	4	
<i>Eucalyptus clelandiorum</i>	500	5	
<i>Maireana</i> ? <i>marginata</i>	5	1	
<i>Olearia muelleri</i>	20	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	1	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	130	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q06
Location MGA 50 327323 mE 6576260 mN

Described by: JW
Date: 12/10/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite, quartz



Vegetation: *Exocarpos aphyllus* tall open shrubland over *Dodonaea lobulata* and *Dodonaea adenophora* low shrubland over *Haloragis trigonocarpa*, *Thysanotus manglesianus* and *Ptilotus obovatus* var. *obovatus* low open herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Dodonaea adenophora</i>	100	12	
<i>Dodonaea lobulata</i>	90	25	
<i>Exocarpos aphyllus</i>	220	11	
<i>Haloragis trigonocarpa</i>	10	11	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	1	
<i>Thysanotus manglesianus</i>	20	2	
<i>Waitzia acuminata</i> var. <i>acuminata</i>	18	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q07
Location MGA 50 328895 mE 6577682 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus torquata* low woodland over *Exocarpos aphyllus* mid sparse shrubland over *Westringia rigida*, *Olearia muelleri* and *Scaevola spinescens* low shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	40	5	
<i>Eucalyptus torquata</i>	400	12	
<i>Exocarpos aphyllus</i>	110	1	
<i>Olearia muelleri</i>	25	11	
<i>Scaevola spinescens</i>	40	6	
<i>Westringia rigida</i>	40	16	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q08
Location MGA 50 328889 mE 6577825 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: N/A



Vegetation: *Eucalyptus campaspe* and *Eucalyptus griffithsii* low open woodland over *Eremophila scoparia*, *Senna artemisioides* subsp. *filifolia* and *Exocarpos aphyllus* mid shrubland over *Senna artemisioides* subsp. *x artemisioides*, *Acacia erinacea* and *Olearia muelleri* low shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia erinacea</i>	40	13	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	200	4	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	35	1	
<i>Eremophila scoparia</i>	140	23	
<i>Eucalyptus campaspe</i>	400	5	
<i>Eucalyptus griffithsii</i>	500	5	
<i>Exocarpos aphyllus</i>	200	8	
<i>Lycium australe</i>	100	0.1	
<i>Maireana georgei</i>	20	0.1	
<i>Myoporum platycarpum</i> subsp. <i>platycarpum</i>	100	1	
<i>Olearia muelleri</i>	35	3	
<i>Santalum acuminatum</i>	55	1	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	100	15	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	140	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q09
Location MGA 50 328932 mE 6578002 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite



Vegetation: *Eucalyptus oleosa* subsp. *oleosa* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia*, *Senna artemisioides* subsp. *filifolia* and *Trymalium myrtillus* subsp. *myrtillus* mid shrubland over *Westringia rigida*, *Eremophila parvifolia* subsp. *auricampi* and *Calothamnus ?gracilis* low shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	25	1	
<i>Alyxia buxifolia</i>	40	1	
<i>Atriplex vesicaria</i>	60	5	
<i>Calothamnus ?gracilis</i>	70	6	
<i>Dodonaea stenozyga</i>	185	5	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	200	15	
<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>	130	5	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	50	12	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	700	10	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	200	11	
<i>Trymalium myrtillus</i> subsp. <i>myrtillus</i>	140	11	
<i>Westringia rigida</i>	25	15	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q10
Location MGA 50 328368 mE 6578263 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: N/A



Vegetation: *Eucalyptus griffithsii* low open woodland over *Acacia jennerae*, *Eremophila oldfieldii* subsp. *angustifolia* and *Exocarpos aphyllus* mid open shrubland over *Eremophila glabra* subsp. *glabra*, *Dodonaea stenozyga* and *Olearia pimeleoides* low shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	110	10	
<i>Aristida contorta</i>	100	1	
<i>Dodonaea stenozyga</i>	100	6	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	60	18	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	120	5	
<i>Eucalyptus griffithsii</i>	400	5	
<i>Exocarpos aphyllus</i>	130	5	
<i>Olearia pimeleoides</i>	25	6	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	10	0.1	
<i>Santalum spicatum</i>	180	3	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	120	5	
<i>Westringia rigida</i>	20	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q20
Location MGA 50 328215 mE 6578113 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Eremophila interstans* subsp. *interstans* mid open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	100	5	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	120	15	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	300	19	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	45	6	
<i>Eucalyptus griffithsii</i>	400	2	
<i>Maireana sedifolia</i>	80	3	
<i>Olearia pimeleoides</i>	30	5	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	100	2	
<i>Sida petrophila</i>	28	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q21
Location MGA 50 326536 mE 6577327 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite



Vegetation: *Dodonaea stenozyga*, *Senna artemisioides* subsp. *filifolia* and *Eremophila oldfieldii* subsp. *angustifolia* mid closed shrubland over *Atriplex vesicaria* low sparse shrubland over *Ptilotus obovatus* var. *obovatus* low open herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	80	5	
<i>Dodonaea stenozyga</i>	180	40	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	140	15	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	28	16	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	180	20	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q22
Location MGA 50 326045 mE 6577335 mN

Described by: JW
Date: 13/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus salmonophloia* low open woodland over *Senna artemisioides* subsp. *x artemisioides* and *Senna artemisioides* subsp. *filifolia* low sparse shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eucalyptus salmonophloia</i>	600	5	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	45	6	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	35	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q23
Location MGA 50 326813 mE 6577450 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus salmonophloia* low woodland over *Senna artemisioides* subsp. *x artemisioides*, *Senna artemisioides* subsp. *filifolia* and *Eremophila parvifolia* subsp. *auricampi* low sparse shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	35	1	
<i>Eucalyptus salmonophloia</i>	1000	19	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	100	2	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	75	2	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q24
Location MGA 50 326578 mE 6577758 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus campaspe* low woodland over *Senna artemisioides* subsp. *x artemisioides* mid sparse shrubland over *Senna artemisioides* subsp. *filifolia* low sparse shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eucalyptus campaspe</i>	400	15	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	130	2	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	45	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q25
Location MGA 50 326770 mE 6578110 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Upper slope
Rock Type: N/A



Vegetation: *Acacia acuminata* tall open shrubland over *Eremophila oldfieldii* subsp. *angustifolia* mid sparse shrubland over *Sida petrophila*, *Solanum lasiophyllum* and *Dodonaea stenozyga* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	400	20	
<i>Austrostipa</i> sp.	10	1	
<i>Dodonaea stenozyga</i>	50	1	
<i>Eremophila georgei</i>	20	0.1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	130	10	
<i>Haloragis trigonocarpa</i>	10	3	
<i>Ptilotus exaltatus</i>	30	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	30	0.1	
<i>Sida petrophila</i>	100	5	
<i>Solanum lasiophyllum</i>	30	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q26
Location MGA 50 327707 mE 6577970 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: N/A



Vegetation: *Eremophila oldfieldii* subsp. *angustifolia*, *Dodonaea lobulata* and *Senna artemisioides* subsp. *x artemisioides* mid shrubland over *Ptilotus obovatus* var. *obovatus* low sparse herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Dodonaea lobulata</i>	140	12	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	190	20	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	1	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	170	12	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	200	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q27
Location MGA 50 327762 mE 6578035 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: N/A



Vegetation: *Eucalyptus torquata* low open woodland over *Eremophila interstans* subsp. *interstans*, *Senna artemisioides* subsp. *filifolia* and *Olearia muelleri* low sparse shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila interstans</i> subsp. <i>interstans</i>	50	2	
<i>Eucalyptus torquata</i>	500	8	
<i>Exocarpos aphyllus</i>	25	0.1	
<i>Maireana tomentosa</i>	10	1	
<i>Olearia muelleri</i>	35	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	55	2	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q28
Location MGA 50 327436 mE 6575925 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Gentle slope
Rock Type: Laterite



Vegetation: *Acacia acuminata* tall open shrubland over *Eremophila oldfieldii* subsp. *angustifolia* mid sparse shrubland over *Dodonaea stenozyga* and *Senna artemisioides* subsp. *filifolia* low closed shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	300	15	
<i>Austrostipa platychaeta</i>	18	2	
<i>Calandrinia baccata</i>	10	2	
<i>Dodonaea stenozyga</i>	80	70	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	120	6	
<i>Goodenia havilandii</i>	18	1	
<i>Haloragis trigonocarpa</i>	18	1	
<i>Lycium australe</i>	10	0.1	
<i>Maireana pentatropis</i>	10	0.1	
<i>Ptilotus exaltatus</i>	25	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	40	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	100	5	
<i>Thysanotus manglesianus</i>	20	3	
<i>Waitzia acuminata</i> var. <i>acuminata</i>	15	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q29
Location MGA 50 328203 mE 6575013 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus torquata* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia*, *Senna artemisioides* subsp. *x artemisioides* and *Dodonaea stenozyga* mid open shrubland over *Ptilotus obovatus* var. *obovatus* and *Ptilotus exaltatus* low sparse herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	45	1	
<i>Dodonaea stenozyga</i>	150	3	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	200	6	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	35	3	
<i>Eucalyptus torquata</i>	400	5	
<i>Olearia muelleri</i>	25	1	
<i>Ptilotus exaltatus</i>	20	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	3	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	190	6	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	190	3	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1Q30
Location MGA 50 328220 mE 6575082 mN

Described by: JW
Date: 14/10/2021
Type: QUADRAT

Landform: Rocky plain
Rock Type: Laterite



Vegetation: *Allocasuarina ?huegeliana* and *Eucalyptus torquata* low woodland over *Santalum spicatum* and *Eremophila oppositifolia* subsp. *angustifolia* mid sparse shrubland over *Dodonaea stenozyga*, *Eremophila glabra* subsp. *glabra* and *Atriplex vesicaria* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Allocasuarina ?huegeliana</i>	400	6	
<i>Atriplex vesicaria</i>	100	3	
<i>Dodonaea stenozyga</i>	80	6	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	90	6	
<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>	190	1	
<i>Eucalyptus torquata</i>	700	5	
<i>Maireana pentatropis</i>	25	0.1	
<i>Ptilotus exaltatus</i>	20	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	1	
<i>Santalum spicatum</i>	150	2	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	100	3	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R01
Location MGA 50 328967 mE 6577693 mN

Described by: JW
Date: 13/10/2021
Type: RELEVE

Landform: Plain
Rock Type: N/A



Vegetation: *Eucalyptus torquata* low open woodland over *Exocarpos aphyllus* tall sparse shrubland over *Eremophila parvifolia* subsp. *auricampi*, *Westringia rigida* and *Senna artemisioides* subsp. *filifolia* low sparse shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	32	3	
<i>Eucalyptus torquata</i>	400	5	
<i>Exocarpos aphyllus</i>	220	6	
<i>Olearia muelleri</i>	40	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	35	2	
<i>Westringia rigida</i>	35	3	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R02
Location MGA 50 328922 mE 6577745 mN

Described by: JW
Date: 13/10/2021
Type: RELEVÉ

Landform: Plain
Rock Type: N/A



Vegetation: *Eucalyptus torquata* low woodland over *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Senna artemisioides* subsp. *filifolia*, *Olearia muelleri* and *Westringia rigida* low open shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila georgei</i>	28	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	200	4	
<i>Eucalyptus torquata</i>	500	16	
<i>Olearia muelleri</i>	25	4	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	40	6	
<i>Westringia rigida</i>	25	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R03
Location MGA 50 329002 mE 6577925 mN

Described by: JW
Date: 13/10/2021
Type: RELEVE

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus torquata* low open woodland over *Eremophila alternifolia* tall sparse shrubland over *Dodonaea stenozyga* and *Eremophila oldfieldii* subsp. *angustifolia* mid sparse shrubland

Condition: Excellent **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	60	2	
<i>Dodonaea stenozyga</i>	120	5	
<i>Eremophila alternifolia</i>	210	2	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	110	5	
<i>Eucalyptus torquata</i>	400	2	
<i>Maireana tomentosa</i>	25	1	
<i>Westringia rigida</i>	25	2	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R04
Location MGA 50 326478 mE 6577408 mN

Described by: JW
Date: 13/10/2021
Type: RELEVÉ

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Senna artemisioides* subsp. *x* *artemisioides* and *Senna artemisioides* subsp. *filifolia* mid open shrubland over *Ptilotus obovatus* var. *obovatus* low open herbland
Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Dodonaea stenozyga</i>	100	12	
<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>	45	1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	15	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	25	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	45	28	
<i>Senna artemisioides</i> subsp. <i>x</i> <i>artemisioides</i>	120	10	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	110	6	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R05
Location MGA 50 326536 mE 6577327 mN

Described by: JW
Date: 13/10/2021
Type: RELEVÉ

Landform: Mid slope
Rock Type: N/A



Vegetation: *Eucalyptus torquata* and *Eucalyptus salmonophloia* low woodland over *Exocarpos aphyllus* mid sparse shrubland over *Senna artemisioides* subsp. *filifolia* and *Senna artemisioides* subsp. *x artemisioides* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eucalyptus salmonophloia</i>	500	10	
<i>Eucalyptus torquata</i>	400	5	
<i>Exocarpos aphyllus</i>	140	2	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	45	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	90	11	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R09
Location MGA 50 326358 mE 6577538 mN

Described by: JW
Date: 14/10/2021
Type: RELEVE

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus celastroides* and *Eucalyptus clelandiorum* low woodland over *Atriplex vesicaria* low sparse shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	40	1	
<i>Eucalyptus celastroides</i>	400	6	
<i>Eucalyptus clelandiorum</i>	400	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R10
Location MGA 50 326982 mE 6578150 mN

Described by: JW
Date: 14/10/2021
Type: RELEVE

Landform: Undulating plain
Rock Type: N/A



Vegetation: *Eucalyptus clelandiorum* low open woodland over *Senna artemisioides* subsp. *x artemisioides*, *Atriplex vesicaria* and *Eremophila interstans* subsp. *interstans* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	28	1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	45	1	
<i>Eucalyptus clelandiorum</i>	1000	5	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	35	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R11
Location MGA 50 327829 mE 6578083 mN

Described by: JW
Date: 14/10/2021
Type: RELEVÉ

Landform: Plain
Rock Type: Laterite, quartz



Vegetation: *Eucalyptus torquata* mid open woodland over *Eremophila interstans* subsp. *interstans* mid open shrubland over *Senna artemisioides* subsp. *x artemisioides*, *Senna artemisioides* subsp. *filifolia* and *Acacia jennerae* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	10	3	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	150	16	
<i>Eucalyptus torquata</i>	1500	5	
<i>Olearia muelleri</i>	35	2	
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	40	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	25	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C1R13
Location MGA 50 327488 mE 6575973 mN

Described by: JW
Date: 14/10/2021
Type: RELEVÉ

Landform: Undulating plain
Rock Type: N/A



Vegetation: *Eucalyptus clelandiorum*, *Eucalyptus salmonophloia* and *Eucalyptus griffithsii* low woodland over *Eremophila interstans* subsp. *interstans* and *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Atriplex nummularia* subsp. *spathulata* and *Dodonaea stenozyga* low open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	45	11	
<i>Dodonaea stenozyga</i>	10	6	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	220	11	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	200	10	
<i>Eucalyptus clelandiorum</i>	500	6	
<i>Eucalyptus griffithsii</i>	300	5	
<i>Eucalyptus salmonophloia</i>	400	6	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q01
Location MGA 50 326721 mE 6571821 mN

Described by: JW
Date: 15/11/2021
Type: QUADRAT

Landform: Scree slope drain line No photo available
Rock Type: Laterite

Vegetation: *Eucalyptus loxophleba* subsp. *lissophloia* and *Eucalyptus campaspe* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* mid sparse shrubland over *Dodonaea adenophora* and *Olearia muelleri* low sparse shrubland
Condition: Good **Disturbance Type:** Fauna tracks/scats, Infrastructure

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Dodonaea adenophora</i>	35	1	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	5	0.1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	130	2	
<i>Eucalyptus campaspe</i>	300	1	
<i>Eucalyptus loxophleba</i> subsp. <i>lissophloia</i>	300	2	
<i>Maireana georgei</i>	15	0.1	
<i>Olearia muelleri</i>	25	0.5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q02
Location MGA 50 324588 mE 6571695 mN

Described by: BD,SW
Date: 15/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Ironstone



Vegetation: *Eucalyptus salmonophloia* low woodland over *Melaleuca ?hamata* and *Eremophila oppositifolia* subsp. *angustifolia* tall open shrubland over *Olearia muelleri* low sparse shrubland

Condition: Very Good **Disturbance Type:** Litter

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	40	0.1	
<i>Acacia hemiteles</i>	160	1	
<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>	40	0.1	
<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>	250	4	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	60	0.1	
<i>Eucalyptus salmonophloia</i>	650	12	
<i>Exocarpos aphyllus</i>	150	0.5	
<i>Melaleuca ?hamata</i>	300	11	
<i>Olearia muelleri</i>	45	5	
<i>Olearia pimeleoides</i>	30	0.1	
<i>Pimelea microcephala</i> subsp. <i>microcephala</i>	50	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	0.1	
<i>Scaevola spinescens</i>	120	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q03
Location MGA 50 327534 mE 6572097 mN

Described by: JW
Date: 15/11/2021
Type: QUADRAT

Landform: Drainage line No photo available
Rock Type: Laterite

Vegetation: *Atriplex ?vesicaria and Senna artemisioides subsp. x artemisioides low sparse shrubland*

Condition: Good **Disturbance Type:** Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	100	5	
<i>Eremophila forrestii</i>	20	0.1	
<i>Eremophila scoparia</i>	130	0.1	
<i>Maireana georgei</i>	5	0.1	
<i>Senna artemisioides subsp. x artemisioides</i>	60	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q04
Location MGA 50 324603 mE 6571566 mN

Described by: BD,SW
Date: 15/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Ironstone



Vegetation: *Eucalyptus ?ravida* low open forest over *Eremophila interstans* subsp. *interstans*, *Melaleuca ?hamata* and *Melaleuca pauperiflora* subsp. *fastigiata* tall open shrubland over *Triodia desertorum* low sparse hummock grassland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	170	0.1	
<i>Acacia hemiteles</i>	140	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	35	0.1	
<i>Dodonaea lobulata</i>	100	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	210	11	
<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>	110	0.1	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	30	0.1	
<i>Eucalyptus ?ravida</i>	550	35	
<i>Exocarpos aphyllus</i>	90	0.1	
<i>Halgania andromedifolia</i>	70	6	
<i>Maireana georgei</i>	30	0.1	
<i>Melaleuca ?hamata</i>	250	2	
<i>Melaleuca pauperiflora</i> subsp. <i>fastigiata</i>	350	1	
<i>Myoporum platycarpum</i> subsp. <i>platycarpum</i>	210	1	
<i>Olearia muelleri</i>	30	0.1	
<i>Phebalium laevigatum</i>	120	1	
<i>Scaevola spinescens</i>	45	5	
<i>Sclerolaena fusiformis</i>	10	0.1	
<i>Triodia desertorum</i>	25	2	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q05
Location MGA 50 327403 mE 6572169 mN

Described by: JW
Date: 15/11/2021
Type: QUADRAT

Landform: Scree slope minor drain No photo available
Rock Type: Laterite

Vegetation: *Eucalyptus campaspe* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Dodonaea stenozyga*, *Alyxia buxifolia* and *Eremophila glabra* subsp. *glabra* low open shrubland
Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Alyxia buxifolia</i>	70	1	
<i>Dodonaea stenozyga</i>	90	15	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	100	1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	200	15	
<i>Eucalyptus campaspe</i>	200	1	
<i>Maireana georgei</i>	15	0.1	
<i>Olearia muelleri</i>	20	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	110	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q06
Location MGA 50 324605 mE 6571453 mN

Described by: BD,SW
Date: 15/11/2021
Type: QUADRAT

Landform: Low rise
Rock Type: Ironstone, Quartz



Vegetation: *Eucalyptus clelandiorum* and *Eucalyptus ?ravida* low open woodland over *Eremophila oppositifolia* subsp. *angustifolia* and *Myoporum platycarpum* subsp. *platycarpum* mid open shrubland over *Scaevola spinescens*, *Dodonaea lobulata* and *Phebalium laevigatum* low open shrubland

Condition: Good **Disturbance Type:** Litter, Historical Clearing

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	120	0.1	
<i>Acacia tetragonophylla</i>	150	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	120	0.1	
<i>Dodonaea lobulata</i>	100	1	
<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>	140	8	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	30	0.1	
<i>Eucalyptus ?ravida</i>	450	2	
<i>Eucalyptus clelandiorum</i>	700	5	
<i>Halgania andromedifolia</i>	100	0.1	
<i>Maireana georgei</i>	15	0.1	
<i>Myoporum platycarpum</i> subsp. <i>platycarpum</i>	190	4	
<i>Olearia muelleri</i>	30	0.5	
<i>Phebalium laevigatum</i>	80	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	0.1	
<i>Scaevola spinescens</i>	90	20	
<i>Solanum lasiophyllum</i>	20	0.1	
<i>Triodia desertorum</i>	30	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q07
Location MGA 50 327549 mE 6571711 mN

Described by: JW
Date: 15/11/2021
Type: QUADRAT

Landform: Scree slope No photo available
Rock Type: Laterite

Vegetation: *Eremophila interstans* subsp. *interstans* tall open shrubland over *Dodonaea stenozyga*, *Senna artemisioides* subsp. *filifolia* and *Alyxia buxifolia* mid open shrubland over *Maireana georgei* low sparse chenopod shrubland
Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Alyxia buxifolia</i>	110	1	
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	25	0.1	
<i>Dodonaea stenozyga</i>	170	5	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	250	15	
<i>Maireana georgei</i>	15	2	
<i>Malvaceae</i> sp.	5	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	190	5	
<i>Solanum cleistogamum</i>	15	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q08
Location MGA 50 324706 mE 6571242 mN

Described by: BD,SW
Date: 15/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Ironstone, Quartz



Vegetation: *Eucalyptus clelandiorum* and *Eucalyptus salmonophloia* mid open woodland over *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Maireana trichoptera*, *Atriplex ?vesicaria* and *Sclerolaena fusiformis* low sparse chenopod shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	30	1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	100	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	160	2	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	40	1	
<i>Eucalyptus clelandiorum</i>	1100	2	
<i>Eucalyptus salmonophloia</i>	1100	2	
<i>Exocarpos aphyllus</i>	40	0.1	
<i>Maireana georgei</i>	15	0.1	
<i>Maireana trichoptera</i>	8	2	
<i>Olearia muelleri</i>	35	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	40	0.1	
<i>Scaevola spinescens</i>	80	1	
<i>Sclerolaena diacantha</i>	10	0.5	
<i>Sclerolaena fusiformis</i>	5	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	100	0.1	
<i>Senna stowardii</i>	60	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q09
Location MGA 50 327372 mE 6571654 mN

Described by: JW
Date: 15/11/2021
Type: QUADRAT

Landform: Slope No photo available
Rock Type: Laterite

Vegetation: *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Dodonaea stenozyga* and *Acacia acuminata* mid sparse shrubland over *Ptilotus obovatus* var. *obovatus* and *Cheilanthes sieberi* subsp. *sieberi* low sparse herbland
Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	140	2	
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	35	0.5	
<i>Dodonaea stenozyga</i>	200	10	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	25	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	25	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q10
Location MGA 50 324887 mE 6571294 mN

Described by: BD,SW
Date: 15/11/2021
Type: QUADRAT

Landform: Drainage line
Rock Type: Calcrete, Ironstone, Quartz



Vegetation: *Eucalyptus salubris* low woodland over *Eremophila oldfieldii* subsp. *angustifolia*, *Dodonaea lobulata* and *Eremophila glabra* subsp. *glabra* mid open shrubland over *Atriplex ?vesicaria* low sparse chenopod shrubland

Condition: Very Good **Disturbance Type:** Litter

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	40	3	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	150	0.1	
<i>Dodonaea lobulata</i>	130	4	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	150	4	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	210	0.1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	180	11	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	50	0.1	
<i>Eucalyptus salubris</i>	900	20	
<i>Exocarpos aphyllus</i>	150	0.1	
<i>Maireana georgei</i>	30	0.1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Olearia muelleri</i>	35	0.1	
<i>Ptilotus exaltatus</i>	3	0.1	
<i>Scaevola spinescens</i>	100	0.1	
<i>Sclerolaena obliquicuspis</i>	10	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	150	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q11
Location MGA 50 326763 mE 6571675 mN

Described by: JW
Date: 15/11/2021
Type: QUADRAT

Landform: Drainage line
Rock Type: Granite, Laterite, Quartz



Vegetation: *Eucalyptus oleosa* subsp. *oleosa* low woodland over *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Dodonaea stenozyga*, *Acacia jennerae* and *Eremophila glabra* subsp. *glabra* low sparse shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia jennerae</i>	35	1	
<i>Dodonaea stenozyga</i>	38	5	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	45	1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	200	1	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	500	15	
<i>Maireana georgei</i>	20	0.1	
<i>Olearia muelleri</i>	40	1	
<i>Ptilotus exaltatus</i>	15	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q12
Location MGA 50 325065 mE 6571260 mN

Described by: BD,SW
Date: 15/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Ironstone, Laterite, Quartz



Vegetation: *Eucalyptus salmonophloia* mid woodland over *Senna artemisioides* subsp. *filifolia* mid isolated shrubs

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila interstans</i> subsp. <i>interstans</i>	400	0.1	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	25	0.1	
<i>Eucalyptus salmonophloia</i>	1500	25	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	150	0.5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q13
Location MGA 50 327631 mE 6571898 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Ridge crest
Rock Type: Laterite



Vegetation: *Eucalyptus oldfieldii* low open woodland over *Dodonaea stenozyga* mid sparse shrubland over *Atriplex vesicaria*, *Senna artemisioides* subsp. *filifolia* and *Maireana pentatropis* low shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	35	15	
<i>Dodonaea stenozyga</i>	110	8	
<i>Eucalyptus oldfieldii</i>	300	5	
<i>Maireana pentatropis</i>	5	5	
<i>Olearia muelleri</i>	20	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	6	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	45	15	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q14
Location MGA 50 325050 mE 6570941 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Ironstone



Vegetation: *Acacia collegialis* and *Eremophila oldfieldii* subsp. *angustifolia* tall sparse shrubland over *Dodonaea lobulata* and *Eremophila alternifolia* mid sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	400	6	
<i>Acacia tetragonophylla</i>	100	0.1	
<i>Chenopodium curvispicatum</i>	10	0.1	
<i>Dodonaea lobulata</i>	180	5	
<i>Eremophila alternifolia</i>	190	4	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	1	
<i>Maireana georgei</i>	20	0.1	
<i>Maireana trichoptera</i>	8	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	25	0.1	
<i>Rhagodia drummondii</i>	5	0.1	
<i>Sclerolaena diacantha</i>	5	0.1	
<i>Sclerolaena fusiformis</i>	10	0.1	
<i>Sida ?petrophila</i>	15	0.1	
<i>Solanum lasiophyllum</i>	10	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q15
Location MGA 50 327026 mE 6571767 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Lower slope
Rock Type: Laterite



Vegetation: *Eucalyptus clelandiorum* and *Eucalyptus oldfieldii* low open woodland over *Atriplex ?vesicaria* and *Eremophila interstans* subsp. *interstans* mid shrubland over *Dodonaea stenozyga*, *Olearia muelleri* and *Maireana pentatropis* low open shrubland

Condition: Good **Disturbance Type:** Fauna tracks/scats, Historical Clearing

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	120	35	
<i>Dodonaea stenozyga</i>	90	6	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	190	20	
<i>Eucalyptus clelandiorum</i>	400	2	
<i>Eucalyptus oldfieldii</i>	60	2	
<i>Maireana georgei</i>	5	0.1	
<i>Maireana pentatropis</i>	20	1	
<i>Olearia muelleri</i>	20	5	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q16
Location MGA 50 324948 mE 6570710 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Ironstone



Vegetation: *Eucalyptus griffithsii* mid woodland over *Eremophila oldfieldii* subsp. *angustifolia* tall sparse shrubland over *Eremophila interstans* subsp. *interstans* mid sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	30	0.1	
<i>Dodonaea lobulata</i>	80	1	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	40	0.5	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	180	3	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	500	2	
<i>Eucalyptus griffithsii</i>	1100	25	
<i>Maireana trichoptera</i>	20	0.1	
<i>Olearia muelleri</i>	40	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	25	0.1	
<i>Rhagodia drummondii</i>	20	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	40	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q17
Location MGA 50 327349 mE 6571149 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, Sandstone, Shale



Vegetation: *Atriplex vesicaria*, *Maireana georgei* and *Maireana pentatropis* low chenopod shrubland over **Asphodelus fistulosus* low sparse herbland

Condition: Poor **Disturbance Type:** Vehicle tracks, Fauna tracks/scats, Historical Clearing, Infrastructure

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>*Asphodelus fistulosus</i>	35	10	
<i>Atriplex vesicaria</i>	100	26	
<i>Maireana georgei</i>	10	15	
<i>Maireana pentatropis</i>	20	10	
<i>*Salvia verbenaca</i>	10	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q18
Location MGA 50 324668 mE 6570076 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Hilltop
Rock Type: Calcrete, Ironstone



Vegetation: *Acacia collegialis* and *Eremophila oldfieldii* subsp. *angustifolia* tall sparse shrubland over *Dodonaea lobulata* mid sparse shrubland over *Senna artemisioides* subsp. *filifolia* low sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	350	5	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	140	0.1	
<i>Dodonaea lobulata</i>	140	5	
<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>	40	0.1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	350	5	
<i>Maireana pentatropis</i>	15	0.1	
<i>Maireana trichoptera</i>	15	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	25	1	
<i>Santalum acuminatum</i>	250	0.1	
<i>Scaevola spinescens</i>	130	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	15	2	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q19
Location MGA 50 328122 mE 6571915 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Plain No photo available
Rock Type: Laterite

Vegetation: *Eucalyptus salmonophloia* low woodland over *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Atriplex vesicaria* and *Atriplex* sp. low open chenopod shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex</i> sp.	35	5	
<i>Atriplex vesicaria</i>	100	20	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	120	5	
<i>Eucalyptus salmonophloia</i>	1000	15	
<i>Maireana georgei</i>	15	0.1	
<i>Salsola australis</i>	5	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q20
Location MGA 50 324559 mE 6569671 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Upper slope
Rock Type: Ironstone



Vegetation: *Eucalyptus salmonophloia* and *Eucalyptus salubris* mid open woodland over *Eremophila interstans* subsp. *interstans* and *Eremophila oldfieldii* subsp. *angustifolia* tall sparse shrubland over *Senna artemisioides* subsp. *filifolia* and *Atriplex nummularia* subsp. *spathulata* mid sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	25	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	120	0.5	
<i>Dodonaea lobulata</i>	60	0.1	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	90	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	350	3	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	1	
<i>Eucalyptus salmonophloia</i>	1200	3	
<i>Eucalyptus salubris</i>	1100	2	
<i>Maireana ?georgei</i>	15	0.1	
<i>Maireana pentatropis</i>	25	0.1	
<i>Maireana sedifolia</i>	100	0.5	
<i>Maireana trichoptera</i>	15	0.1	
<i>Olearia muelleri</i>	30	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	0.1	
<i>Scaevola spinescens</i>	30	0.1	
<i>Sclerolaena fusiformis</i>	15	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	130	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q21
Location MGA 50 327523 mE 6573455 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* and *Eremophila interstans* subsp. *interstans* tall open shrubland over *Senna artemisioides* subsp. *filifolia* and *Dodonaea stenozyga* mid sparse shrubland

Condition: Good **Disturbance Type:** Vehicle tracks, Litter

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Dodonaea stenozyga</i>	120	2	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	220	5	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	290	10	
<i>Eucalyptus griffithsii</i>	500	1	
<i>Maireana</i> ? <i>marginata</i>	30	2	
<i>Maireana georgei</i>	15	1	
<i>Olearia muelleri</i>	25	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	10	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	120	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q22
Location MGA 50 324398 mE 6569324 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Ironstone



Vegetation: *Acacia collegialis* and *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Ptilotus obovatus* var. *obovatus* and **Salvia verbenaca* low herbland

Condition: Good **Disturbance Type:** Grazing, Litter, Fauna tracks/scats, Historical Clearing

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	400	8	
<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>	40	0.1	
<i>Enneapogon caerulescens</i>	4	0.1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	350	4	
<i>Hibiscus sturtii</i> var. <i>grandiflorus</i>	15	0.1	
<i>Maireana trichoptera</i>	20	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	35	
<i>*Salvia verbenaca</i>	30	0.5	
<i>Sclerolaena fusiformis</i>	15	0.1	
<i>Sida petrophila</i>	30	4	
<i>Solanum cleistogamum</i>	10	0.1	
<i>Solanum lasiophyllum</i>	25	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q23
Location MGA 50 327250 mE 6573660 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Mud slope scree
Rock Type: Laterite



Vegetation: *Eucalyptus torquata* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* and *Acacia tetragonophylla* tall open shrubland over *Dodonaea stenozyga*, *Senna artemisioides* subsp. *filifolia* and *Alyxia buxifolia* mid shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia tetragonophylla</i>	250	1	
<i>Alyxia buxifolia</i>	130	1	
<i>Dodonaea stenozyga</i>	190	45	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	15	
<i>Eucalyptus torquata</i>	900	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	170	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q24
Location MGA 50 324535 mE 6569030 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Calcrete, Ironstone



Vegetation: *Eucalyptus salmonophloia* mid open woodland over *Atriplex nummularia* subsp. *spathulata* low sparse chenopod shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	15	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	90	5	
<i>Eucalyptus salmonophloia</i>	1200	6	
<i>Maireana pentatropis</i>	20	0.1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	50	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q25
Location MGA 50 326779 mE 6572716 mN

Described by: JW
Date: 16/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, Limestone, Quartz



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila glabra* subsp. *glabra*, *Atriplex vesicaria* and *Maireana georgei* low shrubland over *Osteocarpum salsuginosum* low sparse herbland

Condition: Poor, Very Poor **Disturbance Type:** Weeds, Vehicle tracks, Litter, Historical Clearing, Infrastructure

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?holocarpa</i>	35	5	
<i>Atriplex vesicaria</i>	38	15	
<i>Enneapogon caerulescens</i>	5	0.1	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	35	30	
<i>Eucalyptus griffithsii</i>	400	2	
<i>Maireana georgei</i>	20	15	
<i>Maireana pentatropis</i>	25	1	
<i>Maireana tomentosa</i>	20	5	
<i>Osteocarpum salsuginosum</i>	15	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q26
Location MGA 50 325151 mE 6568826 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Undulating plain
Rock Type: Ironstone



Vegetation: *Eucalyptus salmonophloia* and *Eucalyptus salubris* mid woodland over *Senna artemisioides* subsp. *filifolia* mid isolated shrubs over *Atriplex ?vesicaria* low isolated chenopod shrubs

Condition: Very Good **Disturbance Type:** Historical Clearing

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	40	0.5	
<i>Dodonaea lobulata</i>	50	0.1	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	40	0.1	
<i>Eucalyptus salmonophloia</i>	1200	15	
<i>Eucalyptus salubris</i>	1100	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	120	0.5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q27
Location MGA 50 326933 mE 6578601 mN

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, Quartz



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* mid open shrubland over *Dodonaea stenozyga*, *Eremophila glabra* subsp. *glabra* and *Santalum spicatum* mid open shrubland over *Atriplex vesicaria*, *Exocarpos aphyllus* and *Pittosporum angustifolium* low sparse shrubland

Condition: Good **Disturbance Type:** Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	42	0.5	
<i>Atriplex vesicaria</i>	60	5	
<i>Dodonaea stenozyga</i>	200	20	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	110	1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	20	
<i>Eucalyptus griffithsii</i>	500	1	
<i>Exocarpos aphyllus</i>	100	1	
<i>Lycium australe</i>	40	0.5	
<i>Maireana georgei</i>	20	0.5	
<i>Pittosporum angustifolium</i>	60	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	2	
<i>Santalum spicatum</i>	170	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	40	0.5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q28
Location MGA 50 325835 mE 6569465 mN

Described by: BD,SW
Date: 16/11/2021
Type: QUADRAT

Landform: Drainage line
Rock Type: Ironstone



Vegetation: *Eucalyptus salubris* mid open forest over *Eremophila interstans* subsp. *interstans* tall sparse shrubland over *Atriplex nummularia* subsp. *spathulata* mid sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	100	2	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	120	2	
<i>Dodonaea lobulata</i>	200	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	300	4	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	50	0.1	
<i>Eucalyptus salubris</i>	1100	35	
<i>Maireana trichoptera</i>	50	0.1	
<i>Olearia muelleri</i>	90	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	40	0.1	
<i>Scaevola spinescens</i>	80	0.1	
<i>Sclerolaena diacantha</i>	10	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q29
Location MGA 50 326944 **mE** 6579386 **mN**

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Plain No photo available
Rock Type: Dolerite, Laterite, Quartz

Vegetation: *Eucalyptus griffithsii* low open woodland over *Acacia hemiteles*, *Dodonaea stenozyga* and *Santalum spicatum* mid shrubland over *Atriplex vesicaria*, *Senna artemisioides* subsp. *filifolia* and *Olearia muelleri* low open shrubland
Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia hemiteles</i>	140	15	
<i>Alyxia buxifolia</i>	35	1	
<i>Atriplex vesicaria</i>	70	5	
<i>Dodonaea stenozyga</i>	120	15	
<i>Eucalyptus griffithsii</i>	700	10	
<i>Olearia muelleri</i>	15	2	
<i>Santalum spicatum</i>	200	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	45	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q31
Location MGA 50 326956 mE 6579776 mN

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Plain No photo available
Rock Type: Laterite

Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila interstans* subsp. *interstans* tall shrubland over *Eremophila oppositifolia* subsp. *angustifolia*, *Exocarpos aphyllus* and *Eremophila glabra* subsp. *glabra* mid sparse shrubland
Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Abutilon cryptopetalum</i>	38	0.1	
<i>Acacia erinacea</i>	40	0.5	
<i>Acacia jennerae</i>	50	5	
<i>Alyxia buxifolia</i>	40	15	
<i>Dodonaea stenozyga</i>	35	1	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	110	0.5	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	220	45	
<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i>	200	1	
<i>Eremophila scoparia</i>	48	0.1	
<i>Eucalyptus griffithsii</i>	500	5	
<i>Exocarpos aphyllus</i>	110	1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Olearia muelleri</i>	20	0.5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	35	0.5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q32
Location MGA 50 325641 mE 6569459 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite, Quartz



Vegetation: *Eucalyptus websteriana* subsp. *websteriana* low open woodland over *Acacia collegialis* tall open shrubland over *Dodonaea lobulata* low open shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	450	11	
<i>Austrostipa trichophylla</i>	5	0.1	
<i>Cheilanthes lasiophylla</i>	5	0.1	
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	8	0.1	
<i>Chrysocephalum puteale</i>	10	0.1	
<i>Dodonaea lobulata</i>	100	13	
<i>Enneapogon caerulescens</i>	3	0.1	
<i>Eremophila alternifolia</i>	30	0.1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	20	0.1	
<i>Eucalyptus websteriana</i> subsp. <i>websteriana</i>	300	1	
* <i>Euphorbia drummondii</i>	5	0.1	
<i>Goodenia havilandii</i>	10	0.1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Olearia muelleri</i>	35	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	1	
<i>Scaevola spinescens</i>	50	0.1	
<i>Sida</i> ? <i>petrophila</i>	10	0.1	
<i>Solanum lasiophyllum</i>	10	0.1	
<i>Waitzia acuminata</i> var. <i>acuminata</i>	10	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q33
Location MGA 50 324395 mE 6579552 mN

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Ridge
Rock Type: Laterite



Vegetation: *Acacia collegialis* tall open shrubland over *Eremophila oldfieldii* subsp. *angustifolia* mid sparse shrubland over *Dodonaea stenozyga* and *Solanum lasiophyllum* low open shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	400	21	
<i>Aristida contorta</i>	5	0.1	
<i>Dodonaea stenozyga</i>	50	20	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	120	10	
<i>Goodenia havilandii</i>	15	0.1	
<i>Solanum lasiophyllum</i>	15	0.5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q34
Location MGA 50 325544 mE 6569645 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite



Vegetation: *Eucalyptus oleosa* subsp. *oleosa* low isolated trees over *Acacia acuminata* and *Acacia collegialis* tall open shrubland over *Dodonaea lobulata* and *Eremophila georgei* mid open shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	450	10	
<i>Acacia collegialis</i>	450	3	
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	5	0.1	
<i>Dodonaea lobulata</i>	120	11	
<i>Eremophila georgei</i>	120	1	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	500	0.5	
<i>Leichhardtia australis</i>	300	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	130	0.1	
<i>Solanum lasiophyllum</i>	3	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q35
Location MGA 50 324374 mE 6579775 mN

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Mld slope
Rock Type: Laterite, Quartz



Vegetation: *Acacia acuminata* tall open shrubland over *Maireana triptera*, *Senna artemisioides* subsp. *filifolia* and *Dodonaea stenozyga* low open shrubland over *Ptilotus obovatus* var. *obovatus* and *Ptilotus polystachyus* low sparse herbland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	400	28	
<i>Cheilanthes lasiophylla</i>	5	0.1	
<i>Dodonaea stenozyga</i>	45	3	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	45	2	
<i>Maireana triptera</i>	25	5	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	1	
<i>Ptilotus polystachyus</i>	35	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	46	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q36
Location MGA 50 324947 mE 6578526 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Upper slope
Rock Type: Granite, Laterite



Vegetation: *Eucalyptus clelandiorum* and *Eucalyptus salmonophloia* mid open woodland over *Eremophila interstans* subsp. *interstans* tall sparse shrubland over *Eremophila parvifolia* subsp. *auricampi* and *Atriplex nummularia* subsp. *spathulata* low sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	40	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	80	0.5	
<i>Cratystylis conocephala</i>	40	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	250	3	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	100	1	
<i>Eucalyptus clelandiorum</i>	1100	3	
<i>Eucalyptus salmonophloia</i>	1100	2	
<i>Exocarpos aphyllus</i>	100	0.1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Roepera ovata</i>	5	0.1	
<i>Scaevola spinescens</i>	50	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	160	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q37
Location MGA 50 323658 mE 6579927 mN

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, Quartz



Vegetation: *Eucalyptus salmonophloia* and *Eucalyptus oleosa* subsp. *oleosa* mid woodland over *Eremophila interstans* subsp. *interstans* and *Atriplex ?vesicaria* mid shrubland over *Senna artemisioides* subsp. *filifolia*, *Chenopodium curvispicatum* and *Olearia muelleri* low open shrubland
Condition: Good **Disturbance Type:** Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	140	15	
<i>Chenopodium curvispicatum</i>	45	2	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	200	25	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	1300	2	
<i>Eucalyptus salmonophloia</i>	1500	15	
<i>Maireana pentatropis</i>	15	0.1	
<i>Maireana sedifolia</i>	45	1	
<i>Olearia muelleri</i>	35	2	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	35	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q38
Location MGA 50 324822 mE 6578130 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite



Vegetation: *Acacia collegialis* and *Eremophila oldfieldii* subsp. *angustifolia* tall open shrubland over *Dodonaea lobulata* and *Eremophila georgei* mid sparse shrubland over *Ptilotus obovatus* var. *obovatus* low isolated herbs

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	450	20	
<i>Aristida contorta</i>	5	0.1	
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	5	0.1	
<i>Dodonaea lobulata</i>	100	7	
<i>Eremophila georgei</i>	130	0.5	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	230	2	
<i>Goodenia havilandii</i>	15	0.1	
<i>Leichhardtia australis</i>	30	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	0.5	
<i>Santalum acuminatum</i>	300	0.1	
<i>Scaevola spinescens</i>	45	0.1	
<i>Sida ?petrophila</i>	10	0.5	
<i>Solanum lasiophyllum</i>	10	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q39
Location MGA 50 323564 mE 6579153 mN

Described by: JW
Date: 17/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Ironstone, Laterite, Quartz



Vegetation: *Eucalyptus celastroides* low woodland over *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Maireana pentatropis*, *Atriplex vesicaria* and *Chenopodium curvispicatum* low sparse chenopod shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	100	2	
<i>Chenopodium curvispicatum</i>	35	1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	120	10	
<i>Eucalyptus celastroides</i>	600	20	
<i>Maireana pentatropis</i>	25	5	
<i>Olearia muelleri</i>	35	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q40
Location MGA 50 324436 mE 6578315 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Laterite, Quartz



Vegetation: *Acacia collegialis* tall open shrubland over *Dodonaea lobulata* and *Eremophila georgei* mid open shrubland over *Ptilotus obovatus* var. *obovatus* low isolated herbs

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	450	15	
<i>Acacia tetragonophylla</i>	210	0.1	
<i>Aristida contorta</i>	10	0.1	
<i>Cheilanthes lasiophylla</i>	5	0.1	
<i>Cheilanthes sieberi</i> subsp. <i>sieberi</i>	8	0.1	
<i>Dodonaea lobulata</i>	170	10	
<i>Eremophila georgei</i>	140	3	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	50	0.1	
<i>Goodenia havilandii</i>	15	0.1	
<i>Leichhardtia australis</i>	10	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	0.5	
<i>Santalum acuminatum</i>	270	0.1	
<i>Scaevola spinescens</i>	120	0.1	
<i>Solanum lasiophyllum</i>	5	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q41
Location MGA 50 323554 mE 6579206 mN

Described by: JW
Date: 18/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Laterite, Quartz



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila interstans* subsp. *interstans*, *Eremophila glabra* subsp. *glabra* and *Acacia hemiteles* mid open shrubland over *Tecticornia halocnemoides*, *Atriplex* sp. and *Maireana trichoptera* low chenopod shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia hemiteles</i>	160	3	
<i>Atriplex</i> sp.	25	5	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	110	5	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	140	15	
<i>Eucalyptus griffithsii</i>	1000	5	
<i>Exocarpos aphyllus</i>	180	1	
<i>Frankenia ?fecunda</i>	5	0.1	
<i>Maireana trichoptera</i>	20	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	60	2	
<i>Tecticornia halocnemoides</i>	100	25	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q42
Location MGA 50 323685 mE 6577907 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Foothills
Rock Type: Ironstone, Quartz



Vegetation: *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Tecticornia halocnemoides* low open samphire shrubland

Condition: Good **Disturbance Type:** Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	50	2	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	90	1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	180	2	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	30	0.1	
<i>Exocarpos aphyllus</i>	30	0.1	
<i>Frankenia ?fecunda</i>	20	0.1	
<i>Lawrenzia repens</i>	3	0.1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Ptilotus exaltatus</i>	2	0.1	
<i>Rhagodia drummondii</i>	15	0.1	
<i>Roepera ovata</i>	8	0.1	
<i>Sclerolaena fusiformis</i>	5	0.1	
<i>Tecticornia halocnemoides</i>	90	11	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q43
Location MGA 50 323723 mE 6578705 mN

Described by: JW
Date: 18/11/2021
Type: QUADRAT

Landform: Ridge crwat
Rock Type: Ironstone, Laterite, Quartz



Vegetation: *Eucalyptus celastroides* low woodland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eucalyptus celastroides</i>	700	25	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R44
Location MGA 50 323751 mE 6577898 mN

Described by: BD,SW
Date: 17/11/2021
Type: RELEVÉ

Landform: Cracking clay No photo available
Rock Type: Ironstone, Quartz

Vegetation: *Streptoglossa liatroides* and *Sclerolaena cuneata* low sparse herbland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Goodenia pinnatifida</i>	5	0.1	
<i>Ptilotus exaltatus</i>	25	0.1	
<i>Sclerolaena cuneata</i>	5	0.5	
<i>Streptoglossa liatroides</i>	2	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q45
Location MGA 50 323830 mE 6584592 mN

Described by: JW
Date: 18/11/2021
Type: QUADRAT

Landform: Plain No photo available
Rock Type: Laterite, Quartz

Vegetation: *Eremophila oldfieldii* subsp. *angustifolia* tall sparse shrubland over *Acacia acuminata* and *Senna artemisioides* subsp. *filifolia* mid sparse shrubland over *Atriplex vesicaria*, *Maireana pentatropis* and *Maireana georgei* low sparse chenopod shrubland
Condition: Poor **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	170	5	
<i>Atriplex vesicaria</i>	45	5	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	220	5	
<i>Maireana georgei</i>	15	0.5	
<i>Maireana pentatropis</i>	15	2	
<i>Ptilotus exaltatus</i>	25	5	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	1	
<i>Rhodanthe chlorocephala</i> subsp. <i>rosea</i>	5	0.1	
* <i>Salvia verbenaca</i>	15	0.5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	140	1	
<i>Solanum lasiophyllum</i>	20	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q46
Location MGA 50 324398 mE 6577581 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Ironstone, Quartz



Vegetation: *Eucalyptus salmonophloia* mid open woodland over *Eremophila interstans* subsp. *interstans* tall sparse shrubland over *Atriplex nummularia* subsp. *spathulata* mid sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	50	2	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	160	6	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	250	1	
<i>Eucalyptus salmonophloia</i>	1300	7	
<i>Maireana trichoptera</i>	15	0.1	
<i>Scaevola spinescens</i>	30	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q48
Location MGA 50 324472 mE 6578839 mN

Described by: BD,SW
Date: 17/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Granite, Ironstone, Quartz



Vegetation: *Eucalyptus salubris* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* and *Acacia collegialis* tall sparse shrubland over *Dodonaea lobulata* and *Senna artemisioides* subsp. *filifolia* mid sparse shrubland

Condition: Good **Disturbance Type:** Litter, Historical Clearing

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia collegialis</i>	300	2	
<i>Aristida contorta</i>	5	0.1	
<i>Dodonaea lobulata</i>	130	4	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	4	
<i>Eucalyptus salubris</i>	600	1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Olearia muelleri</i>	40	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	0.1	
<i>Scaevola spinescens</i>	50	1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	150	3	
<i>Trymalium myrtillus</i> subsp. <i>myrtillus</i>	150	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q50
Location MGA 50 325838 mE 6579783 mN

Described by: BD,SW
Date: 18/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Ironstone, Quartz



Vegetation: *Acacia hemiteles*, *Casuarina pauper* and *Eremophila oldfieldii* subsp. *angustifolia* tall sparse shrubland over *Senna artemisioides* subsp. *filifolia*, *Acacia jennerae* and *Eremophila deserti* mid sparse shrubland over *Scaevola spinescens*, *Chenopodium curvispicatum* and *Maireana trichoptera* low sparse shrubland

Condition: Good **Disturbance Type:** Vehicle tracks, Fauna tracks/scats, Historical Clearing

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia hemiteles</i>	230	4	
<i>Acacia jennerae</i>	130	2	
<i>Aristida contorta</i>	10	0.1	
<i>Atriplex ?vesicaria</i>	40	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	330	0.1	
<i>Austrostipa platychaeta</i>	150	0.1	
* <i>Carrichtera annua</i>	15	0.1	
<i>Casuarina pauper</i>	300	1	
<i>Chenopodium curvispicatum</i>	20	0.5	
<i>Eremophila deserti</i>	170	1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	220	1	
<i>Leichhardtia australis</i>	100	0.1	
<i>Maireana sedifolia</i>	160	2	
<i>Maireana tomentosa</i>	5	0.1	
<i>Maireana trichoptera</i>	30	0.5	
<i>Pimelea microcephala</i> subsp. <i>microcephala</i>	190	1	
<i>Pittosporum angustifolium</i>	200	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	0.1	
<i>Salsola australis</i>	4	0.1	
* <i>Salvia verbenaca</i>	25	0.1	
<i>Scaevola spinescens</i>	100	3	
<i>Sclerolaena fusiformis</i>	10	0.5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	170	3	
<i>Solanum nummularium</i>	35	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q52
Location MGA 50 325464 mE 6579380 mN

Described by: BD,SW
Date: 18/11/2021
Type: QUADRAT

Landform: Claypan
Rock Type: Calcrete, Ironstone, Quartz



Vegetation: *Dodonaea lobulata* and *Eremophila interstans* subsp. *interstans* mid sparse shrubland over *Maireana trichoptera*, *Atriplex vesicaria* and *Maireana georgei* low sparse chenopod shrubland over **Salvia verbenaca*, *Ptilotus exaltatus* and *Sclerolaena fusiformis* low herbland

Condition: Good **Disturbance Type:** Weeds, Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	50	1	
<i>Calandrinia baccata</i>	5	0.1	
<i>*Carrichtera annua</i>	15	1	
<i>*Centaurea melitensis</i>	25	0.1	
<i>Chenopodium curvispicatum</i>	30	0.1	
<i>Dodonaea lobulata</i>	150	4	
<i>Dysphania melanocarpa</i> forma <i>leucocarpa</i>	10	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	180	1	
<i>Lawrenzia repens</i>	5	1	
<i>Maireana georgei</i>	20	1	
<i>Maireana trichoptera</i>	40	3	
<i>Monachather paradoxus</i>	5	0.1	
<i>*Oligocarpus calendulaceus</i>	5	0.1	
<i>Ptilotus exaltatus</i>	35	12	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	20	0.1	
<i>*Salvia verbenaca</i>	20	15	
<i>Sclerolaena fusiformis</i>	8	5	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	150	0.1	
<i>Solanum lasiophyllum</i>	10	0.1	
<i>Streptoglossa liatroides</i>	5	3	
<i>Vittadinia dissecta</i> var. <i>hirta</i>	30	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2Q54
Location MGA 50 324779 mE 6583442 mN

Described by: BD,SW
Date: 18/11/2021
Type: QUADRAT

Landform: Undulating plain
Rock Type: Ironstone, Laterite



Vegetation: *Eucalyptus salmonophloia* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* tall isolated shrubs over *Senna artemisioides* subsp. *filifolia*, *Eremophila interstans* subsp. *interstans* and *Dodonaea lobulata* mid open shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia tetragonophylla</i>	170	0.1	
<i>Aristida contorta</i>	15	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	120	0.1	
<i>Dodonaea lobulata</i>	100	2	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	160	1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	160	3	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	0.5	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	130	1	
<i>Eucalyptus salmonophloia</i>	900	5	
<i>Leichhardtia australis</i>	150	0.1	
<i>Maireana trichoptera</i>	15	0.1	
<i>Olearia muelleri</i>	30	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	30	0.1	
<i>Roepera ovata</i>	5	0.1	
<i>Scaevola spinescens</i>	50	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	140	8	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R05
Location MGA 50 327540 mE 6571802 mN

Described by: JW
Date: 16/11/2021
Type: RELEVE

Landform: Minor creek line No photo available
Rock Type: Laterite

Vegetation: *Eucalyptus torquata* low woodland over *Dodonaea stenozyga* mid sparse shrubland over *Ptilotus obovatus* var. *obovatus* and *Haloragis trigonocarpa* low sparse herbland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Dodonaea stenozyga</i>	110	5	
<i>Eucalyptus torquata</i>	400	20	
<i>Haloragis trigonocarpa</i>	40	2	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	25	6	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	48	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R06
Location MGA 50 327365 mE 6572989 mN

Described by: JW
Date: 16/11/2021
Type: RELEVÉ

Landform: Lake
Rock Type: N/A



Vegetation: *Schinus molle var. areira low woodland over Aristida contorta low grassland over *Heliotropium europaeum, Heliotropium curassavicum and *Sonchus asper low herbland

Condition: Degraded **Disturbance Type:** Weeds, Grazing, Vehicle tracks, Litter, Fauna tracks/scats

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
Aristida contorta	30	70	
Heliotropium curassavicum	18	1	
*Heliotropium europaeum	2	40	
*Schinus molle var. areira	300	15	
*Sonchus asper	70	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R07
Location MGA 50 327376 mE 6570901 mN

Described by: JW
Date: 16/11/2021
Type: RELEVE

Landform: Mid slope
Rock Type: Laterite



Vegetation: *Eucalyptus oleosa* subsp. *oleosa* low woodland over *Atriplex vesicaria* low sparse chenopod shrubland over *Heliotropium curassavicum* low sparse herbland

Condition: Good **Disturbance Type:** Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex vesicaria</i>	60	10	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	700	15	
<i>Heliotropium curassavicum</i>	40	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R09
Location MGA 50 327166 mE 6574087 mN

Described by: JW
Date: 16/11/2021
Type: RELEVÉ

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus torquata* and *Eucalyptus griffithsii* low woodland over *Eremophila oldfieldii* subsp. *angustifolia* and *Santalum acuminatum* tall open shrubland over *Dodonaea stenozyga*, *Senna artemisioides* subsp. *filifolia* and *Alyxia buxifolia* low shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Alyxia buxifolia</i>	50	5	
<i>Dodonaea stenozyga</i>	80	25	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	300	25	
<i>Eucalyptus griffithsii</i>	800	2	
<i>Eucalyptus torquata</i>	1000	15	
<i>Santalum acuminatum</i>	220	2	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	90	18	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R11
Location MGA 50 325721 mE 6569452 mN

Described by: BD,SW
Date: 16/11/2021
Type: RELEVÉ

Landform: Drainage line
Rock Type: Ironstone



Vegetation: *Eucalyptus salmonophloia* mid woodland over *Eremophila interstans* subsp. *interstans* and *Acacia collegialis* tall open shrubland over *Senna artemisioides* subsp. *filifolia* and *Dodonaea lobulata* mid open shrubland
Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	250	0.1	
<i>Acacia collegialis</i>	300	1	
<i>Dodonaea lobulata</i>	120	2	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	250	20	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	250	0.1	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	40	0.1	
<i>Eucalyptus salmonophloia</i>	1200	20	
<i>Olearia muelleri</i>	40	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	40	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	150	15	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R13
Location MGA 50 324764 mE 6579229 mN

Described by: BD,SW
Date: 17/11/2021
Type: RELEVÉ

Landform: Plain No photo available
Rock Type: Calcrete, Laterite

Vegetation: *Maireana georgei* mid sparse chenopod shrubland over *Atriplex nummularia* subsp. *spathulata*, *Maireana tomentosa* and *Atriplex ?vesicaria* low sparse chenopod shrubland over **Centaurea melitensis* and *Ptilotus obovatus* var. *obovatus* low open herbland
Condition: Very Poor **Disturbance Type:** Weeds, Vehicle tracks, Litter, Historical Clearing, Infrastructure

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	25	0.5	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	20	1	
<i>Atriplex quadrivalvata</i>	15	0.1	
* <i>Centaurea melitensis</i>	40	10	
<i>Hibiscus solanifolius</i>	120	0.1	
<i>Maireana georgei</i>	120	2	
<i>Maireana tomentosa</i>	30	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	35	10	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R15
Location MGA 50 324420 mE 6579488 mN

Described by: JW
Date: 17/11/2021
Type: RELEVÉ

Landform: Ridge crst
Rock Type: Laterite, Quartz



Vegetation: *Eucalyptus campaspe* and *Eucalyptus oleosa* subsp. *oleosa* low open forest over *Senna artemisioides* subsp. *filifolia* mid open shrubland over *Olearia muelleri* and *Eremophila glabra* subsp. *glabra* low open shrubland
Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila glabra</i> subsp. <i>glabra</i>	90	1	
<i>Eucalyptus campaspe</i>	1000	25	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	1000	15	
<i>Olearia muelleri</i>	28	10	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	200	18	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C2R17
Location MGA 50 323480 mE 6579368 mN

Described by: JW
Date: 17/11/2021
Type: RELEVE

Landform: Plain
Rock Type: Ironstone, Laterite



Vegetation: *Eucalyptus oleosa* subsp. *oleosa* and *Eucalyptus celastroides* low woodland over *Senna artemisioides* subsp. *filifolia* and *Eremophila interstans* subsp. *interstans* low sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Eremophila interstans</i> subsp. <i>interstans</i>	35	0.5	
<i>Eucalyptus celastroides</i>	500	1	
<i>Eucalyptus oleosa</i> subsp. <i>oleosa</i>	700	10	
<i>Olearia muelleri</i>	25	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	60	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3Q01
Location MGA 50 328133 mE 6574120 mN

Described by: JW
Date: 18/11/2021
Type: QUADRAT

Landform: Plai
Rock Type: Laterite



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila interstans* subsp. *interstans*, *Senna artemisioides* subsp. *filifolia* and *Acacia jennerae* mid open shrubland over *Alyxia buxifolia*, *Westringia rigida* and *Olearia muelleri* low sparse shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	120	1	
<i>Acacia jennerae</i>	140	1	
<i>Alyxia buxifolia</i>	90	2	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	130	5	
<i>Eucalyptus griffithsii</i>	700	5	
<i>Exocarpos aphyllus</i>	150	1	
<i>Maireana tomentosa</i>	10	0.1	
<i>Olearia muelleri</i>	28	0.5	
<i>Ptilotus exaltatus</i>	15	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	120	5	
<i>Westringia rigida</i>	45	2	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3Q02
Location MGA 50 328109 mE 6573080 mN

Described by: BD,SW
Date: 18/11/2021
Type: QUADRAT

Landform: Upper slope
Rock Type: Granite, Laterite



Vegetation: *Eucalyptus torquata* low open forest over *Dodonaea stenozyga* low sparse shrubland

Condition: Good **Disturbance Type:** Litter

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>?Enchylaena tomentosa</i>	20	0.1	
<i>Atriplex ?vesicaria</i>	40	0.1	
<i>Dodonaea lobulata</i>	90	0.1	
<i>Dodonaea stenozyga</i>	80	3	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	180	0.1	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	60	0.1	
<i>Eucalyptus torquata</i>	800	35	
<i>Exocarpos aphyllus</i>	40	0.1	
<i>Maireana trichoptera</i>	10	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	10	0.1	
<i>Rytidosperma</i> sp.	10	0.1	
<i>Scaevola spinescens</i>	50	0.1	
<i>Solanum nummularium</i>	5	0.1	
<i>Trymalium myrtillus</i> subsp. <i>myrtillus</i>	20	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3Q02A
Location MGA 50 327861 mE 6573908 mN

Described by: JW
Date: 18/11/2021
Type: QUADRAT

Landform: Hill top
Rock Type: Granite, Ironstone, Laterite



Vegetation: *Eucalyptus torquata* low open woodland over *Dodonaea stenozyga*, *Acacia acuminata* and *Eremophila oldfieldii* subsp. *angustifolia* mid open shrubland over *Alyxia buxifolia* and *Senna artemisioides* subsp. *filifolia* low sparse shrubland
Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia acuminata</i>	200	3	
<i>Acacia dissona</i> var. <i>dissona</i>	200	1	
<i>Alyxia buxifolia</i>	50	1	
<i>Dodonaea stenozyga</i>	110	12	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	150	3	
<i>Eucalyptus torquata</i>	1000	5	
<i>Maireana georgei</i>	5	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	70	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3Q02B
Location MGA 50 327412 mE 6574092 mN

Described by: JW
Date: 18/11/2021
Type: QUADRAT

Landform: Plain
Rock Type: Ironstone, Laterite



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila oldfieldii* subsp. *angustifolia* mid sparse shrubland over *Atriplex vesicaria*, *Dodonaea stenozyga* and *Acacia dissona* var. *dissona* low open shrubland

Condition: Very Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Acacia dissona</i> var. <i>dissona</i>	25	1	
<i>Acacia hemiteles</i>	100	1	
<i>Atriplex vesicaria</i>	45	15	
<i>Dodonaea stenozyga</i>	25	5	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	25	1	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	150	5	
<i>Eucalyptus griffithsii</i>	700	5	
<i>Maireana pentatropis</i>	15	0.1	
<i>Maireana tomentosa</i>	10	0.5	
<i>Olearia muelleri</i>	25	1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	15	1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3Q06
Location MGA 50 327864 mE 6573343 mN

Described by: BD,SW
Date: 18/11/2021
Type: QUADRAT

Landform: Mid slope
Rock Type: Calcrete, Laterite



Vegetation: *Eucalyptus torquata* low woodland over *Eremophila interstans* subsp. *interstans* tall sparse shrubland over *Dodonaea stenozyga* mid sparse shrubland

Condition: Good **Disturbance Type:** Vehicle tracks, Fauna tracks/scats

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Atriplex ?vesicaria</i>	50	0.1	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	40	0.1	
<i>Austrostipa platychaeta</i>	80	0.1	
<i>Dodonaea stenozyga</i>	150	1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	370	1	
<i>Eremophila parvifolia</i> subsp. <i>auricampi</i>	30	0.1	
<i>Eucalyptus torquata</i>	800	15	
<i>Maireana sedifolia</i>	120	0.1	
<i>Maireana tomentosa</i>	5	0.1	
<i>Maireana trichoptera</i>	15	0.1	
<i>Olearia muelleri</i>	40	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	60	0.1	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3R01
Location MGA 50 328239 mE 6574703 mN

Described by: JW
Date: 18/11/2021
Type: RELEVÉ

Landform: Plain
Rock Type: Laterite



Vegetation: *Eucalyptus griffithsii* low open woodland over *Eremophila interstans* subsp. *interstans* tall open shrubland over *Atriplex vesicaria* and *Exocarpos aphyllus* mid open shrubland

Condition: Good **Disturbance Type:** None

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
<i>Alyxia buxifolia</i>	90	1	
<i>Atriplex vesicaria</i>	110	15	
<i>Dianella revoluta</i> var. <i>divaricata</i>	60	0.5	
<i>Dodonaea stenozyga</i>	15	0.5	
<i>Eremophila glabra</i> subsp. <i>glabra</i>	70	2	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	220	15	
<i>Eucalyptus griffithsii</i>	1000	5	
<i>Exocarpos aphyllus</i>	130	5	
<i>Maireana pentatropis</i>	15	0.5	
<i>Maireana trichoptera</i>	20	0.1	
<i>Olearia muelleri</i>	15	0.5	
<i>Pimelea spiculigera</i> var. <i>thesioides</i>	20	0.1	
<i>Ptilotus exaltatus</i>	35	5	

FLORA SITE SHEET

Project Name 4794 Coolgardie Biological Survey
Site: C3R04
Location MGA 50 328218 mE 6573560 mN

Described by: BD,SW
Date: 18/11/2021
Type: RELEVÉ

Landform: Creek line
Rock Type: Granite, Quartz



Vegetation: *Eucalyptus salmonophloia* mid open forest over *Eremophila interstans* subsp. *interstans* tall sparse shrubland over *Acacia jennerae* mid sparse shrubland

Condition: Good **Disturbance Type:** Weeds, Vehicle tracks

SPECIES LIST

Taxon	Height (cm)	Cover (%)	Notes
? <i>Pimelea</i> sp.	170	0.1	
<i>Acacia jennerae</i>	150	5	
* <i>Asphodelus fistulosus</i>	30	0.1	
<i>Atriplex</i> ? <i>vesicaria</i>	30	2	
<i>Atriplex nummularia</i> subsp. <i>spathulata</i>	50	1	
* <i>Carrichtera annua</i>	20	0.1	
* <i>Centaurea melitensis</i>	20	0.1	
<i>Chloris truncata</i>	30	0.1	
<i>Eragrostis dielsii</i>	3	0.1	
<i>Eremophila interstans</i> subsp. <i>interstans</i>	210	4	
<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i>	140	0.1	
<i>Eucalyptus salmonophloia</i>	1100	40	
<i>Maireana trichoptera</i>	20	0.1	
<i>Olearia muelleri</i>	40	0.1	
* <i>Oligocarpus calendulaceus</i>	5	0.1	
<i>Pittosporum angustifolium</i>	210	0.1	
<i>Ptilotus exaltatus</i>	30	0.1	
<i>Ptilotus obovatus</i> var. <i>obovatus</i>	25	0.1	
<i>Sclerolaena diacantha</i>	5	0.1	
<i>Senna artemisioides</i> subsp. <i>filifolia</i>	120	0.1	
<i>Vittadinia dissecta</i> var. <i>hirta</i>	15	0.1	

Appendix F

Fauna Habitat Assessments

HABC01

Project:	4794 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	899984.8973525156	Northing	6572010.013459269
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open forest (50-80%)	<i>Acacia sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>dodonea sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>ptilotus sp.</i>



Fulcrum photo ID 42151a58-0510-40c7-b5b8-39f62d4c1f75,4ffc3568-888a-4c64-9f70-

HABC02

Project:	4795 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	899822.8621105128	Northing	6571973.766455285
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Disturbance	None observed		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Acacia sp.</i>
Mid stratum	Mid (1-2 m)	Open mallee shrubland (20-50%)	
Ground stratum	Low (>0.5 m)	Open forbland (20-50%)	<i>ptilotus sp.</i>



Fulcrum photo ID 687a6808-3ad9-4430-9631-687bd7c191ca,3a9bdcdb2-3829-4fc9-b373-

HABC03

Project:	4796 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	899951.8547312932	Northing	6571730.697303254
Landform and soil		Rock	
Landform	Plain	Rock type/s	Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Good		
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Hollows - trees,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eucalyptus mallee</i>
Ground stratum	Low (>0.5 m)	Open forbland (20-50%)	<i>eremophila sp, salt Bush, acacia sp.</i>



Fulcrum photo ID cae7ef5d-35d9-43c9-bff0-76d8079f5710,e1c8ac9f-dd84-4e1b-a0b8-

HABC04

Project:	4797 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	899917.5832372475	Northing	6571665.113011075
Landform and soil		Rock	
Landform	Undulating plain	Rock type/s	Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	5 - 25%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Peeling bark,Rock crevices,Woody debris
Introduced fauna	Cattle		
Vegetation			
Upper stratum	Low (<10 m)	Isolated trees (<0.25%)	<i>Eremophila sp.</i>
Mid stratum	Mid (1-2 m)	Shrubland and/or heathland (50-80%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>ptilotus sp.</i>



Fulcrum photo ID 5e27f824-34b2-4807-bc1c-60a160cc8ad3,d582a71f-1bea-4caa-99e9-

HABC05

Project:	4798 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	900074.8952529589	Northing	6571739.672393914
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Litter,Vehicle tracks,Weeds	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Forbland (50-80%)	<i>saltbush sp.</i>



Fulcrum photo ID e29ce514-0173-433b-bb49-ef3ded5af200,1db3699d-1b8b-4cf9-950a-

HABC06

Project:	4799 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	900594.4049126134	Northing	6571695.699349251
Landform and soil		Rock	
Landform	Gorge	Rock type/s	Granite
Soil type	Clay	Surface stone cover	
Soil colour	Brown	Surface stone size classes present	0 - 5%
Condition			
Quality	Highly degraded	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Present
Disturbance	Clearing,Litter,Vehicle tracks	Microhabitats	Exfoliating rock
Introduced fauna	Goat		
Vegetation			
Upper stratum	Absent		
Mid stratum	Absent		
Ground stratum	Low (>0.5 m)	Sparse tussock grassland (0.25-20%)	<i>eraharta sp.</i>



Fulcrum photo ID 7c25ab6a-4d43-4390-897a-cb707b353f57,7ba253fb-c9d3-4b40-885f-

HABC07

Project:	4800 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	901166.9975909423	Northing	6571815.560922004
Landform and soil		Rock	
Landform	Undulating plain	Rock type/s	Granite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Burrows,Leaf litter,Peeling bark,Woody debris
Introduced fauna	Cattle,Rabbit		
Vegetation			
Upper stratum	Low (<10 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp. and eremophila sp.</i>
Mid stratum	Low (0.5-1 m)	Shrubland and/or heathland (50-80%)	<i>dodonaea</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>saltbush and Ptilotus sp.</i>



Fulcrum photo ID 0103efbf-c7eb-4e3f-bb46-943ea4911830,ccb4b5f6-49b0-4d03-8017-

HABC08

Project:	4801 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	901301.0032407076	Northing	6571962.571508318
Landform and soil		Rock	
Landform	Undulating plain	Rock type/s	Granite
Soil type	Sandy loam	Surface stone cover	
Soil colour	Grey,Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	Cattle		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>saltbush sp.</i>



Fulcrum photo ID 212d5260-a83c-4619-b069-494f4f980745,0b6d6cef-7222-4fa8-9bcc-

HABC09

Project:	4802 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	901682.8518081398	Northing	6571863.583170791
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Sandy clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Very good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Litter	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Absent		



Fulcrum photo ID 117fc066-1e33-488f-b006-63d6e0f432e1,e1f0326e-d28b-470e-b3ef-

HABC10

Project:	4803 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	902022.5631623187	Northing	6572062.188234046
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Granite
Soil type	Clay loam	Surface stone cover	
Soil colour	Red	Surface stone size classes present	25 - 50%
Condition			
Quality	Very good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	None observed	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Rock crevices,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Shrubland and/or heathland (50-80%)	<i>acacia sp. eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	



Fulcrum photo ID b36a6fd3-4521-41ec-ba74-6fdeac4098bb,0044da98-5d6a-4f2e-9cba-

HABC11

Project:	4804 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	902400.9875870193	Northing	6571798.486756262
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Calcrete,Granite
Soil type	Sandy loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Disturbance	None observed		
Introduced fauna	Goat,Rabbit		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Shrubland and/or heathland (50-80%)	<i>eremophila sp. dodenea sp. acacia sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>eremophila</i>



Fulcrum photo ID 3c1c6583-ed7c-459a-93f5-8dd769a4d26c,4211c7c8-571a-4138-af6c-

HABC12

Project:	4805 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	902370.029827407	Northing	6571589.932107501
Landform and soil		Rock	
Landform	Plain	Rock type/s	Unknown
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Shrubland and/or heathland (50-80%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	



Fulcrum photo ID aef600dc-722f-4dd1-a355-599e69a6b123,e75ca763-a192-495f-a373-

HABC13

Project:	4806 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	900730.9056804918	Northing	6570131.034563001
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Red	Surface stone size classes present	50 - 75%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance		Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Introduced fauna	Goat,Rabbit		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	Acacia sp.
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	dodonaea sp.
Ground stratum	Low (>0.5 m)	Isolated tussock grasses (<0.25%)	grass species



Fulcrum photo ID 58005a12-2165-49c3-b705-24e2a50985e8,0b2d1291-65d4-48da-b5d6

HABC14

Project:	4807 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	900889.8561838119	Northing	6569940.604672145
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Calcrete,Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Grey	Surface stone size classes present	25 - 50%
Condition			
Quality	Disturbed	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp. salt bush</i>
Ground stratum	Mid (0.5-1 m)	Open forbland (20-50%)	<i>salt bush sp.</i>



Fulcrum photo ID 3b56a64c-51c9-4e00-bdc1-f54d4301f218,c0fddb0c-66e8-40b3-9c48-

HABC15

Project:	4808 Coolgardie Biological Spring Survey		
Date	2021-10-12	Personnel	LC
Easting	901542.5074801195	Northing	6568846.8061696505
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Calcrete,Granite
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Red	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Very good		
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	None observed	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Rock crevices,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>acacia sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>ptilotus sp</i>



Fulcrum photo ID bfb61e82-be29-4f8f-a414-761f945af8e4,e6e2629f-1f45-4bcd-9213-

HABC16

Project:	4809 Coolgardie Biological Spring Survey		
Date	2021-10-13	Personnel	LC
Easting	902295.8342707243	Northing	6571439.575077539
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Calcrete,Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Erosion	Microhabitats	Hollows - trees,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>Saltbush sp.</i>



Fulcrum photo ID b70c7eb9-827e-46df-b8e4-5e4e110d293a,85f73721-6a97-4980-94c7-

HABC17

Project:	4810 Coolgardie Biological Spring Survey		
Date	2021-10-13	Personnel	LC
Easting	901495.8347967644	Northing	6571719.769933704
Landform and soil		Rock	
Landform	Plain	Rock type/s	Calcrete,Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Orange,Yellow	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Burrows,Hollows - logs,Hollows - trees,Leaf litter,Peeling bark,Woody debris
Disturbance	Litter,Vehicle tracks		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>ptilotus sp. and herb sp.</i>



Fulcrum photo ID 372a2bcc-d839-41ff-9db2-d990a2431994,cd2c96ca-6b1d-49e3-b273-

HABC18

Project:	4811 Coolgardie Biological Spring Survey		
Date	2021-10-13	Personnel	LC
Easting	899905.7739034619	Northing	6571300.257774439
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange,Red	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Shrubland and/or heathland (50-80%)	<i>dodenea sp. eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>ptilotus sp.</i>



Fulcrum photo ID 3663874c-d46e-438f-8c20-f9744960ca60,30a5a95b-d86a-4c39-8c3b-

HABC19

Project:	4812 Coolgardie Biological Spring Survey		
Date	2021-10-13	Personnel	LC
Easting	899738.7645235118	Northing	6571320.002878793
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Absent		
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>saltbush sp. and Ptilotus sp.</i>



Fulcrum photo ID 3996f487-e2d9-4d38-9be8-65219fc9ea13,d5e4af84-5d54-464a-b161-

HABC20

Project:	4813 Coolgardie Biological Spring Survey		
Date	2021-10-14	Personnel	LC
Easting	900350.4372829349	Northing	6571342.937353729
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Calcrete,Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Grey	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter,Peeling bark,Woody debris
Disturbance	Erosion,Litter,Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>saltbush sp. and herb sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>saltbush sp.</i>



Fulcrum photo ID cfd77415-c024-4198-bd3a-e011a5fb5404,15466c44-d7ac-4902-a088-

HABC21

Project:	4814 Coolgardie Biological Spring Survey		
Date	2021-10-14	Personnel	LC
Easting	900421.5213021053	Northing	6571645.335509442
Landform and soil		Rock	
Landform	Claypan	Rock type/s	Granite
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Highly degraded	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Rock crevices, Woody debris
Disturbance	Clearing, Erosion, Litter, Overgrazing, Vehicle		
Introduced fauna	Cattle		
Vegetation			
Upper stratum	Mid (10-30 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp.</i>
Mid stratum	Absent		
Ground stratum	Low (>0.5 m)	Tussock grassland (50-80%)	<i>grass sp. and thistle sp. sedges boemia sp.</i>



Fulcrum photo ID bfe042ed-7948-4c78-9fe1-7c8541220392,7696b9b4-9eeb-4df8-8e63-

HABC22

Project:	4815 Coolgardie Biological Spring Survey		
Date	2021-10-14	Personnel	LC
Easting	900235.352356134	Northing	6572033.776216977
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Calcrete, Granite, Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange, Red	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Little or no fire evidence (>5 years)	Microhabitats	Leaf litter, Peeling bark, Rock crevices, Woody debris
Disturbance	Clearing, Litter, Vehicle tracks, Weeds		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Woodland (20-50%)	<i>Acacia sp.</i>
Mid stratum	Low (0.5-1 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>dodenea sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>ptilotus sp. herbs</i>



Fulcrum photo ID 51a9f868-dbf8-4d0d-aa65-cebf271c9d24, bcb04dd4-53d3-407a-9ce9-

HABC23

Project:	4816 Coolgardie Biological Spring Survey		
Date	2021-10-14	Personnel	LC
Easting	900167.7359049395	Northing	6570928.550377593
Landform and soil		Rock	
Landform	Plain	Rock type/s	Calcrete,Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Grey,Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Highly degraded		
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Clearing,Litter,Vehicle tracks	Microhabitats	Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Low (0.5-1 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>saltbush cena sp. herbs sp.</i>



Fulcrum photo ID 4e1b0f41-0592-423d-8408-a19672a9d5c6,f06cb18e-09e9-41d6-a61b-

HABC24

Project:	4817 Coolgardie Biological Spring Survey		
Date	2021-10-14	Personnel	LC
Easting	899663.2610756675	Northing	6571226.542719762
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Calcrete,Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition			
Quality	Very good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Peeling bark,Rock crevices,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila sp. dodenea sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>ptilotus sp. salt bush sp.</i>



Fulcrum photo ID 8f123c2e-9663-4712-845a-fe15deec853c,26ed00e2-46e6-4415-ad80-

C2H2

Project:	4818 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	SW
Easting	897743.7674614477	Northing	6565685.35287808
Landform and soil		Rock	
Landform	Plain	Rock type/s	Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition			
Quality	Good	Habitat Features	
Fire History	Little or no fire evidence (>5 years)	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Peeling bark
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Acacia and Melaleuca sp</i>
Ground stratum	Low (>0.5 m)	Isolated hummock grasses (<0.25%)	<i>Scaveloa spin</i>



Fulcrum photo ID c4cad0bd-4f57-4209-9459-7401a6a867d2

C2H4

Project:	4819 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	SW
Easting	897738.8708299723	Northing	6565557.982810425
Landform and soil		Rock	
Landform	Plain	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	75 - 100%
Condition			
Quality	Very good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Peeling bark
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>Melaleuca sp</i>
Ground stratum	Low (>0.5 m)	Open hummock grassland (20-50%)	<i>Scaveola spin</i>



Fulcrum photo ID a0f87fb2-1a73-43d5-bfb4-bf7e2ce804a4

C2H6

Project:	4820 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	SW
Easting	897718.2217606371	Northing	6565454.049011737
Landform and soil		Rock	
Landform	Plain	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	75 - 100%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Disturbance			
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eucalyptus sp</i>
Ground stratum	Low (>0.5 m)		



Fulcrum photo ID a25ac42a-b0be-4c30-a8b2-8e07a2f5bcd9

C2H8

Project:	4821 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	SW
Easting	897820.0346483674	Northing	6565227.984386012
Landform and soil		Rock	
Landform	Plain	Rock type/s	Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eremophila sp</i>
Ground stratum			



Fulcrum photo ID 93421722-1ad4-4c10-9e11-de2ef1902d8b

C2H10

Project:	4822 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	SW
Easting	898008.8963851782	Northing	6565265.629842697
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Calcrete,Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange,Red	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	Erosion	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>Eremophila</i>
Ground stratum			



Fulcrum photo ID 45de2c23-38f7-401c-8d48-42e59fe7b842

C2H12

Project:	4823 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	SW
Easting	898183.4047518701	Northing	6565240.394553069
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition			
Quality	Good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Erosion	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Low (0.5-1 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>Senna art fil</i>
Ground stratum			



Fulcrum photo ID 08469bcc-a34b-4b23-bf34-76982cd97e9c

HABC2LC01

Project:	4824 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	LC
Easting	899664.2211246493	Northing	6565980.506265143
Landform and soil		Rock	
Landform	Gorge	Rock type/s	Granite,Quartz
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Yellow	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Highly degraded	Water Source	Absent
Fire History	Unknown	Microhabitats	Exfoliating rock,Rock crevices
Disturbance			
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp. salt bush sp.</i>



Fulcrum photo ID 4ebf3251-d69c-4d7e-93c5-260eba7ae549,d2ea3ab1-5502-453c-9997-

HABC2LC02

Project:	4825 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	LC
Easting	899850.8139667105	Northing	6565706.821050039
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Granite,Quartz
Soil type	Sandy clay	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Disturbance			
Introduced fauna	Goat		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>eremophila sp.</i>



Fulcrum photo ID 4b2247e7-9fed-441d-b103-59e6bf65756c,a6bfbb91-71be-4e28-992d-

HABC2LC03

Project:	4826 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	LC
Easting	900665.7028282916	Northing	6565871.541426335
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Sandy clay	Surface stone cover	
Soil colour	Orange,Red	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown		
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Rock crevices,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open forest (50-80%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Mid (0.5-1 m)	Sparse forbland (0.25-20%)	<i>eremophila sp.</i>



Fulcrum photo ID b0752528-0a49-4733-a884-d5b48bd2d627,ce6b6969-95db-4c53-bfdd-

HABC2LC04

Project:	4827 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	LC
Easting	900563.0312287982	Northing	6565632.278082406
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Orange,Red	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Good		
Fire History	Unknown	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Rock crevices,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Woodland (20-50%)	<i>eremophila sp. Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>ptilotus sp.</i>



Fulcrum photo ID 2b2a12b3-b9f2-4fd2-9d5d-b72a773595d9,19860892-f6e1-4a85-843a-

HABC2LC05

Project:	4828 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	LC
Easting	900752.345712022	Northing	6565409.609161669
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Granite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition			
Quality	Good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Erosion,Vehicle tracks	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp.</i>



Fulcrum photo ID bc023295-6531-4172-95cc-e3d96c28e365,3b64dc10-30df-4627-8e7f-

HABC2LC06

Project:	4829 Coolgardie Biological Spring Survey		
Date	2021-11-15	Personnel	LC
Easting	900415.1933355466	Northing	6565440.556371264
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Hollows - logs,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open forest (50-80%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i> , <i>salt bush sp. (in photos)</i>
Ground stratum	Absent		



Fulcrum photo ID 30f77ad9-d6b9-45a3-885d-56cf525debd8,a70c538c-fc2a-4dbd-be76-

C2H14

Project:	4830 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	898138.1000480026	Northing	6564928.758600489
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Ironstone
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	None observed	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	Acacia sp
Ground stratum	Absent		



Fulcrum photo ID 6eb608e3-e562-4314-b5d8-0ecf176e3906

C2H16

Project:	4831 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	898035.707931469	Northing	6564690.336170336
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Ironstone
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	None observed	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eremophila</i>
Ground stratum	Absent		



Fulcrum photo ID 10efecf9-ff45-4474-8f07-5ef50a3af30e

C2H18

Project:	4832 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	897721.3419470013	Northing	6564081.199149741
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Calcrete, Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs, Leaf litter, Peeling bark, Woody debris
Disturbance			
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Eremophila</i>
Ground stratum	Absent		



Fulcrum photo ID 25ba9c30-9a14-438f-952a-4d1cff16da86

C2H20

Project:	4833 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	897584.545833359	Northing	6563668.195859093
Landform and soil		Rock	
Landform	Escarpment	Rock type/s	Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs, Leaf litter, Peeling bark, Woody debris
Disturbance	None observed		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eremophila</i>
Ground stratum			



Fulcrum photo ID 2cd10b5a-17ad-4733-9cfb-f59f28c2f915

C2H22

Project:	4834 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	897407.7797388354	Northing	6563338.106537405
Landform and soil		Rock	
Landform	Lower slope	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - trees,Leaf litter,Peeling bark,Woody debris
Disturbance			
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Acacia sp</i>
Ground stratum	Absent		



Fulcrum photo ID 41f4047f-24e6-432f-a252-5b9c5469bfc3

C2H24

Project:	4835 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	897534.3882230611	Northing	6563041.229803004
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Calcrete,Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Atriplex sp</i>
Ground stratum	Absent		



Fulcrum photo ID 271b8409-d185-4552-a223-f5daf7b9d31a

C2H26

Project:	4836 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	898122.0154905103	Northing	6562781.590976534
Landform and soil		Rock	
Landform	Plain	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Good		
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>Senna sp</i>
Ground stratum			



Fulcrum photo ID 1f602d5d-5404-457f-b29f-7f84c7aaf087

C2H28

Project:	4837 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	898839.090998794	Northing	6563393.36065102
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition			
Quality	Very good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	None observed	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Eremophila</i>
Ground stratum	Absent		



Fulcrum photo ID c4b932de-0fdf-423c-ad63-8302ddd73365

C2H30

Project:	4838 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	SW
Easting	898737.6863762725	Northing	6563394.817201562
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	Erosion	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Acacia</i>
Ground stratum	Absent		



Fulcrum photo ID d16a4f03-6dd4-407a-9184-f1b3cd802bc0

HABC2LC07

Project:	4839 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900704.6350980257	Northing	6565575.750708525
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Granite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Good		
Fire History	Unknown	Water Source	Absent
Disturbance	Erosion	Microhabitats	Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp.</i>



Fulcrum photo ID 51affc75-b427-4809-8ef2-d024d562bb5f,266b71e9-b0ea-422e-8f3d-

HABC2LC08

Project:	4840 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900776.5532258475	Northing	6565713.262605984
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Granite
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	None observed	Microhabitats	Leaf litter,Peeling bark,Rock crevices,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>ptilotus sp.</i>



Fulcrum photo ID	21c66f48-529d-461a-b6c1-d0ce4033c0f2,193ee8c1-7c0d-4271-ac0a-
------------------	---

HABC2LC09

Project:	4841 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900164.9336511727	Northing	6565653.958929246
Landform and soil		Rock	
Landform	Drainage line	Rock type/s	Granite,Quartz
Soil type	Sand	Surface stone cover	
Soil colour	Orange,White	Surface stone size classes present	5 - 25%
Condition			
Quality	Good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Erosion	Microhabitats	Hollows - logs,Hollows - trees,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila sp.</i>
Ground stratum	Mid (0.5-1 m)	Forbland (50-80%)	<i>eremophila sp.</i>



Fulcrum photo ID	0fca10bc-0b3f-4884-a596-1bdc49f0718d,8c8bcab2-c4f1-453a-9933-
------------------	---

HABC2LC10

Project:	4842 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900449.314056684	Northing	6564956.76156202
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Unknown	Microhabitats	Peeling bark,Woody debris
Disturbance	Clearing,Erosion,Litter		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eremophila</i> sp.
Ground stratum	Mid (0.5-1 m)	Forbland (50-80%)	<i>eremophila</i> sp.



Fulcrum photo ID 490f139c-c787-45ce-8b70-4790296ee53c,e21395e7-7464-42cf-bbb9-

HABC2LC11

Project:	4843 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	901276.6564614684	Northing	6565677.347229139
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus</i> sp.
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eremophila</i> sp.
Ground stratum	Low (>0.5 m)	Forbland (50-80%)	<i>eremophila</i> sp.



Fulcrum photo ID bbb0226c-b301-4a5f-ae00-44c3dc787f94,d6688097-4e41-4751-a79f-

HABC2LC12			
Project:	4844 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900519.9630059553	Northing	6566882.003260277
Landform and soil		Rock	
Landform	Claypan	Rock type/s	None
Soil type	Clay	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	
Condition		Habitat Features	
Quality	Disturbed	Water Source	Present
Fire History	Unknown	Microhabitats	
Disturbance	Vehicle tracks,Weeds		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Absent		
Ground stratum	Low (>0.5 m)	Grassland (50-80%)	various weeds
		Fulcrum photo ID	5e418349-31da-4044-8d04-a4990c1f72b0,450a84a8-a05b-4ee2-a0d1-





HABC2LC13			
Project:	4845 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900731.210354395	Northing	6567309.737509467
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	High quality	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Woodland (20-50%)	<i>eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Shrubland and/or heathland (50-80%)	<i>eremophila sp.</i>
Ground stratum	Mid (0.5-1 m)	Open forbland (20-50%)	<i>eremophila sp. amd other mixed herbs</i>
Fulcrum photo ID			570709d4-00c6-43d2-9de6-ef44159e1c8e,2c8bbe5f-630c-45f1-87bb-



HABC2LC14

Project:	4846 Coolgardie Biological Spring Survey		
Date	2021-11-16	Personnel	LC
Easting	900023.0356903123	Northing	6566601.201476825
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Rock crevices,Woody debris
Disturbance	Clearing,Vehicle tracks,Weeds		
Introduced fauna	Goat,Rabbit		
Vegetation			
Upper stratum	Low (<10 m)	Isolated trees (<0.25%)	<i>eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>acacia sp.</i>
Ground stratum	Low (>0.5 m)	Forbland (50-80%)	<i>eremophila sp.</i>



Fulcrum photo ID 2cc3ef18-e4b7-46ee-9c1d-0f4614438dc9,6c342fc0-f7f5-443d-9f15-

HABC2LC15

Project:	4847 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	LC
Easting	900431.2214361653	Northing	6572530.387661901
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Peeling bark,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>dodenea sp.</i>
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>ptilotus sp.</i>



Fulcrum photo ID 4199f50b-b3c4-4fd5-9f8a-3a7d7df69999,761b02e4-9a61-427a-b17e-

HABC2LC16

Project:	4848 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	LC
Easting	900465.6771902259	Northing	6573277.0945564695
Landform and soil		Rock	
Landform	Plain	Rock type/s	Calcrete,Granite
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	Cattle		
Vegetation			
Upper stratum	Low (<10 m)	Open forest (50-80%)	<i>eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Mid (0.5-1 m)	Sparse forbland (0.25-20%)	<i>eremophila sp. mixed herbs</i>



Fulcrum photo ID c208e2cf-2947-4ff4-b0f0-2556bb277948,c4d25400-28ff-4427-be00-

HABC2LC17

Project:	4849 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	LC
Easting	897973.2561752913	Northing	6573513.342421641
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Calcrete,Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open forest (50-80%)	<i>eucalyptus sp., acacia sp.</i>
Mid stratum	Low (0.5-1 m)	Shrubland and/or heathland (50-80%)	<i>dodenea sp., eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>ptilotus sp., eremophila sp.</i>



Fulcrum photo ID 372d8949-1d3b-4870-87bd-7951b013d9e4,6cf82aac-74d5-4f6d-afb6-

HABC2LC18				
Project:	4850 Coolgardie Biological Spring Survey			
Date	2021-11-17	Personnel	LC	
Easting	897208.723548352	Northing	6573995.251239745	
Landform and soil		Rock		
Landform	Plain	Rock type/s	Granite,Quartz	
Soil type	Clay loam	Surface stone cover		
Soil colour	Brown,Orange	Surface stone size classes present	5 - 25%	
Condition		Habitat Features		
Quality	Very good			
Fire History	Unknown	Water Source	Absent	
Disturbance	Litter,Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris	
Introduced fauna	None observed			
Vegetation				
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>eucalyptus</i> sp.	
Mid stratum	Tall (>2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eremophila</i> sp.	
Ground stratum	Tall (1-2 m)	Forbland (50-80%)	<i>eremophila</i> sp.	
Fulcrum photo ID			7081e49e-b513-4cfc-a25e-8219bc15b8f3,4be2e29b-ea2a-4adf-9a60-	



HABC2LC19				
Project:	4851 Coolgardie Biological Spring Survey			
Date	2021-11-17	Personnel	LC	
Easting	897024.5424583412	Northing	6573400.948208064	
Landform and soil		Rock		
Landform	Plain	Rock type/s	Calcrete,Ironstone,Laterite	
Soil type	Sandy loam	Surface stone cover		
Soil colour	Orange	Surface stone size classes present	75 - 100%	
Condition				
Quality	Good	Habitat Features		
Fire History	Unknown	Water Source	Absent	
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris	
Introduced fauna	Rabbit			
Vegetation				
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>eucalyptus</i> sp.	
Mid stratum	Low (0.5-1 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila</i> sp.	
Ground stratum	Low (>0.5 m)	Isolated forbs (<0.25%)	<i>herb in photos</i>	
			Fulcrum photo ID	4635704e-82e0-486f-bffd-6cb63a245066,db940a29-9610-40ed-a43e-



HABC2LC20			
Project:	4852 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	LC
Easting	897207.7894486763	Northing	6573276.328611162
Landform and soil		Rock	
Landform	Plain	Rock type/s	Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Hollows - trees,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eucalyptus sp.</i>
Ground stratum	Mid (0.5-1 m)	Forbland (50-80%)	<i>eremophila sp. eremophila sp.</i>



Fulcrum photo ID	b50692de-a768-4d17-bf33-bb534451e50e,1a36084b-a01d-479b-8a3d
------------------	--



Fulcrum photo ID b50692de-a768-4d17-bf33-bb534451e50e,1a36084b-a01d-479b-8a3d-

C2H32			
Project:	4853 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	898647.6288961614	Northing	6563412.670039611
Landform and soil		Rock	
Landform	Escarpment	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance	None observed	Microhabitats	Hollows - logs,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	Eucalyptus sp.
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	Acacia sp
Ground stratum	Absent		



Fulcrum photo ID	6fed3548-6854-44ad-ace1-79cee900d9bb
------------------	--------------------------------------



Fulcrum photo ID 6fed3548-6854-44ad-ace1-79cee900d9bb

C2H34

Project:	4854 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	898574.3445931688	Northing	6563567.903756286
Landform and soil		Rock	
Landform	Lower slope	Rock type/s	Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs, Leaf litter, Peeling bark, Woody debris
Disturbance			
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	Acacia
Ground stratum	Absent		



Fulcrum photo ID 5ceb13-aadb-470a-b104-5994b272e88d

C2H36

Project:	4855 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	898440.2002370819	Northing	6572518.477781318
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Granite, Laterite
Soil type	Loam	Surface stone cover	
Soil colour	Brown, Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs, Leaf litter, Peeling bark, Woody debris
Disturbance	None observed		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	Eucalyptus sp.
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	Eremophila
Ground stratum	Absent		



Fulcrum photo ID 3ee1eb2e-e0e3-4e77-9780-bb40cd650728

C2H38

Project:	4856 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	898296.4071683842	Northing	6572123.600769359
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Laterite,Quartz
Soil type	Loam	Surface stone cover	
Soil colour	Orange,Red	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Very good		
Fire History	Unknown	Water Source	Absent
Disturbance		Microhabitats	Hollows - logs,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	Acacia
Ground stratum	Absent		



Fulcrum photo ID b933e67c-48b0-421d-8100-6518118a4171

C2H40

Project:	4857 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	897937.4325779848	Northing	6572315.898469241
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Laterite,Quartz
Soil type	Loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition			
Quality	Very good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Hollows - logs,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	Acacia
Ground stratum	Absent		



Fulcrum photo ID 219563fd-c9bd-4cd0-8160-634649c52222

C2H42

Project:	4858 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	897160.8402197923	Northing	6571960.858564717
Landform and soil		Rock	
Landform	Lower slope	Rock type/s	Ironstone,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	75 - 100%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eremophila</i>
Ground stratum	Absent		



Fulcrum photo ID e4db22e5-83d5-4438-9c84-680175928daf

C2H46

Project:	4859 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	897842.0485219336	Northing	6571592.065394482
Landform and soil		Rock	
Landform	Undulating plain	Rock type/s	Ironstone,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Eremophila</i> and <i>Atriplex</i>
Ground stratum	Absent		



Fulcrum photo ID d8c7d48f-634c-4c46-a402-4949c69c1196

C2H48

Project:	4860 Coolgardie Biological Spring Survey		
Date	2021-11-17	Personnel	SW
Easting	897986.3585784157	Northing	6572856.793533593
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Granite,Ironstone,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Disturbed		
Fire History	Unknown	Water Source	Absent
Disturbance		Microhabitats	Hollows - logs,Leaf litter,Peeling bark,Woody debris
Introduced fauna			
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Acacia, eremophila, senna</i>
Ground stratum	Absent		



Fulcrum photo ID 77fa71fc-2061-4776-b46f-5533c5547e69

C2H50

Project:	4861 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	SW
Easting	899410.3384907652	Northing	6573733.02122001
Landform and soil		Rock	
Landform	Plain	Rock type/s	Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition		Habitat Features	
Quality	Disturbed		
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Woody debris
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Absent		
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>Allocasuarina, Acacia</i>
Ground stratum	Absent		



Fulcrum photo ID fc56790e-0b7a-4af8-83ad-43361892dc2a

C2H52

Project:	4862 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	SW
Easting	899024.133330778	Northing	6573326.9469294455
Landform and soil		Rock	
Landform	Claypan	Rock type/s	Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Disturbed	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	Cattle,Rabbit		
Vegetation			
Upper stratum	Absent		
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eremophila</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>Weed</i>



Fulcrum photo ID 3932baf5-4aea-4948-89e6-ec5c8ba929e9

C2H54

Project:	4863 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	SW
Easting	898558.6120445768	Northing	6577434.127282435
Landform and soil		Rock	
Landform	Undulating plain	Rock type/s	Ironstone,Laterite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Very good	Water Source	Absent
Fire History	Unknown	Microhabitats	Hollows - logs,Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	Rabbit		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>Eremophila</i>
Ground stratum	Absent		



Fulcrum photo ID 90af6e6b-2828-4926-bd8c-7ce23e0bbedf

C3H2

Project:	4864 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	SW
Easting	901330.4863598039	Northing	6566876.698387651
Landform and soil		Rock	
Landform	Upper slope	Rock type/s	Granite,Laterite
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Low (0.5-1 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Senna sp</i>
Ground stratum	Absent		



Fulcrum photo ID 9639322a-c25b-452c-9395-2d845a0db67d

C3H6

Project:	4865 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	SW
Easting	901099.8592427254	Northing	6567133.638773368
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Granite,Laterite,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Brown	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	None observed		
Introduced fauna	Cattle		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>Eremophila</i>
Ground stratum	Absent		



Fulcrum photo ID 8c9b4a09-e53a-4d9d-8d80-11e634b91457

HABC2LC21

Project:	4866 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	897544.5825756477	Northing	6572777.829306185
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Ironstone,Quartz
Soil type	Clay	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Good	Water Source	Absent
Fire History	Unknown	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Disturbance	Litter,Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Mid (10-30 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp</i>



Fulcrum photo ID 0f0d98ec-ac0a-44f0-8ad5-c4fc373482ad,537a883e-d403-430b-8f06-

HABC2LC22

Project:	4867 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	897656.784997021	Northing	6572883.799383562
Landform and soil		Rock	
Landform	Mid slope	Rock type/s	Calcrete,Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	25 - 50%
Condition		Habitat Features	
Quality	Highly degraded	Water Source	Absent
Fire History	Unknown	Microhabitats	Rock crevices,Woody debris
Disturbance	Clearing,Erosion,Vehicle tracks		
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Mid (1-2 m)	Isolated shrubs and/or heath shrubs (<0.25%)	<i>eucalyptus sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp.</i>



Fulcrum photo ID 0cc91ff4-82c9-4e1a-8a1d-4d2ee6573b54,5ea1acb4-732a-4029-ace5-

HABC2LC23

Project:	4868 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	897726.5517688439	Northing	6573587.154734112
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Very good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Absent		
Mid stratum	Mid (1-2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila</i> sp.
Ground stratum	Low (>0.5 m)	Open hummock grassland (20-50%)	<i>eremophila</i> sp.



Fulcrum photo ID af1106ec-72c2-4078-a6cf-65fdc8ed54e0,772849ce-8e9f-4ea7-8aff-



Fulcrum photo ID af1106ec-72c2-4078-a6cf-65fdc8ed54e0,772849ce-8e9f-4ea7-8aff-

HABC2LC24

Project:	4869 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	897731.7215430276	Northing	6578600.152287805
Landform and soil		Rock	
Landform	Plain	Rock type/s	Calcrete,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Disturbed	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Leaf litter,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Isolated trees (<0.25%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp.</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp.</i>



Fulcrum photo ID	7264e4af-f5e3-4399-a7c3-718a30cf91f9,043ac726-f8bb-4b81-9e64-
------------------	---



Fulcrum photo ID 7264e4af-f5e3-4399-a7c3-718a30cf91f9,043ac726-f8bb-4b81-9e64-

HABC3LC01

Project:	4870 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	901535.8496579034	Northing	6568495.428023138
Landform and soil		Rock	
Landform	Plain	Rock type/s	Granite,Ironstone,Quartz
Soil type	Clay loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	0 - 5%
Condition			
Quality	Very good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks,Weeds	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Shrubland and/or heathland (50-80%)	<i>eremophila sp.</i>
Ground stratum	Mid (0.5-1 m)	Forbland (50-80%)	<i>eremophila sp.</i>



Fulcrum photo ID 8e57e003-8ef0-4049-8a3d-f7be3316a26a,985a366e-c098-4cbd-9c83-

HABC3LC02

Project:	4871 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	901385.6159884571	Northing	6567944.72848946
Landform and soil		Rock	
Landform	Plain	Rock type/s	Calcrete,Granite,Ironstone
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Orange	Surface stone size classes present	5 - 25%
Condition		Habitat Features	
Quality	Good		
Fire History	Unknown	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	Cattle		
Vegetation			
Upper stratum	Low (<10 m)	Woodland (20-50%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>eremophila sp., sandlewood</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>eremophila sp. and mixed herbs</i>



Fulcrum photo ID 0456611e-322d-43f0-961a-92e3214f4ff6,b383b714-6500-46fb-b290-

HABC3LC03

Project:	4872 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	901116.1473216031	Northing	6567716.167221133
Landform and soil		Rock	
Landform	Undulating plain	Rock type/s	Calcrete,Granite
Soil type	Sandy loam	Surface stone cover	
Soil colour	Brown,Orange,Red	Surface stone size classes present	50 - 75%
Condition		Habitat Features	
Quality	Good		
Fire History	Unknown	Water Source	Absent
Disturbance	Litter,Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Mid (1-2 m)	Sparse shrubland and/or heathland (0.25-20%)	<i>dodenea sp, mixed shrubs</i>
Ground stratum	Low (>0.5 m)	Sparse forbland (0.25-20%)	<i>mixed herbs</i>



Fulcrum photo ID a233588d-1aef-4741-8ae9-5b0d1c66fa0a,42b4783a-a56b-40d7-8240-

HABC3LC04

Project:	4873 Coolgardie Biological Spring Survey		
Date	2021-11-18	Personnel	LC
Easting	900730.2686670613	Northing	6567991.079386459
Landform and soil		Rock	
Landform	Plain	Rock type/s	Calcrete,Granite
Soil type	Sandy loam	Surface stone cover	
Soil colour	Orange	Surface stone size classes present	25 - 50%
Condition			
Quality	Very good	Habitat Features	
Fire History	Unknown	Water Source	Absent
Disturbance	Vehicle tracks	Microhabitats	Leaf litter,Logs > 10 cm,Peeling bark,Woody debris
Introduced fauna	None observed		
Vegetation			
Upper stratum	Low (<10 m)	Open woodland (0.25-20%)	<i>Eucalyptus sp.</i>
Mid stratum	Tall (>2 m)	Open shrubland and/or heathland (20-50%)	<i>eremophila sp.</i>
Ground stratum	Mid (0.5-1 m)	Sparse forbland (0.25-20%)	<i>eremophila sp., mixed herbs</i>



Fulcrum photo ID 20c229a2-37df-4ff5-9e14-6fa99181557f,a6d138ee-2032-4f92-93bc-

Appendix G

Fauna Inventory

Conservation Status: State - Listed under Biodiversity Conservation Act 2016 or Department of Biodiversity, Conservation and Attractions Conservation, Federal - Listed under Environmental Protection and Biodiversity Conservation Act 1999. VU - Vulnerable, MA - Marine. * - Introduced species.

Family	Scientific Name	Common Name	Conservation Status		Method						
			State	Federal	Sighting	Call	Remains	Scat	Tracks	Burrow	Digging
Aves											
Acanthizidae	Smicrornis brevirostris	Weebill				2					
	Acanthiza inornata	Western Thornbill				2					
	Aphelocephala leucopsis	Southern Whiteface				1					
	Calamanthus cautus	Shy Groundwren (Shy Heathwren)				1					
	Acanthiza uropygialis	Chestnut-rumped Thornbill			2	1					
	Sericornis frontalis					1					
Artamidae	Artamus personatus	Masked Woodswallow			2						
Cacatuidae	Cacatua roseicapilla	Galah				1					
	Cacatua leadbeateri	Major Mitchell's Cockatoo				1					
Campephagidae	Coracina novaehollandiae	Black-faced Cuckoo-shrike		MA	2			1			
	Coracina maxima	Ground Cuckoo-shrike				1					
	Lalage tricolor	White-winged Triller				1					
Climacteridae	Climacteris rufus	Rufous Treecreeper				1					
Columbidae	Phaps chalcoptera	Common Bronzewing			1						
	Ocyphaps lophotes	Crested Pigeon			3						
Corvidae	Corvus bennetti	Little Crow				5					
	Corvus orru ceciliae	Western Crow				1					
	Corvus coronoides	Australian Raven			2						
Cracticidae	Strepera versicolor	Grey Currawong			4	2					
Dromaiidae	Dromaius novaehollandiae	Emu						1	2		
Estrildidae	Taeniopygia guttata	Zebra Finch				2					
Falconidae	Falco cenchroides	Australian Kestrel (Nankeen Kestrel)		MA	3						
Maluridae	Malurus splendens	Splendid Fairywren				1					
Meliphagidae	Anthochaera carunculata	Red Wattlebird			1	3					
	Lichmera indistincta	Brown Honeyeater				1					
	Gavicalis virescens	Singing Honeyeater			13	2					

Family	Scientific Name	Common Name	Conservation Status		Method						
			State	Federal	Sighting	Call	Remains	Scat	Tracks	Burrow	Digging
	<i>Manorina flavigula</i>	Yellow-throated Miner				4					
	<i>Acanthagenys rufogularis</i>	Spiny-cheeked Honeyeater			7						
	<i>Melithreptus brevirostris</i>	Brown-headed Honeyeater				1					
	<i>Gliciphila melanops</i>	Tawny-crowned Honeyeater			2	1					
Meropidae	<i>Merops ornatus</i>	Rainbow Bee-eater		MA	1						
Monarchidae	<i>Grallina cyanoleuca</i>	Magpie-lark		MA	2						
Motacillidae	<i>Anthus australis</i>	Australian Pipit			4						
Oreoicidae	<i>Oreoica gutturalis</i>	Crested Bellbird			1	11					
Otididae	<i>Ardeotis australis</i>	Australian Bustard							1		
Pachycephalidae	<i>Pachycephala rufiventris</i>	Rufous Whistler			5	8					
	<i>Pachycephala occidentalis</i>	Western Golden Whistler (Western Whistler)			6						
Petroicidae	<i>Microeca fascians</i>	Jacky Winter			1	1					
Psittacidae	<i>Melopsittacus undulatus</i>	Budgerigar			23						
	<i>Platycercus zonarius</i>	Australian Ringneck			17	2					
Psophodidae	<i>Cinclosoma clarum</i>	Western Chestnut Quail-thrush (Copperback Quail-thrush)			2						
Rhipiduridae	<i>Rhipidura leucophrys</i>	Willie Wagtail			2						
Mammalia											
Bovidae	<i>*Capra hircus</i>	Goat						8	1		
	<i>*Bos taurus</i>	European Cattle						9	7		
Canidae	<i>*Canis familiaris familiaris</i>	Dog						2			
	<i>*Vulpes vulpes</i>	Red Fox						1			
Dasyuridae	<i>Dasyurus geoffroii fortis</i>	Western Quoll, Chuditch	VU	VU				1			
Equidae	<i>*Equus caballus</i>	Horse						1			
Felidae	<i>*Felis catus</i>	Cat					1				
Leporidae	<i>*Oryctolagus cuniculus</i>	Rabbit			1		3	18		1	2
Macropodidae	<i>Macropus fuliginosus melanops</i>	Western Grey Kangaroo			1			39	2		
	<i>Osphranter rufus</i>	Red Kangaroo, Marlu			2						
Reptilia											
Agamidae	<i>Ctenophorus maculatus</i>				2						

Family	Scientific Name	Common Name	Conservation Status		Method						
			State	Federal	Sighting	Call	Remains	Scat	Tracks	Burrow	Digging
	<i>Ctenophorus cristatus</i>	Bicycle Dragon			6						
	<i>Tympanocryptis pseudopsephos</i>	Goldfields pebble-mimic dragons			2						
	<i>Ctenophorus ornatus</i>	Ornate Crevice Dragon			1						
Diplodactylidae	<i>Crenadactylus ocellatus</i>	South-western Clawless Gecko			3						
Gekkonidae	<i>Heteronotia binoei</i>	Bynoe's Gecko			8						
	<i>Gehyra variegata</i>	Variegated gehyra			2						
Scincidae	<i>Tiliqua rugosa</i>	Bobtail			8		2				
Varanidae	<i>Varanus gouldii</i>	Bungarra or Sand Goanna			1						



Part of
SLR 

10 Bermondsey Street West Leederville WA 6007 **t** (+618) 9388 8360 **f** (+618) 9381 2360
PO BOX 14, West Perth WA 6872
w 360environmental.com.au **e** admin@360environmental.com.au

● people ● planet ● professional